



US009315491B2

(12) **United States Patent**
Kuo et al.

(10) **Patent No.:** **US 9,315,491 B2**
(45) **Date of Patent:** **Apr. 19, 2016**

(54) **PYRIMIDINE COMPOUNDS AS MTOR AND PI3K INHIBITORS**

(71) Applicants: **DEVELOPMENT CENTER FOR BIOTECHNOLOGY**, New Taipei (TW); **DCB-USA LLC**, Wilmington, DE (US)

(72) Inventors: **Mann-Yan Kuo**, New Taipei (TW); **Ying-Shuan Lee**, New Taipei (TW); **Paonien Chen**, New Taipei (TW); **Li Jung Chen**, New Taipei (TW); **Yann Yu Lu**, New Taipei (TW); **Yi-Ting Huang**, New Taipei (TW); **Hung-Yi Hsu**, New Taipei (TW); **Ping-Kuei Tsai**, New Taipei (TW)

(73) Assignees: **DEVELOPMENT CENTER FOR BIOTECHNOLOGY**, New Taipei (TW); **DCB-USA LLC**, New Castle, DE (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/192,301**

(22) Filed: **Feb. 27, 2014**

(65) **Prior Publication Data**

US 2014/0178360 A1 Jun. 26, 2014

Related U.S. Application Data

(63) Continuation-in-part of application No. 13/519,535, filed as application No. PCT/IB2010/003347 on Dec. 27, 2010, now abandoned.

(60) Provisional application No. 61/290,437, filed on Dec. 28, 2009.

(51) **Int. Cl.**

A61K 31/5377 (2006.01)
C07D 239/47 (2006.01)
A61K 31/506 (2006.01)
C07D 401/04 (2006.01)
C07D 401/14 (2006.01)
C07D 413/04 (2006.01)
A61K 31/541 (2006.01)
A61K 45/06 (2006.01)
C07D 413/14 (2006.01)
C07D 417/14 (2006.01)
C07D 403/12 (2006.01)
C07D 409/12 (2006.01)
C07D 413/12 (2006.01)

(52) **U.S. Cl.**

CPC **C07D 413/04** (2013.01); **A61K 31/5377** (2013.01); **A61K 31/541** (2013.01); **A61K 45/06** (2013.01); **C07D 239/47** (2013.01); **C07D 401/04** (2013.01); **C07D 401/14** (2013.01); **C07D 403/12** (2013.01); **C07D 409/12** (2013.01); **C07D 413/12** (2013.01); **C07D 413/14** (2013.01); **C07D 417/14** (2013.01)

(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,565,864	A	1/1986	Wade
5,250,530	A *	10/1993	Giencke et al. 514/256
5,420,129	A	5/1995	Breu et al.
7,423,148	B2	9/2008	Nuss et al.
2008/0280889	A1	11/2008	Bilodeau et al.
2009/0156601	A1	6/2009	McDonald et al.

FOREIGN PATENT DOCUMENTS

WO	00/43385	7/2000
WO	01/81338	A1 11/2001
WO	2007/066099	A1 6/2007
WO	2008/023159	A1 2/2008
WO	2008/032086	A1 3/2008
WO	2008/098058	A1 8/2008
WO	2009/019656	A1 2/2009

OTHER PUBLICATIONS

Communication dated Dec. 10, 2014, issued by the Intellectual Property Office of Singapore in corresponding Singapore Application No. 2012044517.

John Hannah et al., "Carba-acyclonucleoside Antiherpetic Agents," J. of Heterocyclic Chemistry, 1989, pp. 1261-1271.

John A. Montgomery et al., "Analogues of 5'-Deoxy-5'-(methylthio)adenosine," J. Med. Chem., 1974, vol. 17, No. 11, pp. 1197-1207.

Michael J. Berridge et al., "Inositol trisphosphate, a novel second messenger in cellular signal transduction," Nature, vol. 312, No. 22, Nov. 1984, pp. 315-321.

Felix Kopp et al., "Functionalization of Unprotected Uracil Derivatives using the Halogen-Magnesium Exchange," Organic Letters, 2007, vol. 9, No. 9, pp. 1639-1641.


(Continued)

Primary Examiner — Samantha Shterengarts

(74) *Attorney, Agent, or Firm* — Sughrue Mion, PLLC

(57) **ABSTRACT**

The present invention relates to pyrimidine compounds of formula (I):



(I)

which are useful in treating mTOR kinase- or PI3K kinase-related diseases.

10 Claims, No Drawings

(56)

References Cited

OTHER PUBLICATIONS

Yasutomi Nishizuka, "Turnover of Inositol Phospholipids and Signal Transduction," Science, vol. 225, No. 21, Sep. 1984, pp. 1365-1370.
International Search Report of PCT/IB10/03347, dated Jun. 14, 2011.

Extended European Search Report dated Apr. 25, 2013 in European Application No. 10840651.3.

S B Gashev et al: "Aminomethylation of substituted 5-hydroxypyrimidines", Chemistry of Heterocyclic Compounds, Jan. 1, 1980, pp. 532-534.

Tilman Lechel et al, "A Concise Synthesis of Alkoxy-Substituted Pyrimidine Derivatives Based upon a Three-Component Access to Functionalized Enamides", Synlett, 2009, No. 07, Apr. 1, 2009, pp. 1059-1062.

Reinhold Zimmer et al, "Preparation of Pyrimidine-N-oxides by Condensation of Functionalized Enamides with Hydroxylamine Hydrochloride", Synlett, 2010, No. 12, Jul. 1, 2010, pp. 1793-1796.

Tilman Lechel et al, "New 5-Alkoxy-pyrimidine Derivatives from [beta]-Alkoxy [beta]-Keto Enamides and Ammonium Salts", European Journal of Organic Chemistry, 2010, No. 13, May 1, 2010, pp. 2555-2564.

Office Action issued on Apr. 24, 2014 from the European Patent Office in European Application No. 10 840 651.3.

* cited by examiner

1

PYRIMIDINE COMPOUNDS AS MTOR AND PI3K INHIBITORS

CROSS REFERENCE TO RELATED APPLICATIONS

This application is a continuation-in-part of U.S. application Ser. No. 13/519,535, filed Jun. 27, 2012, which is a 371 of PCT/IB2010/003347 filed Dec. 27, 2010, which claims priority to U.S. Provisional Application No. 61/290,437, filed Dec. 28, 2009; the contents of all of which are incorporated herein by reference in their entirety.

FIELD OF THE INVENTION

The present invention relates to novel pyrimidine compounds and their use in treating PI3K kinase- and/or mTOR kinase-related diseases.

BACKGROUND OF THE INVENTION

The mammalian target of Rapamycin, mTOR, is a cell-signaling protein that regulates the response of tumor cells to nutrients and growth factors, as well as controlling tumor blood supply through effects on Vascular Endothelial Growth Factor, VEGF. Inhibitors of mTOR starve cancer cells and shrink tumors by inhibiting the effect of mTOR. There are two important effects as mTOR inhibitors bind to the mTOR kinase. First, mTOR is a downstream mediator of the PI3K/Akt pathway. The PI3K/Akt pathway is thought to be over-activated in numerous cancers and may account for the widespread response from various cancers to mTOR inhibitors. The over-activation of the upstream pathway would normally cause mTOR kinase to be over-activated as well. However, in the presence of mTOR inhibitors, this process is blocked. The blocking effect prevents mTOR from signaling to downstream pathways that control cell growth. Over-activation of the PI3K/Akt kinase pathway is frequently associated with mutations in the PTEN gene, which is common in many cancers and may help predict what tumors will respond to mTOR inhibitors. The second major effect of mTOR inhibition is antiangiogenesis via the lowering of VEGF levels. These anticancer drugs have shown exceptional promise in cancer therapy and may change the way many types of cancer are treated.

Several studies have demonstrated that mTOR has a central role in controlling cell growth, proliferation and metabolism. mTOR regulates a wide range of cellular functions, including translation, transcription, mRNA turnover, protein stability, actin cytoskeletal organization and autophagy. mTOR is a member of the phosphoinositide kinase-related kinase (PIKK) family, but is not a phosphorylating phosphoinositide, a phosphorylate protein on serine or a threonine residue. There are two mTOR complexes in mammalian cells. mTOR complex I (mTORC1) is a raptor-mTOR complex, which mainly regulates cell growth in a rapamycin-sensitive manner, whereas mTOR complex II (mTORC2) is a rictor-mTOR complex, which regulates cytoskeletal organization in a rapamycin-insensitive manner.

Kinase subunits of both mTORC1 and mTORC2 regulate cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino-acids. Amino-acid-signaling to mTORC1 is mediated by Rag GTPases, which cause amino-acid-induced relocalization of mTOR within the endomembrane system. Growth-factor-stimulated mTORC1 activation involves AKT1-mediated phosphorylation of TSC1-TSC2, which leads to the

2

activation of the Rheb GTPase that potentially activates the protein kinase activity of mTORC1. Activated mTORC1 up-regulates protein synthesis by phosphorylating key regulators of mRNA translation and ribosome synthesis. mTORC1 phosphorylates eIF4EBP1 and releases it from inhibiting the elongation initiation factor 4E (eIF4E). mTORC1 phosphorylates and activates S6K1 at Thr-421, which then promotes protein synthesis by phosphorylating PDCD4 and targeting it for degradation. mTORC2 is also activated by growth factors, but seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTORC2 plays a critical role in AKT1 Ser-473 phosphorylation, which may facilitate the phosphorylation of the activation loop of AKT1 on Thr-308 by PDK1, which is a prerequisite for full activation. mTORC2 regulates the phosphorylation of SGK1 at Ser-422. mTORC2 also modulates the phosphorylation of PRKCA on Ser-657.

With the recent discovery of rapamycin independent function of mTOR (by mTOR2) in phosphorylation AKT (at S473), which is important in regulation of cell survival and modulation of PKC α , which plays a major role in regulation of actin cytoskeletal organization, it is believed that inhibition of mTOR function by rapamycin is partial. Therefore, a small molecule designed to compete with ATP in the catalytic site of mTOR would be expected to inhibit all of the kinase-dependent functions of mTORC1 and mTORC2, unlike rapalogs that only target mTORC1. Here we describe the discovery of direct mTOR kinase inhibitors which can be used in the treatment of a variety of cancers, including breast, lung, kidney, prostate, blood, liver, and ovarian cancers, and lymphoma and other indications such as rheumatoid arthritis, hamartoma syndromes, transplant rejection, multiple sclerosis and immunosuppression.

Phosphatidylinositol (hereinafter abbreviated as "PI") is one of the phospholipids in cell membranes. In recent years it has become clear that PI also plays an important role in intracellular signal transduction. In particular, it is well recognized in the art that PI (4,5)bisphosphate (PI(4,5)P₂) is degraded into diacylglycerol and inositol(1,4,5)triphosphate by phospholipase C to induce activation of protein kinase C and intracellular calcium mobilization, respectively [M. J. Berridge et al., *Nature*, 312, 315 (1984); Y. Nishizuka, *Science*, 225, 1365 (1984)].

PI3K was originally considered to be a single enzyme, but it has now been clarified that a plurality of subtypes is present in PI3K. Each subtype has its own mechanism for regulating activity. The PI3K family comprises at least 15 different enzymes sub-classified by structural homology and divided into 3 classes based on sequence homology and the product formed by enzyme catalysis. The class I PI3 kinases are composed of 2 subunits: a 110 kd catalytic subunit and an 85 kd regulatory subunit. The regulatory subunits contain SH2 domains and bind to tyrosine residues phosphorylated by growth factor receptors with a tyrosine kinase activity or oncogene products, thereby inducing the PI3K activity of the p110 α catalytic subunit which phosphorylates its lipid substrate. Class I PI3Ks are further divided into two groups, class Ia and class Ib, in terms of their activation mechanism. Class Ia PI3Ks include PI3K p110 α , p110 β and p110 δ subtypes, which transmit signals from tyrosine kinase-coupled receptors. Class Ib PI3K includes a p110 γ subtype activated by a G protein-coupled receptor. PI and PI(4)P are known as substrates for class II PI3Ks. Class II PI3Ks include PI3K C2 α ,


3

C2 β and C2 γ subtypes, which are characterized by containing C2 domains at the C terminus. The substrate for class III PI3Ks is PI only.

Most if not all of the non-rapalog mTOR inhibitors described to date in the scientific literature were developed to inhibit other enzymes, especially class I PI3Ks. Because PI3K regulates mTOR activity, inhibitors that target both enzymes are generally not useful as research tools to study mTOR regulation or function. However, drugs that are dual PI3K/mTOR inhibitors might have a therapeutic advantage over single-target inhibitors in certain disease settings. PI3K inhibitors and mTOR inhibitors are expected to be novel types of medicaments useful against cell proliferation disorders, especially as carcinostatic agents. Thus, it would be advantageous to have new mTOR inhibitors and PI3K inhibitors as potential treatment regimens for mTOR kinase- and PI3K kinase-related diseases.

SUMMARY OF THE INVENTION


It has now been found that a series of novel pyrimidine compounds have activity as inhibitors of mTOR and PI3K. Accordingly, the present invention provides a compound of formula (I):



wherein

R¹ is selected from:

(i) a group of the following formula:



wherein

P is aryl or heteroaryl which is unsubstituted or substituted, wherein when P is a benzoimidazole group, said benzoimidazole group is attached to the pyrimidinyl group in formula (I) through its benzene ring;

Q is selected from —H, —OR, —SR, —Halo, —NR₃R₄, —OS(O)_mR, —OC(O)R, —OC(O)NHR, —S(O)_mNR₃R₄, —NRC(O)R, —NRS(O)_mR, —NRC(O)NR₃R₄, and —NRC(S)NR₃R₄, wherein each R, R₃, and R₄ is independently selected from H, C₁-C₆ alkyl, C₃-C₁₀ cycloalkyl and a 5- to 12-membered carbocyclic group, aryl or heteroaryl group, the group being unsubstituted or substituted; m is 1 or 2; or R₃ and R₄, which are the same or different, are each independently selected from H, C₁-C₆ alkyl which is unsubstituted or substituted, C₃-C₁₀ cycloalkyl which is unsubstituted or substituted, —C(O)R, —C(O)N(R)₂ and —S(O)_mR wherein R and m are as defined above, or R₃ and R₄ together with the nitrogen atom to which they are attached form a saturated 5-, 6- or 7-membered N-containing heterocyclic group which is unsubstituted or substituted; —C(O)R, —C(O)N(R)₂ and —S(O)_mR wherein R and m are as defined above;

Y is selected from —O—(CH₂)_n—, —S—(CH₂)_n—, and —S(O)_m(CH₂)_n— wherein m is 1 or 2, n is 0 or an integer of 1 to 3, and R² is selected from H or a 5- to 12-membered

4

carbocyclic or heterocyclic group which is unsubstituted or substituted, and a group —NR₃R₄ wherein R₃ and R₄ are as defined above;

Z is selected from (i) halo, —(CH₂)_s, —COOR, —(CH₂)_sCHO, —(CH₂)_sCH₂OR, —(CH₂)_sCONR₃R₄, —(CH₂)_sCH₂NR₃R₄, —NR₃R₄ and —O(CH₂)_sNR₃R₄ wherein s is 0 or an integer of 1 to 2 and wherein R, R₃ and R₄ are as defined above; (ii) substituted or unsubstituted heteroaryl, (iii) substituted or unsubstituted heterocyclyl, (iv) substituted or unsubstituted aryl, and (v) substituted or unsubstituted C₁-C₆-alkyl; and W is selected from (i) NR₅R₆, wherein R₅ and R₆ form, together with the N atom to which they are attached, a morpholine ring which is unsubstituted or substituted, (ii) substituted or unsubstituted heteroaryl, (iii) substituted or unsubstituted heterocyclyl, (iv) substituted or unsubstituted aryl, and (v) substituted or unsubstituted C₁-C₆-alkyl; or a stereoisomer, or a tautomer, or an N-oxide, or a pharmaceutically acceptable salt, or an ester, or a pro-drug, or a hydrate, or a solvate thereof.


Provided herein are pharmaceutical compositions comprising the compound of formula (I), or a stereoisomer, or a tautomer, or an N-oxide, or a pharmaceutically acceptable salt, or an ester, or a prodrug, or a hydrate, or a solvate thereof, or a pharmaceutically acceptable carrier or diluent.

Further provided herein is a method for treating mTOR kinase-/PI3K kinase-related diseases, which comprises administering to a subject an effective amount of the compound of formula (I), or a stereoisomer, or a tautomer, or an N-oxide, or a pharmaceutically acceptable salt, or an ester, or a prodrug, or a hydrate, or a solvate thereof.

Additionally provided herein is a method for preparing the compound of formula (I), or a stereoisomer, or a tautomer, or an N-oxide, or a pharmaceutically acceptable salt, or an ester, or a prodrug, or a hydrate, or a solvate thereof.

DETAILED DESCRIPTION OF THE INVENTION


The present invention relates to a compound of formula (I):



wherein

R¹ is selected from:

(i) a group of the following formula:



wherein

P is aryl or heteroaryl which is unsubstituted or substituted, wherein when P is a benzoimidazole group, said benzoimidazole group is attached to the pyrimidinyl group in formula (I) through its benzene ring;

Q is selected from —H, —OR, —SR, —Halo, —NR₃R₄, —OS(O)_mR, —OC(O)R, —OC(O)NHR, —S(O)_mNR₃R₄, —NRC(O)R, —NRS(O)_mR, —NRC(O)NR₃R₄, and —NRC(S)NR₃R₄, wherein each R, R₃, and R₄ is independently selected from H, C₁-C₆ alkyl, C₃-C₁₀ cycloalkyl and a

5


5- to 12-membered carbocyclic group, aryl or heteroaryl group, the group being unsubstituted or substituted; m is 1 or 2; or R₃ and R₄, which are the same or different, are each independently selected from H, C₁-C₆ alkyl which is unsubstituted or substituted, C₃-C₁₀ cycloalkyl which is unsubstituted or substituted, —C(O)R, —C(O)N(R)₂ and —S(O)_mR wherein R and m are as defined above, or R₃ and R₄ together with the nitrogen atom to which they are attached form a saturated 5-, 6- or 7-membered N-containing heterocyclic group which is unsubstituted or substituted; —C(O)R, —C(O)N(R)₂ and —S(O)_mR wherein R and m are as defined above;

Y is selected from —O—(CH₂)_n—, —S—(CH₂)_n—, and —S(O)_m(CH₂)_n— wherein m is 1 or 2, n is 0 or an integer of 1 to 3, and R² is selected from H or a 5- to 12-membered carbocyclic or heterocyclic group which is unsubstituted or substituted, and a group —NR₃R₄ wherein R₃ and R₄ are as defined above;

Z is selected from (i) halo, —(CH₂)_s—COOR, —(CH₂)_sCHO, —(CH₂)_sCH₂OR, —(CH₂)_sCONR₃R₄, —(CH₂)_sCH₂NR₃R₄, —NR₃R₄ and —O(CH₂)_sNR₃R₄ wherein s is 0 or an integer of 1 to 2 and wherein R, R₃ and R₄ are as defined above; (ii) substituted or unsubstituted heteroaryl, (iii) substituted or unsubstituted heterocyclyl, (iv) substituted or unsubstituted aryl, and (v) substituted or unsubstituted C₁-C₆-alkyl; and W is selected from (i) NR₅R₆, wherein R₅ and R₆ form, together with the N atom to which they are attached, a morpholine ring which is unsubstituted or substituted, (ii) substituted or unsubstituted heteroaryl, (iii) substituted or unsubstituted heterocyclyl, (iv) substituted or unsubstituted aryl, and (v) substituted or unsubstituted C₁-C₆-alkyl; or a stereoisomer, or a tautomer, or an N-oxide, or a pharmaceutically acceptable salt, or an ester, or a pro-drug, or a hydrate, or a solvate thereof.

Preferably, P is a heteroaryl group selected from:

- (a) an indazole group which is unsubstituted or substituted;
- (b) an indole group which is unsubstituted or substituted; and
- (c) a benzimidazole group of the formula



which is unsubstituted or substituted.

A C₁-C₆ alkyl group is linear or branched. Preferably, the alkyl is a C₁-C₄ alkyl group. A C₁-C₆ alkyl group can be unsubstituted or substituted with one or more groups Z as defined above. Examples of a C₁-C₆ alkyl group include, but are not limited to, methyl, ethyl, propyl (including all isomeric forms), n-propyl, isopropyl, butyl (including all isomeric forms), n-butyl, isobutyl, sec-butyl, t-butyl, pentyl (including all isomeric forms), and hexyl (including all isomeric forms).

A halogen is F, Cl, Br or I. Preferably, it is F, Cl or Br. A C₁-C₆ alkyl group substituted by halogen may be denoted by the term “halo-C₁-C₆ alkyl,” which means an alkyl group in which one or more hydrogen atoms are replaced by halo. A halo-C₁-C₆ alkyl group preferably contains one, two or three halo groups. A preferred halo-C₁-C₆ alkyl group is trifluoromethyl.

A C₃-C₁₀ cycloalkyl group may be saturated or unsaturated but a non-aromatic, and/or bridged, and/or non-bridged, and/or fused bicyclic group having 3 to 10 carbon atoms. The

6

cycloalkyl group preferably has 3 to 8, more preferably has 3 to 6 carbon atoms. Examples of a C₃-C₁₀ cycloalkyl group include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cyclohexadienyl, cycloheptyl, and cycloheptenyl. A C₃-C₁₀ cycloalkyl group can be unsubstituted or substituted with one or more groups Z as defined above.

An unsaturated 5- to 12-membered carbocyclic group is a 5-, 6-, 7-, 8-, 9-, 10-, 11- or 12-membered carbocyclic ring containing at least one unsaturated bond. It is a monocyclic or fused bicyclic ring system. The group is aromatic or non-aromatic, for instance a 5- to 12-membered aryl group. Examples of an unsaturated 5- to 12-membered carbocyclic group include, but are not limited to, phenyl, naphthyl, indanyl, indenyl and tetrahydronaphthyl groups. The group is unsubstituted or substituted with one or more groups Z as defined above.

An aryl group refers to a 5- to 12-membered, monovalent monocyclic aromatic group and/or monovalent polycyclic aromatic group that contains at least one aromatic carbon ring. Preferably, it is monocyclic or bicyclic. Examples of aryl groups include, but are not limited to, phenyl, naphthyl, fluorenyl, azulenyl, anthryl, phenanthryl, pyrenyl, biphenyl, and terphenyl. The group is unsubstituted or substituted with a group Z as defined above.

A saturated 5-, 6-, or 7-membered N-containing heterocyclic ring refers to a monovalent monocyclic non-aromatic ring system or monovalent polycyclic ring system that contains at least one N and the remaining ring atoms are carbon atoms. Examples of such heterocyclic ring include, but are not limited to, piperidine, piperazine, morpholine or pyrrolidine. The ring typically contains one nitrogen atom and either an additional N atom or an O atom, or no additional heteroatoms. The ring is unsubstituted or substituted on one or more ring carbon atoms and/or on any additional N atom present in the ring. Examples of suitable substituents include one or more groups Z as defined above and a C₁-C₆ alkyl group. When the ring is piperazine, it is typically unsubstituted or substituted, typically on the second ring nitrogen atom, by —C(O)R, —C(O)N(R)₂ or —S(O)_mR, or by C₁-C₆ alkyl which is unsubstituted or substituted by C₁-C₆ alkoxy or OH.

An unsaturated 5- to 12-membered heterocyclic group is typically heteroaryl. Heteroaryl can be a 5- to 12-membered aromatic group having 1, 2, 3, or 4 heteroatoms selected from O, N and S. Typically it contains one N atom and 0, 1, 2 or 3 additional heteroatoms selected from 0, S and N. It may be, for example, furan, thiophene, pyrrole, indole, isoindole, pyrazole, imidazole, benzothiophene, benzothiazole, benzofuran, isoxazole, oxazole, oxadiazole, thiazole, isothiazole, thiadiazole, dihydroimidazole, pyridine, pyridine, quinoline, isoquinoline, quinoxaline, thienopyrazine, pyran, pyrimidine, pyridazine, pyrazine, triazine, triazole or tetrazole. The group can be unsubstituted or substituted with one or more groups Z as defined above.

Specific examples of the compounds of the invention include:

- 5-Ethoxy-2-(3-hydroxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0037)
- 5-Ethoxy-2-(4-hydroxy-3-methoxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0038)
- 5-Ethoxy-2-(3-fluoro-4-methoxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0039)
- 2-(3-Amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0040)
- 3-[4-(2-Dimethylamino-ethoxy)-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl]-phenol (MTR-0043)

3-[4-(2-Dimethylamino-ethoxy)-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl]-phenol (MTR-0046)
 3-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0047)
 2-(3,5-Difluoro-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0053)
 5-Ethoxy-2-(1H-indol-5-yl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0054)
 5-Ethoxy-2-(1H-indol-6-yl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0055)
 2-Benzo[1,3]dioxol-5-yl-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0056)
 5-Ethoxy-2-(1H-indazol-4-yl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0057)
 5-Ethoxy-2-(2-fluoro-3-methoxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0058)
 2-{3-[3-(4-Chloro-3-trifluoromethyl-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0059)
 5-Ethoxy-6-morpholin-4-yl-2-[3-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0060)
 2-(4-Amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0068)
 5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0069)
 2-{4-[3-(4-Chloro-3-trifluoromethyl-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0070)
 4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0071)
 2-(3-Hydroxy-phenyl)-5-methoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0073)
 5-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-1H-indole (MTR-0074)
 2-(1H-Indazol-4-yl)-5-methoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0075)
 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-chloro-3-trifluoromethyl-phenyl)-urea (MTR-0076)
 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(6-chloro-pyridin-3-yl)-urea (MTR-0077)
 N-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-acetamide (MTR-0078)
 2-{4-[3-(6-Chloro-pyridin-3-yl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0079)
 5-Ethoxy-2-[4-(3-ethyl-ureido)-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0081)
 [4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0083)
 4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-1H-indazole (MTR-0084)
 Methanesulfonic acid 3-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester (MTR-0086)
 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea (MTR-0091)
 3-(4-Chloro-5-methanesulfonyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenol (MTR-0094)
 N-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-methane sulfonamide (MTR-0096)
 4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenol (MTR-0098) Methanesulfonic acid 4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester (MTR-0100)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-thioureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0102)
 2-(4-Benzenesulfonylamino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0103)
 1-(2-Chloro-ethyl)-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0108)
 5-Ethoxy-2-(4-hydroxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0111)
 5-Ethoxy-2-(4-ethylcarbamoyloxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0112)
 5-Ethoxy-6-morpholin-4-yl-2-(4-phenylcarbamoyloxy-phenyl)-pyrimidine-4-carboxylic acid ethyl ester (MTR-0113)
 (2-Chloro-ethyl)-carbamic acid 4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester (MTR-0114)
 Benzenesulfonic acid 4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester (MTR-0115)
 3-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenol (MTR-0116)
 5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid (MTR-0118)
 N-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-benzenesulfonamide (MTR-0119)
 5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid diethylamide (MTR-0120)
 5-Ethoxy-6-morpholin-4-yl-2-(4-ureido-phenyl)-pyrimidine-4-carboxylic acid ethyl ester (MTR-0121)
 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-dimethylamino-phenyl)-urea (MTR-0123)
 1-{4-[5-Ethoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-phenyl-urea (MTR-0124)
 5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid amide (MTR-0125)
 Phenyl-carbamic acid 4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester (MTR-0127)
 1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea (MTR-0128)
 1-Ethyl-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0129)
 Ethyl-carbamic acid 4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester (MTR-0130)
 4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-benzenesulfonamide (MTR-0131)
 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-ethyl-urea (MTR-0132)
 5-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-pyridin-2-ylamine (MTR-0133)
 1-[5-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-pyridin-2-yl]-3-phenyl-urea (MTR-0134)
 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-fluoro-phenyl)-urea (MTR-0135)
 1-[5-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-pyridin-2-yl]-3-ethyl-urea (MTR-0136)
 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3-fluoro-phenyl)-urea (MTR-0137)
 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3,4-difluoro-phenyl)-urea (MTR-0138)
 5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid (2-morpholin-4-yl-ethyl)-amide (MTR-0139)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid (2-pyrrolidin-1-yl-ethyl)-amide (MTR-0140)

1-{4-[5-Ethoxy-4-morpholin-4-yl-6-(pyrrolidine-1-carbonyl)-pyrimidin-2-yl]-phenyl}-3-phenyl-urea (MTR-0141)

1-{4-[5-Ethoxy-4-morpholin-4-yl-6-(piperidine-1-carbonyl)-pyrimidin-2-yl]-phenyl}-3-phenyl-urea (MTR-0142)

1-[5-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-pyridin-2-yl]-3-(3-fluoro-phenyl)-urea (MTR-0143)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid (2-diethylamino-ethyl)-amide (MTR-0144)

1-{4-[5-Ethoxy-4-(4-methanesulfonyl-piperazine-1-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-phenyl-urea (MTR-0145)

2-(6-Amino-pyridin-3-yl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0146)

5-Ethoxy-2-{4-[3-(3-fluoro-phenyl)-ureido]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0147)

5-Ethoxy-6-morpholin-4-yl-2-[6-(3-phenyl-ureido)-pyridin-3-yl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0148)

5-Ethoxy-2-{4-[3-(4-fluoro-phenyl)-ureido]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0149)

2-{4-[3-(3,4-Difluoro-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0150)

1-(4-Fluoro-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0152)

4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0153)

4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0154)

4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenol (MTR-0155)

5-Ethoxy-2-{6-[3-(4-fluoro-phenyl)-ureido]-pyridin-3-yl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0156)

2-{6-[3-(3,4-Difluoro-phenyl)-ureido]-pyridin-3-yl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0157)

5-Ethoxy-2-{6-[3-(3-fluoro-phenyl)-ureido]-pyridin-3-yl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0158)

1-Ethyl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0159)

4-[5-Ethoxy-4-(4-methanesulfonyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-1H-indazole (MTR-0160)

4-[5-Ethoxy-4-(4-methanesulfonyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenylamine (MTR-0161)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea (MTR-0162)

4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-benzenesulfonamide (MTR-0163)

1-(3-Fluoro-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0164)

1-(3,4-Difluoro-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0165)

[2-(4-Amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidin-4-yl]-methanol (MTR-0166)

1-(4-Fluoro-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thio urea (MTR-0167)

1-(3-Fluoro-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thiourea (MTR-0168)

1-{4-[5-Ethoxy-4-(4-methanesulfonyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-phenyl-urea (MTR-0169)

4-[5-Ethoxy-4-(4-methyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-1H-indazole (MTR-0170)

4-[5-Ethoxy-4-(4-methyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenylamine (MTR-0171)

1-{4-[5-Ethoxy-4-(4-methyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-phenyl-urea (MTR-0172)

1-{4-[4-(4-Methanesulfonyl-piperazin-1-yl)-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-phenyl-urea (MTR-0173)

4-(5-Ethoxy-4-morpholin-4-yl-6-morpholin-4-ylmethyl-pyrimidin-2-yl)-1H-indazole (MTR-0174)

4-(5-Ethoxy-4-morpholin-4-yl-6-morpholin-4-ylmethyl-pyrimidin-2-yl)-phenylamine (MTR-0175)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-morpholin-4-ylmethyl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea (MTR-0176)

1-(3,4-Difluoro-phenyl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0177)

1-(4-Fluoro-phenyl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0178)

3-[5-Ethoxy-4-(4-methanesulfonyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenol (MTR-0180)

2-(4-Amino-phenyl)-5-methoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0181)

5-Methoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0182)

5-Methoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid (MTR-0183)

1-{4-[5-Methoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-phenyl-urea (MTR-0184)

1-{4-[4-(4-Methanesulfonyl-piperazine-1-carbonyl)-5-methoxy-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-phenyl-urea (MTR-0185)

1-{4-[5-Ethoxy-4-(4-methyl-piperazine-1-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-phenyl-urea (MTR-0186)

[2-(4-Amino-phenyl)-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-4-yl]-acetic acid methyl ester (MTR-0187)

{5-Methylsulfanyl-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidin-4-yl}-acetic acid methyl ester (MTR-0188)

{5-Methylsulfanyl-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidin-4-yl}-acetic acid (MTR-0189)

2-{5-Methylsulfanyl-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidin-4-yl}-acetamide (MTR-0192)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-piperidin-1-ylmethyl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea (MTR-0193)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-methyl-urea (MTR-0195)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-morpholin-4-yl-urea (MTR-0196)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0197)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-morpholin-4-yl-ethyl)-urea (MTR-0198)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(2-dimethylamino-ethoxy)-phenyl]-urea (MTR-0199)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-thiazol-2-yl-urea (MTR-0200)

11

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-methyl-piperazin-1-yl)-urea (MTR-0201)

5-Ethoxy-2-[4-(3-methyl-ureido)-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0202)

5-Ethoxy-2-[4-[(morpholine-4-carbonyl)-amino]-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0203)

2-[4-[3-(3-Dimethylamino-propyl)-ureido]-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0204)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-pyridin-3-yl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0205)

5-Ethoxy-2-[4-[3-(4-methyl-piperazin-1-yl)-ureido]-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0206)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-morpholin-4-yl-ethyl)-urea (MTR-0207)

1-(3-Dimethylamino-propyl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0208)

1-(4-Methyl-piperazin-1-yl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0209)

1-Methyl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0210)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3-fluoro-4-morpholin-4-yl-phenyl)-urea (MTR-0211)

4-[3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido]-benzoic acid ethyl ester (MTR-0212)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0213)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3-trifluoromethyl-phenyl)-urea (MTR-0214)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-methyl-piperazin-1-yl)-phenyl]-urea (MTR-0215)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-thiomorpholin-4-yl-phenyl)-urea (MTR-0216)

5-Ethoxy-6-morpholin-4-yl-2-[4-[3-(2-morpholin-4-yl-ethyl)-ureido]-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0217)

5-Ethoxy-6-morpholin-4-yl-2-(2-phenylamino-1H-benzimidazol-5-yl)-pyrimidine-4-carboxylic acid ethyl ester (MTR-0218)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-morpholin-4-yl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0219)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-thiazol-2-yl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0220)

5-Ethoxy-2-[4-[3-(3-fluoro-4-morpholin-4-yl-phenyl)-ureido]-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0221)

5-Ethoxy-6-morpholin-4-yl-2-[4-[3-(3-oxo-morpholin-4-yl)-phenyl]-ureido]-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0222)

5-Ethoxy-6-morpholin-4-yl-2-[4-[3-(3-trifluoromethyl-phenyl)-ureido]-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0223)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-morpholin-4-yl-urea (MTR-0224)

12

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0225)

1-(3-Fluoro-4-morpholin-4-yl-phenyl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0226)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0227)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0228)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-methyl-urea (MTR-0229)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-pyridin-2-yl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0230)

5-Ethoxy-2-(4-[3-[4-(4-methyl-piperazin-1-yl)-phenyl]ureido]-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0231)

5-Ethoxy-6-morpholin-4-yl-2-[4-[3-(4-thiomorpholin-4-yl-phenyl)-ureido]-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0232)

5-Ethoxy-6-morpholin-4-yl-2-[4-[3-(4-morpholin-4-yl-phenyl)-ureido]-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0233)

2-[4-[3-(2-Amino-phenyl)-thioureido]-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0234)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-morpholin-4-yl-phenyl)-urea (MTR-0235)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-urea (MTR-0236)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(6-morpholin-4-yl-pyridin-3-yl)-urea (MTR-0237)

1-[4-[4-(4-Methyl-piperazin-1-yl)-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl]-3-pyridin-3-yl-urea (MTR-0238)

1-(3,4-Dimethoxy-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0239)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3-trifluoromethyl-phenyl)-urea (MTR-0240)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0241)

1-(3-Fluoro-4-morpholin-4-yl-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0242)

2-[4-(1H-Benzimidazol-2-ylamino)-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0243)

2-[4-[3-(4-Carbamoyl-phenyl)-ureido]-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0244)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-pyridin-3-yl-ureido)-phenyl]-pyrimidine-4-carboxylic acid (MTR-0245)

1-[4-(4-Methyl-piperazin-1-yl)-phenyl]-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0246)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-thiomorpholin-4-yl-phenyl)-urea (MTR-0247)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-morpholin-4-yl-phenyl)-urea (MTR-0248)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-thiazol-2-yl-urea (MTR-0249)

13

2-{4-[3-(3-Acetylamino-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0250)

2-{4-[3-(3-Carbamoyl-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0251)

5-Ethoxy-6-morpholin-4-yl-2-(4-{3-[3-(2,2,2-trifluoroacetylamino)-phenyl]-ureido}-phenyl)-pyrimidine-4-carboxylic acid ethyl ester (MTR-0252)

1-{4-[5-Ethoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-pyridin-3-yl-urea (MTR-0253)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-2-yl-urea (MTR-0254)

2-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-thiophene-3-carboxylic acid methyl ester (MTR-0255)

1-Benzo[1,3]dioxol-5-yl-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0256)

4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide (MTR-0257)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3,4-dimethoxy-phenyl)-urea (MTR-0258)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-morpholin-4-yl-ethyl)-urea (MTR-259)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-penta-2,4-dienyl]-3-(4-methyl-piperazin-1-yl)-urea (MTR-0260)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-morpholin-4-yl-phenyl)-urea (MTR-0261)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-thiomorpholin-4-yl-phenyl)-urea (MTR-0262)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3-trifluoromethyl-phenyl)-urea (MTR-0263)

4-{3-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzoic acid ethyl ester (MTR-0264)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(6-morpholin-4-yl-pyridin-3-yl)-urea (MTR-0265)

1-[6-(4-Methyl-piperazin-1-yl)-pyridin-3-yl]-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0266)

4-{3-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide (MTR-0267)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[5-(4-methyl-piperazin-1-yl)-pyridin-2-yl]-urea (MTR-0268)

N-(3-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-2,2,2-trifluoroacetamide (MTR-0269)

N-(3-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-acetamide (MTR-0270)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-methyl-piperazine-1-carbonyl)-phenyl]-urea (MTR-0271)

[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester (MTR-0272)

5-Ethoxy-6-morpholin-4-yl-2-{4-[3-(4-sulfamoyl-phenyl)-ureido]-phenyl}-pyrimidine-4-carboxylic acid ethyl ester (MTR-0273)

5-Ethoxy-2-(4-{3-[4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ureido}-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0274)

14

5-Ethoxy-2-(4-{3-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-ureido}-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0275)

5-Ethoxy-6-morpholin-4-yl-2-{4-[3-(6-morpholin-4-yl-pyridin-3-yl)-ureido]-phenyl}-pyrimidine-4-carboxylic acid ethyl ester (MTR-0276)

2-(4-{3-[4-(2-Dimethylamino-ethylcarbamoyl)-phenyl]-ureido}-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0277)

5-Ethoxy-2-(4-{3-[4-(morpholine-4-carbonyl)-phenyl]-ureido}-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0278)

{4-[5-Ethoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-carbamic acid phenyl ester (MTR-0279)

1-{4-[5-Ethoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0280)

[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester (MTR-0281)

4-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide (MTR-0282)

4-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzenesulfonamide (MTR-0283)

4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-2-nitro-phenylamine (MTR-0284)

1-(2-Amino-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thiourea (MTR-0285)

[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0286)

1-(2-Amino-phenyl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thiourea (MTR-0287)

4-{3-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzenesulfonamide (MTR-0288)

(1H-Benzimidazol-2-yl)-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-amine (MTR-0289)

[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester (MTR-0290)

1-(6-Bromo-pyridin-3-yl)-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0291)

1-(2-Amino-phenyl)-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thiourea (MTR-0292)

4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thiourea}-benzoic acid (MTR-0293)

5-Ethoxy-2-(4-{3-[5-(4-methyl-piperazin-1-yl)-pyridin-2-yl]-ureido}-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0294)

5-Ethoxy-2-{4-[3-(2-methoxycarbonyl-thiophen-3-yl)-ureido]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0296)

5-Ethoxy-2-{4-[3-(1H-indazol-4-yl)-ureido]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0297)

5-Ethoxy-2-{4-[3-(4-methyl-1H-benzotriazol-5-yl)-ureido]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0298)

5-Ethoxy-6-morpholin-4-yl-2-{4-[3-(2-oxo-2,3-dihydro-benzooxazol-5-yl)-ureido]-phenyl}-pyrimidine-4-carboxylic acid ethyl ester (MTR-0299)

2-(4-Amino-3-fluoro-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0300)

5-Ethoxy-2-(3-fluoro-4-phenoxy-carbonylamino-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0301)

15

5-Ethoxy-2-(3-fluoro-4-{3-[4-(3-oxo-morpholin-4-yl)-phenyl]-ureido}-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0302)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-thiourea (MTR-0303)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-thiourea (MTR-0304)

(1H-Benzimidazol-2-yl)-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-amine (MTR-0305)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-isopropyl-urea (MTR-0306)

[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid (MTR-0307)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-carbonyl)-phenyl]-urea (MTR-0308)

4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-benzene-1,2-diamine (MTR-0309)

(1H-Benzimidazol-2-yl)-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-amine (MTR-0310)

5-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-1H-benzimidazol-2-ylamine (MTR-0311)

1-[4-(2-Dimethylamino-ethoxy)-phenyl]-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0312)

3-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide (MTR-0313)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-thiazol-2-yl-urea (MTR-0314)

1-(3,4-Dimethoxy-phenyl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0315)

4-{3-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzoic acid (MTR-0316)

1-Benzol[1,3]dioxol-5-yl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0317)

1-[5-(4-Methyl-piperazin-1-yl)-pyridin-2-yl]-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0318)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-thiourea (MTR-0319)

2-(4-Amino-2-fluoro-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0320)

5-Ethoxy-2-(2-fluoro-4-phenoxy-carbonylamino-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0321)

5-Ethoxy-2-(2-fluoro-4-{3-[4-(3-oxo-morpholin-4-yl)-phenyl]-ureido}-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0322)

4-{3-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide (MTR-0323)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-thiomorpholin-4-yl-phenyl)-urea (MTR-0324)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0325)

4-{3-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzenesulfonamide (MTR-0326)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-2-fluoro-phenylamino-urea (MTR-0327)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(5-morpholin-4-yl-pyridin-2-yl)-urea (MTR-0328)

16

1-(4-Amino-phenyl)-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0329)

1-tert-Butyl-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0330)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-amino-urea (MTR-0331)

4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenylamine (MTR-0332)

1-tert-Butylamino-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0333)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[5-(4-methyl-piperazin-1-yl)-pyridin-2-yl]-urea (MTR-0334)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-thiourea (MTR-0335)

[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0336)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-methyl-piperazin-1-yl)-phenyl]-urea (MTR-0337)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-urea (MTR-0338)

1-Isopropyl-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0339)

1-(6-Bromo-pyridin-3-yl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0340)

1-Isopropyl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0341)

1-tert-Butyl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0342)

1-[4-(2-Dimethylamino-ethoxy)-phenyl]-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0343)

2-{4-[3-(4-Carbamoyl-phenyl)-ureido]-3-fluoro-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0344)

2-{4-[3-(4-Carbamoyl-phenyl)-ureido]-2-fluoro-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0345)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(1H-indazol-4-yl)-urea (MTR-0346)

N-(4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-methanesulfonamide (MTR-0347)

1,3-Bis-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0349)

1-(4-Amino-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0350)

4-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzoic acid (MTR-0351)

1-(6-Bromo-pyridin-3-yl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0352)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(6-morpholin-4-yl-pyridin-3-yl)-urea (MTR-0353)

[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenyl]-carbamic acid phenyl Ester (MTR-0354)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0355)

[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenyl]-urea (MTR-0356)

4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenylamine (MTR-0357)

4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzenesulfonamide (MTR-0358)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-3-fluoro-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0359)

4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0360)

2-[4-(3-Benzo[1,3]dioxol-5-yl-ureido)-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0361)

5-Ethoxy-2-{4-[3-(4-ethoxycarbonyl-phenyl)-ureido]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0362)

2-(4-{3-[4-(2-Dimethylamino-ethoxy)-phenyl]-ureido}-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0363)

4-(3-{4-[5-Ethoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-ureido)-benzamide (MTR-0364)

5-Ethoxy-2-{4-[3-(4-methanesulfonyl-phenyl)-ureido]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0365)

5-Ethoxy-2-[4-(3-isopropyl-ureido)-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0366)

2-[4-(3-tert-Butyl-ureido)-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0367)

N-(4-{3-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-acetamide (MTR-0368)

5-Ethoxy-2-{4-[3-(4-methanesulfonylamino-phenyl)-ureido]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0369)

2-{4-[3-(4-Acetylamino-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0370)

N-(4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-acetamide (MTR-0371)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzooxazol-5-yl)-urea (MTR-0372)

6-Chloro-5-methylsulfanyl-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-ylamine (MTR-0373)

1,3-Bis-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0374)

3-Fluoro-4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0375)

N-(4-{3-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-methanesulfonamide (MTR-0376)

N-(3-{3-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-acetamide (MTR-0377)

1-[3-Fluoro-4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0378)

N-(4-{3-[3-Fluoro-4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-acetamide (MTR-0379)

N-(4-{3-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-acetamide (MTR-0380)

1-[4-(5-Methanesulfonyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0381)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-carbonyl)-phenyl]-urea (MTR-0382)

2-Fluoro-4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0383)

3-Fluoro-4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0384)

4-{3-[2-Fluoro-4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide (MTR-0385)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-2-yl-urea (MTR-0386)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-methanesulfonyl-phenyl)-urea (MTR-0387)

[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0388)

1-(6-Chloro-5-methylsulfanyl-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-yl)-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0389)

[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester (MTR-0390)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0391)

4-{3-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide (MTR-0392)

N-(4-{3-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-methanesulfonamide (MTR-0394)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzooxazol-5-yl)-urea (MTR-0395)

4-(5-Ethoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0396)

[4-(5-Ethoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]carbamic acid phenyl ester (MTR-0397)

1-[4-(5-Ethoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0398)

1-[4-(5-Ethoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea (MTR-0399)

4-{3-[4-(5-Ethoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide (MTR-0400)

1-[4-(5-Methanesulfonyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0401)

2'-Amino-5-ethoxy-6-morpholin-4-yl-[2,5']bipyrimidinyl-4-carboxylic acid ethyl ester (MTR-0402)

5-Ethoxy-6-morpholin-4-yl-2'-[3-[4-(3-oxo-morpholin-4-yl)-phenyl]-ureido]-[2,5']bipyrimidinyl-4-carboxylic acid ethyl ester (MTR-0403)

1-(4-Methanesulfonyl-phenyl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0404)

1-[2-Fluoro-4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0405)

1-[3-Fluoro-4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0406)

4-{3-[3-Fluoro-4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide (MTR-0407)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-thiourea (MTR-0408)

4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenyl]-ureido}-benzamide (MTR-0409)

19

4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenyl]-ureido}-benzenesulfonamide (MTR-0410)

4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-3-fluoro-phenyl]-ureido}-benzamide (MTR-0411)

4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-3-fluoro-phenyl]-ureido}-benzenesulfonamide (MTR-0412)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-methanesulfonyl-piperazin-1-yl)-phenyl]-urea (MTR-0413)

N-(5-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-pyridin-2-yl)-acetamide (MTR-0415)

2'-[3-(4-Carbamoyl-phenyl)-ureido]-5-ethoxy-6-morpholin-4-yl-[2,5']bipyrimidinyl-4-carboxylic acid ethyl ester (MTR-0417)

2'-[3-(4-Acetylamino-phenyl)-ureido]-5-ethoxy-6-morpholin-4-yl-[2,5']bipyrimidinyl-4-carboxylic acid ethyl ester (MTR-0418)

5-Ethoxy-2-(4-{3-[4-(4-methanesulfonyl-piperazin-1-yl)-phenyl]-ureido}-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0419)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-methanesulfonyl-piperazin-1-yl)-phenyl]-urea (MTR-0420)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-carbonyl)-phenyl]-urea (MTR-0421)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzooxazol-6-yl)-urea (MTR-0422)

5-Ethoxy-6-morpholin-4-yl-2-{4-[3-(2-oxo-2,3-dihydro-benzooxazol-6-yl)-ureido]-phenyl}-pyrimidine-4-carboxylic acid ethyl ester (MTR-0423)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-methanesulfonyl-phenyl)-urea (MTR-0424)

5-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-1,3-dihydro-benzoimidazol-2-one (MTR-0425)

N-(3-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-acetamide (MTR-0426)

N-(3-Dimethylamino-propyl)-4-{3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide (MTR-0427)

5-Methoxy-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-ylamine (MTR-0428)

4-[3-(5-Methoxy-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-yl)-ureido]-benzamide (MTR-0429)

1-[4-(4-Methanesulfonyl-piperazin-1-yl)-phenyl]-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0430)

5-Methylsulfanyl-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-ylamine (MTR-0431)

4-[3-(6-Chloro-5-methylsulfanyl-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-yl)-ureido]-benzenesulfonamide (MTR-0432)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzooxazol-6-yl)-urea (MTR-0433)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-sulfonyl)-phenyl]-urea (MTR-0435)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-dimethylamino-piperidine-1-carbonyl)-phenyl]-urea (MTR-0436)

4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenylamine (MTR-0437)

20

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea (MTR-0438)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-thiourea (MTR-0440)

N-(5-{3-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-pyridin-2-yl)-acetamide (MTR-0442)

2-{4-[3-(6-Acetylamino-pyridin-3-yl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0443)

N-[3-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-methanesulfonamide (MTR-0444)

N-[3-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-2,4-difluoro-benzenesulfonamide (MTR-0445)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0446)

1-[4-(5-Ethoxy-4-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0447)

4-(5-Methoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0448)

N-(4-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-methanesulfonamide (MTR-0449)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzooxazol-6-yl)-urea (MTR-0450)

1-(5-Methoxy-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-yl)-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0451)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-sulfonyl)-phenyl]-urea (MTR-0452)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0453)

1-[4-(Morpholine-4-carbonyl)-phenyl]-3-[4-[4-morpholin-4-yl-5-(2-morpholin-4-yl-ethoxy)-pyrimidin-2-yl]-phenyl]-urea (MTR-0454)

1-{4-[4-Morpholin-4-yl-5-(2-morpholin-4-yl-ethoxy)-pyrimidin-2-yl]-phenyl}-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0455)

4-(3-{4-[4-Morpholin-4-yl-5-(2-morpholin-4-yl-ethoxy)-pyrimidin-2-yl]-phenyl}-ureido)-benzamide (MTR-0456)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0457)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzooxazol-6-yl)-urea (MTR-0459)

4-(4-Chloro-5-methoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0460)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-carbonyl)-phenyl]-urea (MTR-0461)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0462)

N-(4-{3-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-methane sulfonamide (MTR-0463)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0464)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea (MTR-0465)

4-(5-Methylsulfanyl-2,6-di-pyridin-3-yl-pyrimidin-4-yl)-morpholine (MTR-0466)

N-[3-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-methane sulfonamide (MTR-0467)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-thiourea (MTR-0468)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0469)

21

1-[4-(4-Methanesulfonyl-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0470)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-sulfonyl)-phenyl]-urea (MTR-0471)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0472)

1-[4-(5-Methoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-carbonyl)-phenyl]-urea (MTR-0473)

1-[4-(5-Methoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0474)

1-[4-(4-Chloro-5-methoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0476)

1-Ethyl-3-[4-(5-methylsulfonyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0477)

1-Methyl-3-[4-(5-methylsulfonyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0478)

1-Isopropyl-3-[4-(5-methylsulfonyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0479)

1-[4-(5-Methoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0480)

1-[4-(4-Morpholin-4-yl-5-(2-morpholin-4-yl-ethoxy)-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0481)

4-(5-Methoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0482)

4-(5-Methoxy-4-morpholin-4-yl-6-(4-aminophenyl-pyrimidin-2-yl)-phenylamine (MTR-0483)

1-[4-(4-Chloro-5-methylsulfonyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-isoxazol-3-yl-urea (MTR-0484)

1-Isoxazol-3-yl-3-[4-(5-methylsulfonyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0485)

N-(5-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-pyridin-2-yl)-acetamide (MTR-0486)

1-[4-(4-Methanesulfonyl-piperazin-1-yl)-phenyl]-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0487)

1-[4-(4-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0488)

1-[4-(4-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0489)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0490)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0491)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0492)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0493)

1-[4-(4-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0494)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0495)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0496)

1-[4-(5-Methoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0497)

1-[4-(5-Methoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0498)

1-[4-(5-Methoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0499)

1-[4-(4-Chloro-5-methoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0500)

22

1-[4-(5-Methoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0501)

1-[4-(5-Methoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0502)

1-[4-(4-Chloro-5-methoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-isoxazol-3-yl-urea (MTR-0503)

1-Isoxazol-3-yl-3-[4-(5-methoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0504)

1-[4-(4-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-isoxazol-3-yl-urea (MTR-0505)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-isoxazol-3-yl-urea (MTR-0506)

1-Isoxazol-3-yl-3-[4-(5-methoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0507)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-isoxazol-3-yl-urea (MTR-0508)

1-Isoxazol-3-yl-3-[4-(5-methylsulfonyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0509)


or a stereoisomer, or a tautomer, or an N-oxide, or a pharmaceutically acceptable salt, or an ester, or a prodrug, or a hydrate, or a solvate thereof.

The pyrimidine compounds of formula (I), or a stereoisomer, or a tautomer, or an N-oxide, or a pharmaceutically acceptable salt, or an ester, or a prodrug, or a hydrate, or a solvate thereof, may be prepared by any process known to be applicable to the preparation of chemically related compounds.


According to the invention, suitable pharmaceutically acceptable salts of formula (I) include hydrochloride, hydrobromide, hydroiodide, sulfate, phosphate, mesylate, besylate, acetate, oxalate, citrate, lactate, tartrate, succinate, methane-sulfonate, trifluoroacetate, and maleate salts. The preferred salt is a hydrochloride salt. When the compounds of the invention bear a free carboxy substituent, the salts include both the above-mentioned acid addition salts and the salts of sodium, potassium, calcium and ammonium. The latter are prepared by treating the free pyrimidine of formula (I), or the acid addition salt thereof, with the corresponding metal base or ammonia.

The pyrimidine compounds of formula (I) may be prepared by any suitable synthetic routes. Examples of the routes can be those set out in Schemes 1 to 10 below.

Scheme 1




23



In Scheme 1, R^1 , R^2 , Y and W are as defined above for formula (I). A compound of formula (1) which is a known compound or is prepared by methods known in the literature is converted into a compound of formula (2) by treatment with a strong base. The base is typically sodium methoxide, sodium ethoxide, potassium methoxide or potassium ethoxide. A compound of formula (3) is prepared by treatment of the compound of formula (2) with urea in an appropriate solvent such as ethanol. Compounds of formula (4) may be prepared by treating compounds of formula (3) with phosphorous oxychloride in the presence of an N,N-dialkylaniline. Compounds of formula (5) may be prepared by treating compounds of formula (4) with an amine of formula HW in an inert solvent in the presence of a base. Compounds of formula (Ia) may be prepared by the Suzuki coupling of a compound of formula (5) with a boronic acid or a boronic ester.

Scheme 2




In Scheme 2, $YR^2=SCH_3$, $Z=Cl$ or Br , W and R^1 are as defined above for formula (I). Compounds of formula (6) may be prepared by treating Barbituric acid with dimethyl sulfoxide. Compounds of formula (7) may be prepared by treating compounds of formula (6) with phosphorous oxychloride or phosphorous oxybromide in the presence of an N,N-dialkyl-

24


laniline. Compounds of formula (8) may be prepared by treating compounds of formula (7) with an amine of formula HW in an inert solvent in the presence of a base. Compounds of formula (I) may be prepared by the Suzuki coupling of a compound of formula (8) with a boronic acid or a boronic ester.


Scheme 3



In Scheme 3, $YR^2=SCH_3$, W and R^1 are as defined above for formula (I). The compound of formula (3) wherein $YR^2=SCH_3$ is prepared from 5-Iodouracil according to the method in *Organic Letters*, 2007, 9, 1639. Compounds of formula (4) may be prepared by treating compounds of formula (3) with phosphorous oxychloride in the presence of an N,N-dialkylaniline. Compounds of formula (5) may be prepared by treating compounds of formula (4) with an amine of formula HW in an inert solvent in the presence of a base. Compounds of formula (Ia) may be prepared by the Suzuki coupling of a compound of formula (5) with a boronic acid or a boronic ester.


Scheme 4





5
10
15

Scheme 5




In Scheme 4, $YR^2 = -OCH_3$ or $-OEt$, $Z = -C(O)NR_3R_4$, R^1 and W are as defined above for formula (I). A compound of formula (9), which is a known compound or is prepared by methods known in the literature, is converted into a compound of formula (10) by treatment with a strong base. The base is typically sodium methoxide, sodium ethoxide, potassium methoxide or potassium ethoxide. A compound of formula (11) is prepared by treatment of the compound of formula (10) with urea in an acid solution. Compounds of formula (11) may be converted to compounds of formula (12) in the presence of a base solution. Compounds of formula (13) may be prepared by the acetylation of a compound of

55
60
65

In Scheme 5, $YR^2 = -OCH_3$ or $-OEt$, $Z = -CH_2NR_3R_4$, R^1 and W are as defined above for formula (I). Compounds of formula (18) can be obtained from compounds of formula (15) by treatment with sodium borohydride or other reducing agent. Des s-Martin periodinate or other oxidizing agent is used to oxidize compounds of formula (18) to compounds of formula (19). Reductive amination of compounds of formula (19) using the appropriate amine and sodium triacetoxyborohydride proceeds smoothly to yield compounds of formula (20). Compounds of formula (I) may be prepared by the Suzuki coupling of a compound of formula (20) with a boronic acid or a boronic ester.

27

Scheme 6




In Scheme 6, $YR^2 = SCH_3$, $Z = -CH_2C(O)NR_3R_4$, R^1 and W are as defined above for formula (I). Compounds of formula (7) may be prepared by the method of scheme 2. Compounds of formula (21) may be prepared by reacting together compounds of formula (7) and sodium methoxide in MeOH. Compounds of formula (22) may be prepared by reacting

28

together compounds of formula (21) and dimethyl malonate in the presence of sodium hydride. Compounds of formula (23) can be obtained by demethylation and decarboxylation of compounds of formula (22) in the presence of an excess of alkali hydroxide. Compounds of formula (24) may be prepared by treating compounds of formula (23) with phosphorous oxychloride or phosphorous oxybromide in the presence of an N,N-dialkylaniline. Compounds of formula (25) may be prepared by treating compounds of formula (24) with an amine of formula HW in an inert solvent in the presence of a base. Compounds of formula (26) may be prepared by the Suzuki coupling of a compound of formula (25) with a boronic acid or a boronic ester. Compounds of formula (27) can be obtained by hydrolysis of compounds of formula (26). Compounds of Formula (I) can be prepared by coupling of compounds of formula (27) with an amine by one of the standard methods of amide bond formation.

Scheme 7




In Scheme 7, $R_7 = OMe$, OEt , $YR^2 = OCH_3$, OEt , $Z = Cl$, Br , R^1 and W are as defined above for formula (I). Compounds of formula (30) are prepared from carbonic acid diethyl ester


29

(28, $R_7=OEt$) according to the method in J. of Heterocyclic Chemistry, 1989, 1261-1271 or from compounds of formula (29) according to the method in J. Med. Chem., 1974, 1197. Compounds of formula (6) are prepared by treating the compounds of formula (30) with urea in an appropriate solvent such as ethanol. Compounds of formula (7) may be prepared by treating compounds of formula (6) with phosphorous oxychloride in the presence of an N,N-dialkylaniline. Compounds of formula (8) may be prepared by treating compounds of formula (7) with an amine of formula HW in an inert solvent in the presence of a base. Compounds of formula (I) may be prepared by the Suzuki coupling of a compound of formula (8) with a boronic acid or a boronic ester.

Scheme 8




30



In Scheme 8, $R_7=OMe$, OEt , $YR^2=OCH_3$, OEt , $Z=Cl$, Br , W is as defined above for formula (I). The use of diverse amidines in the cyclization reaction with substituted 1,3-dicarbonyl compound can obtain pyrimidines bearing a carbon-carbon linkage in position 2. The desired amidines are either commercially available or can be obtained from known procedures by one skilled in the art. Compounds of formula (31) or (35) may be prepared by treating appropriate amidine with 1,3-dicarbonyl compound (2) or (30). Compounds of formula (32) or (36) may be prepared by treating compounds of formula (31) or (35) with phosphorous oxychloride in the presence of an N,N-dialkylaniline. Compounds of formula (33) or (37) may be prepared by treating compounds of formula (32) or (36) with an amine of formula HW in an inert solvent in the presence of a base. Compounds of formula (34) or (38) may be prepared by the reduction of compounds of formula (33) or (37) with a hydrogen/palladium on carbon.


31


32



Substitution at the 4-Cl is not limited to an amino group, as described in Scheme 1-8. 4-Cl can also bear a carbon linker. In Scheme 9, R^2 , Y , W (W_a , W_b), $R_g = NR_3R_4$ are as defined above for formula (I). Each chlorine atom of the compound of formula (4) is selectively replaced at different conditions. Compounds of formula (4) are reacted with different amines (W_a =morpholine or substituted morpholine derivatives) to yield the compound of formula (5a) and with aryl and heteroaryl boronic acid (ester) (W_b) by Suzuki coupling in the presence of palladium catalyst to yield the compound of formula (5b). The second chlorine atom is replaced with 4-aminoaryl and aminoheteroaryl boronic acid(ester) in the presence of palladium catalyst to yield (Ia) and (Ib) respectively. The amino group is converted to the urea derivatives by

three different procedures depending upon the availability of the starting material. Some of the examples shown here are converted into the urea derivatives by reacting (Ia) or (Ib) with an appropriately substituted isocyanate or thioisocyanate derivative. Some of the urea derivatives reported here are prepared by reacting (Ia) or (Ib) with triphosgene in presence of triethylamine and an appropriately substituted primary amine derivative. The corresponding carbamate derivatives are prepared by reacting (Ia), (Ib) or a substituted amine derivative with a phenyl chloroformate reagent. The phenyl N-substituted carbamates are reacted with different substituted amine, heteroalkyl amine or heteroaryl aniline to yield the compound of formula (Ia-a) or (Ib-b).





Substitution at the 6-Cl is not limited to an amino group, as described in Scheme 2, 7 and 8. 6-Cl can also bear an oxygen or a carbon linker. In Scheme 10, R^2 , Y, W(Wa, Wb, Wc) and $R_g = NR_3R_4$ are as defined above for formula (I). Each chlorine atom of the compound of formula (7) is selectively replaced at different conditions. Substitution at the chlorine position is not limited to an amino group. Chlorine position can also bear an oxygen or a carbon linker. 2,4,6-Trisubstituted pyrimidines can be obtained via standard procedure (i.e. SNAr, Mitsunobu, Suzuki, Stille and Heck couplings). Compounds of formula (7) are reacted with different amines (Wa=morpholine or substituted morpholine derivatives) to yield the compound of formula (8a) and with aryl and heteroaryl boronic acid (ester) (Wb) by Suzuki coupling in the presence of palladium catalyst to yield the compound of formula (8b). The second chlorine atom is replaced with 4-aminoaryl and aminoheteroaryl boronic acid(ester) in the presence of palladium catalyst to yield (Ic) and (Id) respectively. The third chlorine atom is replaced with alkyl, alkene, alkyne, aryl or heteroaryl by organomagnesium or organozinc or organoboronic ester reagents to yield (Ic-c) and (Id-d) respectively. The amino group of (Ic-c) and (Id-d) is converted to the urea derivatives by three different procedures as described above (Scheme 9) to yield the compound of formula (Ic-c-c) and (Id-d-d). Compounds of formula (8a) and (8b) also are reacted with different amines and alcohols to give (Ie), (If), (Ig), and (Ih), respectively. Compounds of formula (Ie), (If), (Ig) and (Ih) then underwent the same procedures as described above to yield compounds of formula (Ie-e-e), (If-f-f), (Ig-g-g) and (Ih-h-h).

It has now been found that a series of novel pyrimidine compounds have inhibitory activity against the PI3K enzymes and the class IV kinase mTOR. It is now well understood that deregulation of oncogenes and tumour-suppressor genes contributes to the formation of malignant tumours, for example, by way of increased cell proliferation or increased cell survival. It is also now known that signaling pathways mediated by the PI3K/mTOR families have a central role in a number of cell processes, including proliferation and survival, and deregulation of these pathways is a causative factor in a wide spectrum of human cancers and other diseases.

The compounds of the present invention have been found to be inhibitors of mTOR kinase and PI3 kinase. The pharmacological inhibitors of mTOR kinase and PI3 kinase should be of therapeutic value for treatment of various forms of cancer comprising solid tumors such as carcinomas, sarcomas, leukaemias and lymphoid malignancies. Accordingly, a compound of the present invention can be used to treat a disease or disorder arising from abnormal cell growth, function or behaviour associated with mTOR kinase and PI3 kinase.

Also within the scope of this invention are a pharmaceutical composition that contains an effective amount of at least one of the pyrimidine compounds of formula (I) or a stereoisomer, or a tautomer, or an N-oxide, or a pharmaceutically acceptable salt, or an ester, or a prodrug, or a hydrate, or a solvate thereof together with a pharmaceutically acceptable carrier, a method for treating a PI3K kinase-/mTOR kinase-related disease (e.g., cancer) by administering to a subject in need of this treatment an effective amount of the pyrimidine

35

compounds of formula (I), and a method of decreasing the activity of at least one PI3K kinase and mTOR kinase by contacting the at least one PI3K kinase and mTOR kinase with at least one of the pyrimidine compounds of formula (I).

As used herein, the term “PI3 kinase-/mTOR kinase-related disease” refers to a disease or condition that is characterized by abnormal PI3 and/or mTOR activity or a disease or condition that can be treated with changes to the activity of at least one of PI3 and mTOR. Abnormal PI3 and/or mTOR activity can arise as the result of elevated PI3 and/or mTOR expression level, or presence of PI3 and/or mTOR expression that does not occur in normal conditions. PI3 kinase-/mTOR kinase-related diseases described herein include, but are not limited to, cancer, diabetes, immune disorders, hyper-proliferation disorders, hyperproliferative disorders of the kidney, renal disease, von Hippel-Lindau disease, restenosis, fibrosis, psoriasis, osteoarthritis, rheumatoid arthritis, inflammatory disorders, immunological disorders such as autoimmune diseases (e.g., AIDS, lupus, etc.), cardiovascular disorders (e.g. atherosclerosis), and blood vessel proliferative disorders such as abnormal vasculogenesis.

The term “treating” refers to administering a pyrimidine compound of formula (I) to a subject that has a PI3 kinase-/mTOR kinase-related disease, or has a symptom of or a predisposition toward it, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve, affect or reduce the risk of the disorder, or the symptoms of or the predisposition toward the disorder. For example, treating cancer refers to treatment resulting in inhibition of cancer growth or cancer cell growth, regression in cancer growth (i.e. reducing the size of a detectable cancer), or disappearance of a cancer. The term “an effective amount” refers to the amount of the active agent that is required to confer the intended therapeutic effect in the subject. Effective amounts may vary, as recognized by those skilled in the art, depending on route of administration, the excipient used, and the possibility of co-usage with other agents. The subject in need of the treatment can be a mammal. The term “mammal” refers to human or nonhuman mammal, for example, dogs, cats, pigs, cows, sheep, goats, horses, rats, or mice.

Cancer that can be treated by the methods of the invention is any abnormal cell or tissue growth, for example, a tumor, whether malignant, pre-malignant, or non-malignant. It is characterized by uncontrolled proliferation of cells that may or may not invade the surrounding tissue and, hence, may or may not metastasize to new body sites. Cancer encompasses carcinomas, which are cancers of epithelial cells; carcinomas include squamous cell carcinomas, adenocarcinomas, melanomas, and hepatomas. Cancer also encompasses sarcomas, which are tumors of mesenchymal origin; sarcomas include osteogenic sarcomas, leukemias, and lymphomas. Cancers may involve one or more neoplastic cell type. The term cancer includes, as non-limiting examples, lung cancer, colon cancer, colorectal cancer, breast cancer, prostate cancer, liver cancer, pancreatic cancer, bladder cancer, gastric cancer, renal cancer, salivary gland cancer, ovarian cancer, uterine body cancer, cervical cancer, oral cancer, skin cancer, brain cancer, lymphoma, and leukemia. The cancers also include Epidermal Growth Factor Receptor (EGFR) dependent cancers or cancers that resist to EGFR targeting agent.

The compounds described herein can be administered to a mammal in conjunction with radiation therapy, immunotherapy, monoclonal antibody therapy, hormonal therapy, chemotherapy using other agents, and/or surgery. “In conjunction with” means that the therapies do not need to occur at the same time, and can be in succession, or alternate with each other and/or periods of rest and recovery.

36

In one embodiment, a PI3 kinase-/mTOR kinase-related disease, such as cancer, can be treated with a method comprising administering an effective amount of at least a pyrimidine compound of formula (I) and at least one chemotherapeutic agent to a mammal. Non-limiting examples of chemotherapeutic agent include protein kinase inhibitors other than the compound described herein (e.g., imatinib mesylate, gefitinib, dasatinib, erlotinib, lapatinib, sunitinib, nilotinib, and sorafenib; antibodies, including, e.g., trastuzumab, rituximab, cetuximab, and bevacizumab; mitoxantrone; dexamethasone; prednisone; and temozolomide), alkylating agents (e.g., melphalan, chlorambucil, busulfan, thiopeta, ifosfamide, carmustine, lomustine, semustine, streptozocin, decarbazine, and cyclophosphamide), mitotic inhibitors, antimetabolites (e.g., capecitabine, gemcitabine, 5-fluorouracil or 5-fluorouracil/leucovorin, fludarabine, cytarabine, mercaptopurine, thioguanine, pentostatin, and methotrexate), cell cycle inhibitors, enzymes, hormones, anti-hormones, growth-factor inhibitors, plant alkaloids and terpenoids, topoisomerase inhibitors (e.g., etoposide, teniposide, camptothecin, topotecan, irinotecan, doxorubicin, and daunorubicin), antitumor antibiotics (e.g., actinomycin D, bleomycin, mitomycin C, adriamycin, daunorubicin, idarubicin, doxorubicin and pegylated liposomal doxorubicin), vinca alkaloids (e.g., vincristine and vinblastin), taxanes (e.g., paclitaxel and docetaxel), platinum agents (e.g., cisplatin, carboplatin, and oxaliplatin), thalidomide and related analogs (e.g., CC-5013 and CC-4047), monoclonal antibodies, antiangiogenic agents, and combinations thereof.

To practice the method of this invention, the above-described pharmaceutical composition can be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term “parenteral” as used herein includes subcutaneous, intracutaneous, intravenous, intramuscular, intraarticular, intraarterial, intrasynovial, intrasternal, intrathecal, intraleisional, and intracranial injection or infusion techniques. In one embodiment, the pharmaceutical composition of this invention is administered intravenously. The pharmaceutically acceptable carriers may include, but are not limited to, water, Ringer’s solution, isotonic sodium chloride solution or phosphate buffered saline, and solutions containing thickening and solubilizing agents, such as glucose, polyethylene glycol, and polypropylene glycol and mixtures thereof.

A sterile injectable composition can be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation can also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent. Among the acceptable vehicles and solvents that can be employed are mannitol, water, Ringer’s solution and isotonic sodium chloride solution. Fatty acids, such as oleic acid and its glyceride derivatives, are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions can also contain a long-chain alcohol diluent or dispersant, or carboxymethyl cellulose or similar dispersing agents. Other commonly used surfactants such as Tweens or Spans or other similar emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms can also be used for purposes of formulation.

A composition for oral administration can be any orally acceptable dosage form including, but not limited to, capsules, tablets, emulsions and aqueous suspensions, dispersions and solutions. In the case of tablets for oral use, carriers

37

that are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions or emulsions are administered orally, the active ingredient can be suspended or dissolved in an oily phase combined with emulsifying or suspending agents. If desired, certain sweetening, flavoring, or coloring agents can be added. A nasal aerosol or inhalation composition can be prepared according to techniques well known in the art of pharmaceutical formulation. The composition of the present invention may also be administered in the form of suppositories for rectal administration.

Suitable in vitro assays can be used to preliminarily evaluate the efficacy of the pyrimidine compounds of formula (I) in anticancer activities such as inhibiting growth of tumor cells. The compounds can further be examined for their efficacy in treating cancer. For example, a compound can be administered to an animal (e.g., a mouse model) having cancer and its therapeutic effects then assessed. Based on the results, an appropriate dosage range and administration route can also be determined.


The invention will be further described in the Examples as follows. The examples given below are intended to be illustrative only and not to limit the invention. Any modifications and variations that can be easily made by those skilled in the art fall within the scope of the disclosure of the specification and the appended claims of the present invention.

EXAMPLES

Example 1


Preparation of Compounds of Formula (I) in Scheme 4

2-Ethoxy-malonic acid diethyl ester



A mixture of diethyl oxalate (39.23 ml, 1.12 eq.) and ethyl ethoxyacetate (34.95 ml, 1 eq.) was added dropwise to a slurry of sodium ethoxide (18.64 g, 1.07 eq.) in toluene (100 ml) at 45-50° C. After dropping, the resulting solution was heated to 70-80° C. for 2 hrs and poured into 70 ml of 14% HCl with cooling. The resultant mixture was extracted with EA and the combined organic layers were washed with brine and dried in vacuo to give 55.86 g (93.7%) of a product.


2-Methoxy-malonic acid diethyl ester



38

A mixture of diethyl oxalate (27.5 ml, 1 eq.) and methyl methoxyacetate (20 ml, 1 eq.) was added dropwise to a slurry of sodium methoxide (16.4 g, 1.5 eq.) in toluene (350 ml) at 45-50° C. After dropping, the resulting solution was heated to 70-80° C. for 2 hrs and poured into 70 ml of 14% HCl with cooling. The resultant mixture was extracted with EA and the combined organic layers were washed with brine and dried in vacuo to give 16.7 g (40.5%) of a product.


(2,5-Dioxo-imidazolidin-4-ylidene)-ethoxy-acetic acid ethyl ester



A mixture of diethyl 2-ethoxy-3-oxo-succinate (55.86 g, 1 eq) and urea (14.41 g, 1 eq.) was refluxed for 2.5 h in 1 M HCl—AcOH (1200 ml, 5 eq.) and allowed to become cool. The cooled mixture was evaporated to dryness. The residues were dissolved in MeOH, heated aside for crystallization. After crystallization, the precipitates were filtered to give a product which was an off-white solid (15.11 g, 27.5%).


¹H NMR (500 MHz, DMSO-d₆): δ 1.21-1.27 (m, 6H), 3.82-3.86 (m, 2H), 4.24-4.29 (m, 2H), 9.79 (s, 1H), 11.33 (s, 1H)

(2,5-Dioxo-imidazolidin-4-ylidene)-methoxy-acetic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.23-1.28 (m, 3H), 3.61 (s, 3H), 4.25-4.29 (m, 2H), 9.85 (s, 1H), 11.39 (s, 1H)

5-Ethoxy-2,6-dihydroxy-pyrimidine-4-carboxylic acid




39

(2,5-dioxo-imidazolidin-4-ylidene)-ethoxy-acetic acid ethyl ester (16.02 g, 1 eq.) was dissolved in aq. 1 N KOH (281 ml, 4 eq.) and refluxed for 3.5 hrs. The mixture was cooled to 0° C. and carefully acidified with conc. HCl. After overnight cooling (4° C.), it led to the formation of white precipitate. The mixture was filtered and the precipitate was dried in vacuo. The product obtained was a white solid (7.76 g, 55%).


¹H NMR (500 MHz, DMSO-d₆): δ 1.17-1.23 (m, 3H), 3.89-3.93 (m, 2H), 10.60 (s, 1H), 11.42 (s, 1H)

2,6-Dihydroxy-5-methoxy-pyrimidine-4-carboxylic acid



¹H NMR (500 MHz, DMSO-d₆): δ 3.61 (s, 3H), 10.53 (s, 1H), 11.39 (s, 1H)


5-Ethoxy-2,6-dihydroxy-pyrimidine-4-carboxylic acid ethyl ester



Acetyl chloride (100 ml, 36 eq.) was added dropwise to absolute EtOH (780 ml) at 0° C. After stirring the resulting solution for 20 min at r.t., 5-ethoxy-2,6-dihydroxy-pyrimidine-4-carboxylic acid (7.76 g, 1 eq.) was added in one portion and the mixture was refluxed overnight. The volatiles were evaporated in vacuo to give 5-ethoxy-2,6-dihydroxy-pyrimidine-4-carboxylic acid ethyl ester as an off-white solid (8.85 g, 100%).

¹H NMR (500 MHz, DMSO-d₆): δ 1.18-1.30 (m, 6H), 3.91-3.95 (m, 2H), 4.25-4.31 (m, 2H), 10.74 (s, 1H), 11.51 (s, 1H)


5-Methoxy-2,6-dihydroxy-pyrimidine-4-carboxylic acid ethyl ester



40

¹H NMR (500 MHz, DMSO-d₆): δ 1.23-1.30 (m, 3H), 3.65 (s, 3H), 4.27-4.32 (m, 2H), 10.80 (s, 1H), 11.54 (s, 1H)


2,6-Dichloro-5-ethoxy-pyrimidine-4-carboxylic acid ethyl ester



N,N-dimethylaniline (6.85 ml) was added to a stirred solution of 5-ethoxy-2,6-dihydroxy-pyrimidine-4-carboxylic acid ethyl ester (8.85 g) in POCl₃ (265 ml) and the mixture was refluxed overnight. Excess POCl₃ was evaporated in vacuo and the residue was poured into ice-water and extracted with ether. The combined ethereal layers were washed with brine, dried and evaporated in vacuo. Purification by flash chromatography gave a product (8.33 g, 81%).


¹H NMR (500 MHz, DMSO-d₆): δ 1.23-1.35 (m, 3H), 4.13-4.18 (m, 2H), 4.39-4.43 (m, 2H)

2,6-Dichloro-5-methoxy-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.26-1.35 (m, 3H), 3.93 (s, 3H), 4.40-4.48 (m, 2H)

2-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




2,6-dichloro-5-ethoxy-pyrimidine-4-carboxylic acid ethyl ester (2 g, 1 eq.) was stirred in toluene (80 ml) and a solution of morpholine (0.73 ml, 1.1 eq.) in toluene (20 ml) was added dropwise in about 10 minutes at -10-0° C., after stirring the resulting solution for 3 h at r.t. Water was added and extracted with EA. The combined organic layers were washed with

41

brine, dried and evaporated in vacuo. Purification by flash chromatography gave a product (1.79 g, 75%).


¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.41 (m, 6H), 3.76-3.78 (m, 4H), 3.87-3.89 (m, 4H), 3.92-3.96 (m, 2H), 4.39-4.43 (m, 2H)

2-Chloro-5-methoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ 1.38-1.42 (m, 3H), 3.74 (s, 3H), 3.77-3.79 (m, 4H), 3.87-3.89 (m, 4H), 4.40-4.44 (m, 2H)

5-Ethoxy-2-(3-hydroxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



A mixture of 2-chloro-5-ethoxy-6-morpholine-4-yl-pyrimidine-4-carboxylic acid ethyl ester (152 mg, 1 eq.), 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenol (137.9 mg, 1.3 eq.), dioxane (6 ml), H₂O (1 ml), Pd(PPh₃)₄ (56 mg, 0.1 eq.) and sodium carbonate (102 mg, 2 eq.) was heated to reflux overnight. The solvent was removed in vacuo. The residue was extracted with EA, and the organic layer washed with brine and dried. The crude was purified by chromatography to give a product (99.6 mg, 55%).


¹H NMR (500 MHz, CDCl₃-d₁): δ 1.36-1.39 (m, 3H), 1.44-1.46 (m, 3H), 3.85-3.87 (m, 4H), 3.91-3.93 (m, 4H), 3.99-4.03 (m, 2H), 4.46-4.50 (m, 2H), 6.94-6.95 (m, 1H), 7.27-7.33 (m, 1H), 7.82-7.83 (m, 1H), 7.91-7.92 (m, 1H)

Compounds of the following Examples were synthesized following the synthetic method described above. The bionic acids or bionic esters are readily recognizable by one skilled

42


in the art and are commercially available from Aldrich, Acros Organics and Maybridge Chemical Company Ltd.

5-Ethoxy-2-(4-hydroxy-3-methoxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.36 (m, 3H), 1.42-1.45 (m, 3H), 3.82-3.88 (m, 8H), 3.96-3.99 (m, 5H), 4.47-4.49 (m, 2H), 5.84 (s, 1H), 6.94-6.96 (m, 1H), 7.89-7.91 (m, 2H)

5-Ethoxy-2-(3-fluoro-4-methoxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ 1.34-1.36 (m, 3H), 1.42-1.45 (m, 3H), 3.82-3.84 (m, 4H), 3.88-3.89 (m, 4H), 3.95 (s, 3H), 3.94-4.00 (m, 2H), 4.44-4.49 (m, 2H), 6.96-6.99 (m, 1H), 8.04-8.11 (m, 2H)


2-(3-Amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ 1.34-1.36 (m, 3H), 1.42-1.44 (m, 3H), 3.74 (s, 2H), 3.81-3.84 (m, 4H), 3.88-3.90 (m, 4H), 3.96-4.00 (m, 2H), 4.44-4.49 (m, 2H), 6.75-6.76 (m, 1H), 7.20-7.26 (m, 1H), 7.68-7.74 (m, 1H)


43

2-(3,5-Difluoro-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, CDCl₃-d₁): δ 1.35-1.38 (m, 3H), 1.43-1.46 (m, 3H), 3.83-3.84 (m, 4H), 3.89-3.91 (m, 4H), 3.98-4.02 (m, 2H), 4.45-4.50 (m, 2H), 6.84-6.88 (m, 1H), 7.84-7.86 (m, 2H)

5-Ethoxy-2-(1H-indol-6-yl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ 1.34-1.37 (m, 3H), 1.43-1.46 (m, 3H), 3.84-3.86 (m, 4H), 3.91-3.93 (m, 4H), 3.98-4.02 (m, 2H), 4.46-4.50 (m, 2H), 6.57 (s, 1H), 7.26-7.28 (m, 1H), 7.65-7.66 (m, 1H), 8.13-8.14 (m, 1H), 8.33 (s, 1H), 8.45 (s, 1H)

5-Ethoxy-2-(1H-indol-5-yl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, CDCl₃-d₁): δ 1.34-1.37 (m, 3H), 1.43-1.46 (m, 3H), 3.84-3.86 (m, 4H), 3.91-3.93 (m, 4H), 3.97-

44


4.02 (m, 2H), 4.46-4.50 (m, 2H), 6.63 (s, 1H), 7.21-7.22 (m, 1H), 7.39-7.41 (m, 1H), 8.23-8.28 (m, 2H), 8.67 (s, 1H)

5-Ethoxy-2-(1H-indazol-4-yl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, CDCl₃-d₁): δ 1.37-1.40 (m, 3H), 1.46-1.49 (m, 3H), 3.87-3.88 (m, 4H), 3.93-3.95 (m, 4H), 4.03-4.07 (m, 2H), 4.48-4.52 (m, 2H), 7.45-7.48 (m, 1H), 7.58-7.59 (m, 1H), 8.21-8.22 (m, 1H), 8.98 (s, 1H)

2-Benzo[1,3]dioxol-5-yl-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.36 (m, 3H), 1.42-1.45 (m, 3H), 3.82-3.83 (m, 4H), 3.87-3.88 (m, 4H), 3.95-3.99 (m, 2H), 4.44-4.48 (m, 2H), 6.00 (s, 1H), 6.84-6.86 (m, 1H), 7.81 (s, 1H), 7.92-7.94 (m, 1H)

5-Ethoxy-2-(2-fluoro-3-methoxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, CDCl₃-d₁): δ 1.35-1.38 (m, 3H), 1.41-1.44 (m, 3H), 3.79-3.80 (m, 4H), 3.90-3.92 (m, 4H), 3.98-


45

4.03 (m, 2H), 4.43-4.47 (m, 2H), 7.00-7.04 (m, 1H), 7.10-7.13 (m, 1H), 7.57-7.59 (m, 1H)

2-(4-Amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, CDCl₃-d₁): δ 1.32-1.35 (m, 3H), 1.42-1.44 (m, 3H), 3.82-3.87 (m, 10H), 3.94-3.99 (m, 2H), 4.43-4.48 (m, 2H), 6.68-6.70 (m, 2H), 8.15-8.16 (m, 2H) 5-Methoxy-2-(3-hydroxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, CDCl₃-d₁): δ 1.44-1.47 (m, 3H), 3.81 (s, 3H), 3.86-3.88 (m, 4H), 3.91-3.93 (m, 4H), 4.47-4.51 (m, 2H), 5.33 (s, 1H), 6.94-6.95 (m, 1H), 7.29-7.34 (m, 1H), 7.82-7.83 (m, 1H), 7.91-7.93 (m, 1H)

5-Methoxy-2-(1H-indazol-4-yl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester

**46**


¹H NMR (500 MHz, CDCl₃-d₁): δ 1.45-1.48 (m, 3H), 3.83 (s, 3H), 3.87-3.87 (m, 4H), 3.92-3.94 (m, 4H), 4.48-4.53 (m, 2H), 7.44-7.47 (m, 1H), 7.58-7.60 (m, 1H), 8.20-8.21 (m, 1H), 8.98 (m, 1H)

2-(4-hydroxy-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.36 (m, 3H), 1.41-1.43 (m, 3H), 3.82-3.84 (m, 4H), 3.86-3.89 (m, 4H), 3.95-3.99 (m, 2H), 4.43-4.47 (m, 2H), 5.79 (s, 1H), 6.84-6.86 (m, 2H), 8.21-8.22 (m, 2H)

2-(6-Amino-pyridin-3-yl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.36 (m, 3H), 1.42-1.45 (m, 3H), 3.81-3.83 (m, 4H), 3.87-3.88 (m, 4H), 3.95-3.99 (m, 2H), 4.44-4.48 (m, 2H), 4.67 (s, 2H), 6.51-6.52 (m, 1H), 8.34-8.36 (m, 1H), 9.01-9.02 (m, 1H)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester




47

Phenyl isocyanate (0.22 ml, 1.5 eq.) was added to a stirred solution of 2-(4-Amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (500 mg, 1 eq.) in toluene (25 ml) and the mixture was reacted for 4 h at 80° C. The reaction mixture was cooled, the solvent was removed in vacuo, and the residue was washed with EA and filtered to give a product (550 mg, 83%).

¹H NMR (500 MHz, Aetone-d₆): δ 1.31-1.40 (m, 6H), 3.81-3.83 (m, 4H), 3.88-3.90 (m, 4H), 3.98-4.03 (m, 2H), 4.39-4.43 (m, 2H), 6.98-7.01 (m, 1H), 7.27-7.30 (m, 2H), 7.54-7.56 (m, 2H), 7.62-7.64 (m, 2H), 8.17 (s, 1H), 8.27-8.29 (m, 2H), 8.33 (s, 1H)

5-Ethoxy-2-[4-(3-ethyl-ureido)-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




Ethyl isocyanate (0.012 ml, 1.4 eq.) was added to a stirred solution of 2-(4-amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (40 mg, 1 eq.) in toluene (2 ml) and the mixture was reacted overnight at 80° C. The reaction mixture was extracted with EA, washed with brine. The crude was purified by chromatography to give a product (12 mg, 25.2%).

¹H NMR (500 MHz, Aetone-d₆): δ 1.10-1.13 (m, 3H), 1.31-1.39 (m, 6H), 3.25 (m, 2H), 3.81-3.82 (m, 4H), 3.86-3.87 (m, 4H), 4.00-4.01 (m, 2H), 4.39-4.41 (m, 2H), 5.85 (m, 1H), 7.55-7.57 (m, 2H), 8.09 (s, 1H), 8.21-8.23 (m, 2H)

The urea compounds of the following Examples were synthesized following the synthetic method described above by the different isocyanate that are commercially available.

2-{3-[3-(4-Chloro-3-trifluoromethyl-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, Aetone-d₆): δ 1.32-1.35 (m, 3H), 1.38-1.41 (m, 3H), 3.81-3.83 (m, 4H), 3.89-3.91 (m, 4H), 4.02-

48


4.05 (m, 2H), 4.40-4.44 (m, 2H), 7.38-7.41 (m, 1H), 7.55-7.57 (m, 1H), 7.76-7.79 (m, 1H), 7.82-7.84 (m, 2H), 8.01-8.03 (m, 1H), 8.16-8.17 (m, 1H), 8.36 (s, 1H), 8.43 (s, 1H), 8.58 (s, 1H)

5-Ethoxy-6-morpholin-4-yl-2-[3-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, Aetone-d₆): δ 1.32-1.34 (m, 3H), 1.38-1.41 (m, 3H), 3.81-3.83 (m, 4H), 3.89-3.91 (m, 4H), 4.02-4.05 (m, 2H), 4.41-4.44 (m, 2H), 6.99 (m, 1H), 7.26-7.30 (m, 2H), 7.55-7.57 (m, 2H), 7.84-7.86 (m, 1H), 7.98-8.00 (m, 1H), 8.15 (s, 1H), 8.27 (s, 1H), 8.34-8.36 (m, 1H)

2-{4-[3-(4-Chloro-3-trifluoromethyl-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, Aetone-d₆): δ 1.33-1.42 (m, 6H), 3.83-3.85 (m, 4H), 3.90-3.92 (m, 4H), 4.03-4.05 (m, 2H), 4.41-4.45 (m, 2H), 7.58-7.60 (m, 1H), 7.64-7.66 (m, 2H), 7.78-7.80 (m, 1H), 8.18 (m, 1H), 8.31-8.32 (m, 2H), 8.52 (s, 1H), 8.63 (s, 1H)


5-Ethoxy-2-[4-[3-(3-fluoro-phenyl)-ureido]-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



49


¹H NMR (500 MHz, Aetone-d₆): δ 1.32-1.34 (m, 3H), 1.37-1.40 (m, 3H), 3.81-3.83 (m, 4H), 3.88-3.90 (m, 4H), 4.00-4.02 (m, 2H), 4.40-4.42 (m, 2H), 6.70-6.80 (m, 1H), 7.10-7.20 (m, 1H), 7.21-7.31 (m, 1H), 7.62-7.64 (m, 3H), 8.28-8.30 (m, 2H), 8.38 (s, 2H)

5-Ethoxy-2-{4-[3-(4-fluoro-phenyl)-ureido]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, Aetone-d₆): δ 1.30-1.35 (m, 3H), 1.38-1.41 (m, 3H), 3.81-3.84 (m, 4H), 3.88-3.90 (m, 4H), 4.01-4.04 (m, 2H), 4.40-4.44 (m, 2H), 7.06-7.10 (m, 2H), 7.56-7.64 (m, 4H), 8.22-8.34 (m, 3H)

2-{4-[3-(3,4-Difluoro-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, Aetone-d₆): δ 1.30-1.35 (m, 3H), 1.38-1.41 (m, 3H), 3.82-3.84 (m, 4H), 3.89-3.91 (m, 4H), 4.00-4.04 (m, 2H), 4.40-4.44 (m, 2H), 7.20-7.25 (m, 2H), 7.62-7.64 (m, 2H), 7.80-7.90 (m, 1H), 8.29-8.30 (m, 2H), 8.38 (s, 1H)


2-{4-[3-(6-Chloro-pyridin-3-yl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



50

¹H NMR (500 MHz, Aetone-d₆): δ 1.31-1.40 (m, 6H), 3.80-3.82 (m, 4H), 3.88-3.90 (m, 4H), 4.00-4.03 (m, 2H), 4.39-4.43 (m, 2H), 7.37-7.39 (m, 1H), 7.62-7.64 (m, 2H), 8.11-8.13 (m, 1H), 8.28-8.30 (m, 2H), 8.47-8.49 (m, 1H)


5-Ethoxy-6-morpholin-4-yl-2-(4-ureido-phenyl)-pyrimidine-4-carboxylic acid ethyl ester



Trimethylsilyl isocyanate (0.1 ml, 5 eq.) was added to a stirred solution of 2-(4-amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (60 mg, 1 eq.) in THF (2 ml) and the mixture was reacted to reflux overnight. The reaction mixture was extracted with EA and washed with brine. The crude was purified by chromatography to give a product (26.2 mg, 40%).

¹H NMR (500 MHz, Aetone-d₆): δ 1.31-1.40 (m, 6H), 3.80-3.82 (m, 4H), 3.86-3.88 (m, 4H), 3.99-4.02 (m, 2H), 4.38-4.42 (m, 2H), 5.55 (s, 2H), 7.57-7.59 (m, 2H), 8.22-8.24 (m, 2H), 8.31 (s, 1H)


5-Ethoxy-6-morpholin-4-yl-2-[6-(3-phenyl-ureido)-pyridin-3-yl]-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, Aetone-d₆): δ 1.33-1.36 (m, 3H), 1.39-1.41 (m, 3H), 3.82-3.84 (m, 4H), 3.92-3.94 (m, 4H), 4.02-4.04 (m, 2H), 4.42-4.44 (m, 2H), 7.00-7.10 (m, 1H), 7.31-7.34 (m, 3H), 8.55-8.65 (m, 1H), 8.90-9.00 (s, 1H), 9.24-9.25 (m, 1H), 11.40 (s, 1H)


51

5-Ethoxy-2-{6-[3-(4-fluoro-phenyl)-ureido]-pyridin-3-yl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, Acetone-d₆): δ 1.32-1.41 (m, 6H), 3.82-3.84 (m, 4H), 3.92-3.94 (m, 4H), 4.02-4.05 (m, 2H), 4.41-4.45 (m, 2H), 7.08-7.12 (m, 2H), 7.35-7.36 (m, 1H), 7.71-7.73 (m, 2H), 8.59-8.61 (m, 1H), 8.97 (s, 1H), 9.23-9.24 (m, 1H), 11.51 (s, 1H)

2-{6-[3-(3,4-Difluoro-phenyl)-ureido]-pyridin-3-yl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, Acetone-d₆): δ 1.33-1.41 (m, 6H), 3.82-3.83 (m, 4H), 3.92-3.93 (m, 4H), 4.00-4.05 (m, 2H), 4.41-4.45 (m, 2H), 7.25-7.38 (m, 3H), 7.89-7.92 (m, 1H), 8.60-8.62 (m, 1H), 9.06 (s, 1H), 9.24 (s, 1H), 11.78 (s, 1H)

5-Ethoxy-2-{6-[3-(3-fluoro-phenyl)-ureido]-pyridin-3-yl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, Acetone-d₆): δ 1.32-1.41 (m, 6H), 3.82-3.84 (m, 4H), 3.92-3.94 (m, 4H), 4.00-4.05 (m, 2H), 4.41-

52

4.45 (m, 2H), 6.79-6.82 (m, 1H), 7.33-7.38 (m, 3H), 7.73-7.75 (m, 1H), 8.60-8.63 (m, 1H), 9.03 (s, 1H), 9.26 (s, 1H)


5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-thioureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester



Phenyl isothiocyanate (0.02 ml, 1.5 eq.) was added to a stirred solution of 2-(4-amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (40 mg, 1 eq.) in CHCl₃ (3 ml) and the mixture was reacted overnight at r.t. The reaction mixture was extracted with EA and washed with brine. The crude was purified by chromatography to give a product (32.6 mg, 60%).

¹H NMR (500 MHz, Acetone-d₆): δ 1.32-1.40 (m, 6H), 3.81-3.83 (m, 4H), 3.89-3.91 (m, 4H), 4.01-4.04 (m, 2H), 4.39-4.43 (m, 2H), 7.16-7.19 (m, 1H), 7.35-7.38 (m, 2H), 7.55-7.57 (m, 2H), 7.67-7.70 (m, 2H), 8.30-8.32 (m, 2H), 9.19 (s, 1H), 9.23 (s, 1H)

2-(4-Benzenesulfonylamino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




Phenyl sulfonyl chloride (0.02 ml, 1.3 eq.), Et₃N (1 drop) was added to a stirred solution of 2-(4-amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (40 mg, 1 eq.) in CH₂Cl₂ (2 ml) and the mixture was reacted overnight at r.t. The reaction mixture was extracted with EA and washed with brine. The crude was purified by chromatography to give a product (13 mg, 23.7%).

¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.36 (m, 3H), 1.40-1.44 (m, 3H), 3.81-3.87 (m, 8H), 4.10-4.14 (m, 2H), 4.43-4.47 (m, 2H), 6.79 (s, 1H), 7.12-7.13 (m, 2H), 7.41-7.44 (m, 2H), 7.51-7.53 (m, 1H), 7.77-7.79 (m, 2H), 8.20-8.22 (m, 2H)

53


5-Ethoxy-2-(4-ethylcarbamoyloxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



Ethyl isocyanate (0.013 ml, 1.5 eq.) was added to a stirred solution of 2-(4-hydroxy-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (40 mg, 1 eq.) in toluene (3 ml) and the mixture was reacted to reflux overnight. The reaction mixture was extracted with EA and washed with brine. The crude was purified by chromatography to give a product (12 mg, 25.2%).


¹H NMR (500 MHz, CDCl₃-d₁): δ 1.23-1.27 (m, 3H), 1.34-1.37 (m, 3H), 1.39-1.45 (m, 3H), 3.31-3.34 (m, 2H), 3.82-3.83 (m, 4H), 3.88-3.89 (m, 4H), 3.96-4.11 (m, 2H), 4.44-4.49 (m, 2H), 5.07 (s, 1H), 7.17-7.19 (m, 2H), 8.31-8.33 (m, 2H)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenylureido)-loxy-phenyl]-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, Acetone-d₆): δ 1.32-1.40 (m, 6H), 3.81-3.83 (m, 4H), 3.90-3.92 (m, 4H), 4.02-4.05 (m, 2H), 4.40-4.44 (m, 2H), 7.07-7.10 (m, 1H), 7.30-7.37 (m, 4H), 7.62-7.64 (m, 2H), 8.39-8.40 (m, 2H), 9.28 (s, 1H)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenylureido)-phenyl]-pyrimidine-4-carboxylic acid amide




5-ethoxy-6-morpholin-4-yl-2-[4-(3-phenylureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (50 mg, 1 eq.), 7N NH₃/MeOH (9 ml) and THF (0.5 ml) were placed in high

54

pressure reacted bottle and reacted to reflux for 2 days. The reaction mixture was cooled, solvent removed in vacuo, and the residue washed with heating MeOH and filtered to give a product (37.7 mg, 80%).

¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 3.75-3.76 (m, 4H), 3.80-3.81 (m, 4H), 3.92-3.96 (m, 2H), 6.97-6.99 (m, 1H), 7.27-7.31 (m, 2H), 7.46-7.47 (m, 2H), 7.54-7.55 (m, 2H), 7.60 (s, 1H), 7.95 (s, 1H), 8.21-8.3 (m, 2H), 8.70 (s, 1H), 8.90 (s, 1H)


5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenylureido)-phenyl]-pyrimidine-4-carboxylic acid



1N KOH (10.5 ml, 5 eq.) was added to a stirred solution of 5-ethoxy-6-morpholin-4-yl-2-[4-(3-phenylureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (1.03 g, 1 eq.) in MeOH (10 ml) and the mixture was refluxed for 4 h. The mixture was cooled to 0° C. and carefully acidified with conc. HCl. The mixture was filtered and the precipitate was dried in vacuo, and a product was obtained as a white solid (0.93 g, 97%).

¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 3.75-3.76 (m, 4H), 3.81-3.82 (m, 4H), 3.90-3.95 (m, 2H), 6.96-6.99 (m, 1H), 7.27-7.30 (m, 2H), 7.46-7.48 (m, 2H), 7.55-7.57 (m, 2H), 8.17-8.19 (m, 2H), 9.09 (s, 1H), 9.29 (s, 1H)

Potassium Salt of 5-ethoxy-6-morpholin-4-yl-2-[4-(3-phenylureido)-phenyl]-pyrimidine-4-carboxylic acid




1N KOH (5 ml, 5 eq.) was added to a stirred solution of 5-ethoxy-6-morpholin-4-yl-2-[4-(3-phenylureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (500 mg, 1 eq.) in MeOH (5 ml) and the mixture was refluxed for 4 h. The mixture was cooled to r.t. and filtered. The precipitate was

55

washed with MeOH and dried in vacuo, and a product was obtained as a white solid (400 mg, 78%).

¹H NMR (500 MHz, DMSO-d₆): δ 1.23-1.26 (m, 3H), 3.74 (m, 8H), 4.01-4.03 (m, 2H), 6.94-6.95 (m, 1H), 7.25-7.28 (m, 2H), 7.55-7.57 (m, 2H), 7.69-7.70 (m, 2H), 8.19-8.20 (m, 2H), 10.17 (s, 1H), 10.30 (s, 1H)


5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid diethylamide



EDC (50 mg, 1.5 eq.), HOBT (40 mg, 1.5 eq.), Et₂NH (0.0275 ml, 1.5 eq.) and Et₃N (0.05 ml, 2.0 eq.) were added to a stirred solution of 5-ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid (80 mg, 1 eq.) in dry DMF (2 ml) and the mixture reacted overnight at r.t. The mixture was quenched with H₂O, extracted with EA, and washed with brine. The crude was purified by chromatography to give a product (43 mg, 48%).

¹H NMR (500 MHz, DMSO-d₆): δ 1.07-1.10 (m, 3H), 1.15-1.18 (m, 3H), 1.22-1.24 (m, 3H), 3.14-3.16 (m, 2H), 3.45-3.46 (m, 2H), 3.76-3.81 (m, 4H), 3.87-3.90 (m, 4H), 4.02-4.03 (m, 2H), 6.98-6.99 (m, 1H), 7.27-7.30 (m, 2H), 7.45-7.47 (m, 2H), 7.54-7.56 (m, 2H), 8.17-8.19 (m, 2H), 8.71 (s, 1H), 8.90 (s, 1H)


1-{4-[5-Ethoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-phenyl-urea



¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.27 (m, 3H), 3.28-3.30 (m, 2H), 3.56-3.57 (m, 2H), 3.67 (m, 4H), 3.76-3.82 (m, 8H), 3.90-3.91 (m, 2H), 6.98 (m, 1H), 7.28-7.31 (m, 2H), 7.46-7.47 (m, 2H), 7.54-7.56 (m, 2H), 8.17-8.18 (m, 2H), 8.71 (s, 1H), 8.90 (s, 1H)


56

1-{4-[5-Ethoxy-4-morpholin-4-yl-6-(pyrrolidine-1-carbonyl)-pyrimidin-2-yl]-phenyl}-3-phenyl-urea




¹H NMR (500 MHz, DMSO-d₆): δ 1.21-1.24 (m, 3H), 1.84-1.88 (m, 4H), 3.25-3.27 (m, 2H), 3.47-3.49 (m, 2H), 3.75-3.81 (m, 8H), 3.90-3.91 (m, 2H), 6.98 (m, 1H), 7.29-7.30 (m, 2H), 7.45-7.47 (m, 2H), 7.53-7.55 (m, 2H), 8.16-8.18 (m, 2H), 8.70 (s, 1H), 8.90 (s, 1H)

1-{4-[5-Ethoxy-4-morpholin-4-yl-6-(piperidine-1-carbonyl)-pyrimidin-2-yl]-phenyl}-3-phenyl-urea



¹H NMR (500 MHz, DMSO-d₆): δ 1.23-1.26 (m, 3H), 1.50-1.64 (m, 6H), 3.20 (m, 2H), 3.62 (m, 2H), 3.76-3.80 (m, 8H), 3.87-3.91 (m, 2H), 6.97-6.99 (m, 1H), 7.27-7.30 (m, 2H), 7.45-7.47 (m, 2H), 7.53-7.55 (m, 2H), 8.16-8.18 (m, 2H), 8.76 (s, 1H), 8.95 (s, 1H)


1-{4-[5-Ethoxy-4-(4-methanesulfonyl-piperazine-1-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-phenyl-urea



57


¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.27 (m, 3H), 2.95 (s, 3H), 3.13 (m, 2H), 3.23 (m, 2H), 3.40 (m, 2H), 3.77-3.82 (m, 10H), 3.88-3.92 (m, 2H), 6.98 (m, 1H), 7.27-7.30 (m, 2H), 7.46-7.47 (m, 2H), 7.54-7.55 (m, 2H), 8.17-8.19 (m, 2H), 8.71 (s, 1H), 8.91 (s, 1H)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid (2-morpholin-4-yl-ethyl)-amide




¹H NMR (500 MHz, DMSO-d₆): δ 1.23-1.27 (m, 6H), 2.36-2.63 (m, 6H), 3.3-3.41 (m, 4H), 3.59-3.60 (m, 4H), 3.74-3.75 (m, 4H), 3.81-3.82 (m, 4H), 3.92-3.95 (m, 2H), 6.97-6.99 (m, 1H), 7.27-7.30 (m, 2H), 7.46-7.48 (m, 2H), 7.54-7.56 (m, 2H), 8.22-8.24 (m, 2H), 8.54-8.57 (m, 1H), 8.70 (s, 1H), 8.91 (s, 1H)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid (2-pyrrolidin-1-yl-ethyl)-amide



¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.27 (m, 6H), 1.70 (m, 4H), 2.57-2.60 (m, 4H), 3.75-3.76 (m, 4H), 3.81-3.82 (m, 4H), 3.90-3.95 (m, 2H), 6.96-6.99 (m, 1H), 7.27-7.30 (m, 2H), 7.47-7.49 (m, 2H), 7.56-7.57 (m, 2H), 8.20-8.22 (m, 2H), 8.55-8.57 (m, 1H), 9.10 (s, 1H), 9.30 (s, 1H)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid (2-diethylamino-ethyl)-amide




¹H NMR (500 MHz, DMSO-d₆): δ 0.98-1.01 (m, 6H), 1.24-1.27 (m, 3H), 2.54-2.58 (m, 8H), 3.75-3.76 (m, 4H),

58


3.81-3.82 (m, 4H), 3.92-3.95 (m, 2H), 6.98 (m, 1H), 7.27-7.30 (m, 2H), 7.46-7.48 (m, 2H), 7.55-7.56 (m, 2H), 8.21-8.23 (m, 2H), 8.50 (m, 1H), 9.00 (s, 1H), 9.20 (s, 1H)

5-Methoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, CDCl₃-d₁): δ 1.42-1.44 (m, 3H), 3.76 (s, 3H), 3.81-3.873 (m, 8H), 4.45-4.50 (m, 2H), 7.05-7.07 (m, 1H), 7.12 (s, 1H), 7.22 (m, 1H), 7.27-7.33 (m, 4H), 7.37-7.39 (m, 2H), 8.20-8.21 (m, 2H)

5-Methoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid



¹H NMR (500 MHz, DMSO-d₆): δ 3.72 (s, 3H), 3.77-3.82 (m, 8H), 6.99-7.00 (m, 1H), 7.28-7.30 (m, 2H), 7.46-7.48 (m, 2H), 7.55-7.57 (m, 2H), 8.18-8.19 (m, 2H), 8.78 (s, 1H), 8.98 (s, 1H)


1-[4-[5-Methoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl]-3-phenyl-urea



59


¹H NMR (500 MHz, DMSO-d₆): δ 3.28 (m, 2H), 3.33 (m, 4H), 3.56 (m, 2H), 3.67-3.68 (m, 7H), 3.76-3.82 (m, 8H), 6.98-6.99 (m, 1H), 7.28-7.30 (m, 2H), 7.46-7.47 (m, 2H), 7.54-7.56 (m, 2H), 8.17-8.19 (m, 2H), 8.71 (s, 1H), 8.91 (s, 1H)

1-{4-[4-(4-Methanesulfonyl-piperazine-1-carbonyl)-5-methoxy-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-phenyl-urea




¹H NMR (500 MHz, DMSO-d₆): δ 2.95 (s, 3H), 3.12 (m, 2H), 3.24 (m, 2H), 3.39 (m, 2H), 3.68 (s, 3H), 3.77-3.82 (m, 10H), 6.97-6.99 (m, 1H), 7.27-7.30 (m, 2H), 7.46-7.47 (m, 2H), 7.54-7.56 (m, 2H), 8.17-8.19 (m, 2H), 8.71 (s, 1H), 8.91 (s, 1H)

1-{4-[5-Ethoxy-4-(4-methyl-piperazine-1-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-phenyl-urea



¹H NMR (500 MHz, DMSO-d₆): δ 1.23-1.27 (m, 6H), 2.20 (s, 3H), 2.29 (m, 2H), 2.37-2.38 (m, 2H), 3.24-3.26 (m, 2H), 3.65 (m, 2H), 3.76-3.81 (m, 8H), 3.87-3.91 (m, 2H), 6.97-6.99 (m, 1H), 7.27-7.30 (m, 2H), 7.46-7.47 (m, 2H), 7.54-7.56 (m, 2H), 8.16-8.18 (m, 2H), 8.731 (s, 1H), 8.93 (s, 1H)


5-Ethoxy-6-morpholin-4-yl-2-(4-phenoxy-carbonylamino-phenyl)-pyrimidine-4-carboxylic acid ethyl ester



60

Phenyl chloroformate (0.95 ml, 1.5 eq.) was added to a stirred solution of 2-(4-Amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (1.15 g, 1 eq.) in EA (20 ml) and NaHCO₃(sat) (20 ml). The mixture was reacted at room temperature for 2 hrs. The reaction mixture was diluted with NaHCO₃(sat) and extracted with EA. The organic solution was washed with brine, was dried (MgSO₄), filtered and concentrated under reduced pressure to give a crude carbamate (1.21 g, 80%).


5-Ethoxy-2-[4-(3-methyl-ureido)-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



2M Methyl amine (1.1 ml, 10 eq.) was added to a stirred solution of 5-ethoxy-6-morpholin-4-yl-2-(4-phenoxy-carbonylamino-phenyl)-pyrimidine-4-carboxylic acid ethyl ester (110 mg, 1 eq.) in dioxane (3 ml), and the mixture was reacted at 80° C. overnight. The reaction mixture was concentrated and purified by flash chromatography to give a product (47.5 mg, 50%).

¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.35 (m, 3H), 1.40-1.43 (m, 3H), 2.73-2.75 (m, 3H), 3.81-3.82 (m, 4H), 3.86-3.87 (m, 4H), 3.94-3.98 (m, 2H), 4.43-4.48 (m, 2H), 5.25 (s, 1H), 7.2 (s, 1H), 7.36-7.37 (m, 2H), 8.21-8.22 (m, 2H)

5-Ethoxy-2-{4-[(morpholine-4-carbonyl)-amino]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



Morpholine (0.071 ml, 4 eq.) was added to a stirred solution of 5-ethoxy-6-morpholin-4-yl-2-(4-phenoxy-carbonylamino-phenyl)-pyrimidine-4-carboxylic acid ethyl ester (100 mg, 1 eq.) and Et₃N (0.085 ml, 3 eq.) in dioxane (3 ml), the mixture was reacted at 80° C. overnight. The reaction


61

mixture was concentrated, and purified by flash chromatography to give a product (59 mg, 60%).

¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.36 (m, 3H), 1.42-1.44 (m, 3H), 3.49-3.51 (m, 4H), 3.74-3.75 (m, 4H), 3.83-3.84 (m, 4H), 3.87-3.88 (m, 4H), 3.95-3.99 (m, 2H), 4.44-4.48 (m, 2H), 6.45 (s, 1H), 7.42-7.43 (m, 2H), 8.26-8.28 (m, 2H)


The urea compounds of the following examples were synthesized following the synthetic method described above by the appropriate amine with 5-ethoxy-6-morpholin-4-yl-2-(4-phenoxycarbonylamino-phenyl)-pyrimidine-4-carboxylic acid ethyl ester.

2-{4-[3-(3-Dimethylamino-propyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.36 (m, 3H), 1.41-1.44 (m, 3H), 1.66-1.69 (m, 2H), 2.20-2.22 (s, 6H), 2.39-2.41 (m, 2H), 3.34 (s, 2H), 3.81-3.83 (m, 4H), 3.87-3.89 (m, 4H), 3.96-3.99 (m, 2H), 4.43-4.47 (m, 2H), 7.40-7.42 (m, 2H), 8.24-8.26 (m, 2H)


5-Ethoxy-6-morpholin-4-yl-2-[4-(3-pyridin-3-yl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.27 (m, 3H), 1.32-1.35 (m, 3H), 3.75-3.76 (m, 4H), 3.82-3.83 (m, 4H), 3.89-3.93 (m, 2H), 4.35-4.40 (m, 2H), 7.32-7.34 (m, 1H), 7.56-7.58 (m, 2H), 7.95-7.97 (m, 1H), 8.17-8.21 (m, 3H), 8.61 (s, 1H), 8.91 (s, 1H), 9.01 (s, 1H)

62


5-Ethoxy-2-{4-[3-(4-methyl-piperazin-1-yl)-ureido]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



4-methyl-piperazin-1-ylamine (0.52 ml, 2 eq.) was added to a stirred solution of 5-ethoxy-6-morpholin-4-yl-2-(4-phenoxycarbonylamino-phenyl)-pyrimidine-4-carboxylic acid ethyl ester (100 mg, 1 eq.) and Et₃N (0.085 ml, 3 eq.) in dioxane (3 ml), the mixture was reacted at 80° C. overnight. The reaction mixture was concentrated, and purified by flash chromatography to give a product (49.2 mg, 45%).


¹H NMR (500 MHz, CDCl₃-d₁): δ 1.32-1.35 (m, 3H), 1.41-1.44 (m, 3H), 2.34 (s, 3H), 2.46-2.49 (m, 4H), 3.54-3.56 (m, 4H), 3.81-3.83 (m, 4H), 3.87-3.88 (m, 4H), 4.10-4.12 (m, 2H), 4.44-4.46 (m, 2H), 6.67 (s, 1H), 7.41-7.43 (m, 2H), 8.24-8.25 (m, 2H)

5-Ethoxy-6-morpholin-4-yl-2-{4-[3-(2-morpholin-4-yl-ethyl)-ureido]-phenyl}-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.36 (m, 3H), 1.41-1.44 (m, 3H), 2.48 (s, 4H), 2.52-2.54 (m, 2H), 3.36-3.38 (m, 2H), 3.70 (s, 4H), 3.82-3.83 (m, 4H), 3.87-3.89 (m, 4H), 3.96-3.99 (m, 2H), 4.43-4.47 (m, 2H), 5.34 (s, 1H), 7.39-7.41 (m, 2H), 8.26-8.28 (m, 2H)


5-Ethoxy-6-morpholin-4-yl-2-[4-(3-morpholin-4-yl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester



63


¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.35 (m, 3H), 1.41-1.44 (m, 3H), 3.49-3.51 (m, 4H), 3.74-3.75 (m, 4H), 3.81-3.83 (m, 4H), 3.87-3.88 (m, 4H), 3.95-3.99 (m, 2H), 4.43-4.48 (m, 2H), 6.50 (s, 1H), 7.41-7.43 (m, 2H), 8.25-8.27 (m, 2H)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-thiazol-2-yl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.75-3.76 (m, 4H), 3.82-3.83 (m, 4H), 3.89-3.93 (m, 2H), 4.35-4.40 (m, 2H), 7.13 (s, 1H), 7.38-7.39 (m, 1H), 7.58-7.60 (m, 2H), 8.19-8.21 (m, 2H), 9.21 (s, 1H)

5-Ethoxy-2-{4-[3-(3-fluoro-4-morpholin-4-yl-phenyl)-ureido]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.36 (m, 3H), 1.42-1.45 (m, 3H), 2.99-3.01 (m, 4H), 3.82-3.84 (m, 4H), 3.85 (m, 4H), 3.87-3.88 (m, 4H), 3.95-3.99 (m, 2H), 4.45-4.50 (m, 2H), 6.81-6.84 (m, 1H), 6.94-6.97 (m, 2H), 7.07 (s, 1H), 7.17-7.20 (m, 1H), 7.35-7.37 (m, 2H), 8.21-8.23 (m, 2H)

5-Ethoxy-6-morpholin-4-yl-2-(4-{3-[4 (3 oxo-morpholin-4-yl)-phenyl]-ureido}-phenyl)-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.70-3.71 (m, 2H), 3.75-3.76 (m, 4H),

64


3.82-3.83 (m, 4H), 3.89-3.93 (m, 2H), 3.96 (m, 2H), 4.18 (m, 2H), 4.35-4.40 (m, 2H), 7.29-7.31 (2H), 7.48-7.50 (m, 2H), 7.55-7.57 (m, 2H), 8.17-8.18 (m, 2H), 8.24 (s, 1H), 8.97 (s, 1H)

5-Ethoxy-6-morpholin-4-yl-2-{4-[3-(3-trifluoromethyl-phenyl)-ureido]-phenyl}-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.36 (m, 3H), 1.42-1.45 (m, 3H), 3.80-3.81 (m, 4H), 3.86 (m, 4H), 3.94-3.98 (m, 2H), 4.47-4.51 (m, 2H), 7.20-7.22 (m, 1H), 7.29-7.32 (m, 3H), 7.53-7.55 (m, 3H), 7.70 (s, 1H), 8.11-8.12 (m, 2H)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-pyridin-2-yl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.76 (m, 4H), 3.82-3.83 (m, 4H), 3.91-3.92 (m, 2H), 4.38-4.39 (m, 2H), 7.04 (m, 1H), 7.53-7.54 (m, 1H), 7.62-7.63 (m, 2H), 7.77 (m, 1H), 8.19-8.20 (m, 2H), 8.30 (m, 1H), 9.49 (s, 1H), 10.68 (s, 1H)


5-Ethoxy-2-(4-{3-[4-(4-methyl-piperazin-1-yl)-phenyl]-ureido}-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



65


¹H NMR (500 MHz, DMSO-d₆): δ 1.21-1.24 (m, 3H), 1.28-1.31 (m, 3H), 2.17 (s, 3H), 2.41 (m, 4H), 3.01 (s, 4H), 3.71 (m, 4H), 3.77 (m, 4H), 3.86-3.87 (m, 2H), 4.33-4.34 (m, 2H), 6.83-6.85 (m, 2H), 7.25-7.27 (m, 2H), 7.48-7.50 (m, 2H), 8.10-8.12 (m, 2H), 8.43 (s, 1H), 8.81 (s, 1H)

5-Ethoxy-6-morpholin-4-yl-2-[4-[3-(4-thiomorpholin-4-yl-phenyl)-ureido]-phenyl]-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.28 (m, 3H), 1.32-1.35 (m, 3H), 2.67-2.69 (m, 4H), 3.38-3.40 (m, 4H), 3.01 (m, 4H), 3.75-3.76 (m, 4H), 3.81-3.82 (m, 4H), 3.89-3.93 (m, 2H), 4.36-4.40 (m, 2H), 6.88-6.89 (m, 2H), 7.31-7.32 (m, 2H), 7.53-7.55 (m, 2H), 8.15-8.16 (m, 2H), 8.48 (s, 1H), 8.85 (s, 1H)


5-Ethoxy-6-morpholin-4-yl-2-[4-[3-(4-morpholin-4-yl-phenyl)-ureido]-phenyl]-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.01-3.03 (m, 4H), 3.72-3.75 (m, 8H), 3.81-3.82 (m, 4H), 3.89-3.93 (m, 2H), 4.36-4.40 (m, 2H), 6.88-6.90 (m, 2H), 7.31-7.32 (m, 2H), 7.52-7.54 (m, 2H), 8.15-8.16 (m, 2H), 8.47 (s, 1H), 8.84 (s, 1H)

66


2-[4-[3-(2-Amino-phenyl)-thioureido]-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



To a stirred solution of 1,1-thiocarbonyldiimidazole (216.5 mg, 1.5 eq.), imidazole (164.52 mg, 3 eq.) and CH₃CN (10 ml) at 0° C., 2-(4-amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (300 mg, 1 eq.) dissolved in CH₃CN (10 ml) was added dropwise over 10 min. After 10 min, the cooling bath was removed. After 3 h, benzene-1,2-diamine (175.2 mg, 2 eq.) was added and the reaction was heated to 50° C. for 3 h, and then stirred at ambient temperature for 16 h. The solvent was removed by evaporation. The residue was chromatographed to give a thiourea (120 mg, 29%)

¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.75-3.76 (m, 4H), 3.82-3.83 (m, 4H), 3.89-3.93 (m, 2H), 4.36-4.40 (m, 2H), 6.56-6.59 (m, 1H), 6.75-6.76 (m, 1H), 6.96-6.99 (m, 1H), 7.09-7.10 (m, 1H), 7.68-7.69 (m, 2H), 8.17-8.19 (m, 2H), 9.17 (s, 1H), 9.82 (s, 1H)

2-[4-(1H-Benzoimidazol-2-ylamino)-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




A solution of thiourea (120 mg, 1 eq.) in THF (5 ml) and DCC (56.85 mg, 1.2 eq.) was heated under reflux with stirring for 4 h. The cooled solution was stirred overnight and filtered to give a product (33.7 mg, 30%).

¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.75-3.77 (m, 4H), 3.82-3.83 (m, 4H), 3.89-3.93 (m, 2H), 4.36-4.40 (m, 2H), 7.00-7.03 (m, 2H), 7.30-7.32 (m, 2H), 7.37-7.39 (m, 2H), 7.85-7.86 (m, 2H), 8.20-8.21 (m, 2H), 9.76 (s, 1H), 11.00 (s, 1H)


67

2-{4-[3-(4-Carbamoyl-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.76 (m, 4H), 3.81-3.82 (m, 4H), 3.91-3.92 (m, 2H), 4.36-4.40 (m, 2H), 7.21 (m, 1H), 7.52-7.58 (m, 4H), 7.82-7.84 (m, 3H), 8.18-8.19 (m, 2H), 8.99 (s, 1H), 9.05 (s, 1H)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-pyridin-3-yl-ureido)-phenyl]-pyrimidine-4-carboxylic acid



¹H NMR (500 MHz, DMSO-d₆): δ 1.26-1.28 (m, 3H), 3.75-3.76 (m, 4H), 3.81-3.82 (m, 4H), 3.91-3.95 (m, 2H), 4.37-4.39 (m, 2H), 7.56-7.60 (m, 2H), 7.89 (m, 1H), 8.18-8.23 (m, 2H), 8.32-8.34 (m, 1H), 8.48-8.49 (m, 1H), 9.08 (m, 1H), 9.77 (s, 1H), 10.23 (s, 1H)


2-{4-[3-(3-Acetylamino-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 2.04 (s, 3H), 3.75-3.76 (m, 4H), 3.81-3.82 (m, 4H), 3.89-3.93 (m, 2H), 4.36-4.40 (m, 2H), 7.17-7.21 (m, 3H), 7.54-7.56 (m, 2H), 7.79 (m, 1H), 8.16-8.18 (m, 2H), 8.76 (s, 1H), 8.85 (s, 1H), 9.94 (s, 1H)


68

2-{4-[3-(3-Carbamoyl-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.75-3.76 (m, 4H), 3.81-3.82 (m, 4H), 3.90-3.92 (m, 2H), 4.37-4.38 (m, 2H), 7.35 (m, 2H), 7.47-7.48 (m, 1H), 7.56-7.58 (m, 2H), 7.63-7.64 (m, 1H), 7.91-7.94 (m, 2H), 9.17-8.19 (m, 2H), 8.86 (s, 1H), 8.99 (s, 1H)

5-Ethoxy-6-morpholin-4-yl-2-(4-{3-[3-(2,2,2-trifluoroacetyl-amino)-phenyl]-ureido}-phenyl)-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.28 (m, 3H), 1.32-1.35 (m, 3H), 2.67-2.69 (m, 4H), 3.38-3.40 (m, 4H), 3.01 (m, 4H), 3.75-3.76 (m, 4H), 3.81-3.82 (m, 4H), 3.89-3.93 (m, 2H), 4.36-4.40 (m, 2H), 7.57-7.59 (m, 3H), 7.72-7.74 (m, 1H), 7.84-7.85 (m, 1H), 8.18-8.20 (m, 2H), 8.58 (s, 1H), 9.11 (s, 1H), 9.27 (s, 1H)


1-{4-[5-Ethoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-pyridin-3-yl-urea



69


¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.27 (m, 3H), 3.28-3.29 (m, 2H), 3.55-3.57 (m, 2H), 3.66 (m, 4H), 3.76 (m, 4H), 3.81 (m, 4H), 3.88-3.92 (m, 2H), 7.32-7.35 (m, 1H), 7.55-7.57 (m, 2H), 7.95-7.97 (m, 1H), 8.17-8.21 (m, 3H), 8.61-8.62 (m, 1H), 8.95 (s, 1H), 9.10 (s, 1H)

5-Ethoxy-6-morpholin-4-yl-2-{4-[3-(4-sulfamoyl-phenyl)-ureido]-phenyl}-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.75-3.76 (m, 4H), 3.81-3.82 (m, 4H), 3.89-3.93 (m, 2H), 4.36-4.40 (m, 2H), 7.22 (s, 1H), 7.56-7.58 (m, 2H), 7.62-7.63 (m, 2H), 7.73-7.75 (m, 2H), 8.18-8.19 (m, 2H), 9.12 (s, 1H), 9.18 (s, 1H)

2-{4-[3-(4-Carboxy-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.75-3.76 (m, 4H), 3.81-3.82 (m, 4H), 3.89-3.93 (m, 2H), 4.36-4.40 (m, 2H), 7.56 (m, 4H), 7.86-7.88 (2H), 8.16-8.18 (m, 2H), 9.26 (s, 1H), 9.33 (s, 1H)

5-Ethoxy-2-(4-{3-[4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ureido}-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




EDC (53.7 mg, 1.5 eq.), HOBt (42.9 mg, 1.5 eq.), 1-methyl-piperazine (0.03 ml, 1.5 eq.) and Et₃N (0.053 ml, 2.0 eq.)

70

were added to a stirred solution of 2-{4-[3-(4-carboxy-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (80 mg, 1 eq.) in dry DMF (2 ml) and the mixture reacted overnight at r.t. The mixture was quenched with H₂O, extracted with EA, and washed with brine. The crude was purified by chromatography to give a product (34.5 mg, 30%).


¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 2.19 (s, 3H), 2.31 (m, 4H), 3.50 (m, 4H), 3.75-3.76 (m, 4H), 3.82 (m, 4H), 3.89-3.93 (m, 2H), 4.35-4.40 (m, 2H), 7.32-7.34 (m, 2H), 7.52-7.57 (m, 4H), 8.16-8.18 (m, 2H), 9.27 (s, 1H), 9.29 (s, 1H).

5-Ethoxy-2-(4-{3-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-ureido}-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 2.22 (s, 3H), 2.42 (m, 4H), 3.40 (m, 4H), 3.75-3.76 (m, 4H), 3.81-3.82 (m, 4H), 3.89-3.92 (m, 2H), 4.35-4.39 (m, 2H), 6.81-6.82 (m, 1H), 7.53-7.54 (m, 2H), 7.69-7.70 (m, 1H), 8.14-8.16 (m, 3H), 8.52 (s, 1H), 8.99 (s, 1H)


5-Ethoxy-6-morpholin-4-yl-2-{4-[3-(6-morpholin-4-yl-pyridin-3-yl)-ureido]-phenyl}-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.71 (m, 4H), 3.75 (m, 4H), 3.82 (m, 4H), 3.89-3.91 (m, 2H), 4.35-4.40 (m, 2H), 6.82-6.84 (m, 1H), 7.53-7.55 (m, 2H), 7.72-7.74 (m, 1H), 8.15-8.18 (m, 3H), 8.51 (s, 1H), 8.96 (s, 1H)


71

2-(4-{3-[4-(2 Dimethylamino-ethylcarbamoyl)-phenyl]-ureido}-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 2.17 (s, 6H), 2.37-2.39 (m, 2H), 3.75-3.76 (m, 4H), 3.82 (m, 4H), 3.89-3.93 (m, 2H), 4.35-4.40 (m, 2H), 7.54-7.59 (m, 4H), 7.77-7.79 (m, 2H), 8.16-8.18 (m, 3H), 9.5 (b, 2H)

5-Ethoxy-2-(4-{3-[4-(morpholine-4-carbonyl)-phenyl]-ureido}-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.50 (m, 4H), 3.59 (m, 4H), 3.75-3.76 (m, 4H), 3.82 (m, 4H), 3.89-3.93 (m, 2H), 4.35-4.40 (m, 2H), 7.35-7.38 (m, 2H), 7.52-7.57 (m, 4H), 8.17-8.18 (m, 2H), 8.95 (s, 1H), 8.99 (s, 1H)

4-[5-Ethoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl-carbamic acid phenyl ester




¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.27 (m, 3H), 3.28-3.30 (m, 2H), 3.55-3.57 (m, 2H), 3.66 (s, 4H), 3.75-3.76

72


(m, 4H), 3.80-3.82 (m, 2H), 3.88-3.92 (m, 2H), 7.24-7.29 (m, 3H), 7.42-7.46 (m, 2H), 7.60-7.62 (m, 2H), 8.20-8.22 (m, 2H), 10.45 (s, 1H)

5 1-[4-[5-Ethoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.27 (m, 3H), 3.29 (m, 2H), 3.56 (m, 2H), 3.67-3.69 (m, 6H), 3.75-3.96 (m, 12H), 4.18 (s, 2H), 7.28-7.30 (m, 2H), 7.47-7.50 (m, 2H), 7.54-7.56 (m, 2H), 8.16-8.18 (m, 2H), 8.80 (s, 1H), 8.94 (s, 1H)

5-Ethoxy-2-(4-{3-[5-(4-methyl-piperazin-1-yl)-pyridin-2-yl]-ureido}-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 2.22 (s, 3H), 2.46 (m, 4H), 3.10 (s, 4H), 3.75-3.76 (m, 4H), 3.81-3.83 (m, 4H), 3.89-3.93 (m, 2H), 4.35-4.39 (m, 2H), 7.44-7.46 (m, 2H), 7.59-7.60 (m, 2H), 7.96 (s, 1H), 8.17-8.19 (m, 2H), 9.21 (s, 1H), 10.5 (b, 1H)


5-Ethoxy-2-[4-[3-(2-methoxycarbonyl-thiophen-3-yl)-ureido]-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



73


¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.36 (m, 3H), 1.42-1.45 (m, 3H), 3.83-3.88 (m, 11H), 3.96-4.00 (m, 2H), 4.045-4.49 (m, 2H), 7.19 (m, 1H), 7.46-7.51 (m, 3H), 8.05 (s, 1H), 8.29-8.31 (m, 2H), 9.72 (s, 1H)

5-Ethoxy-2-{4-[3-(1H-indazol-4-yl)-ureido]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.75-3.76 (m, 4H), 3.82-3.83 (m, 4H), 3.89-3.92 (m, 2H), 4.36-4.40 (m, 2H), 7.14-7.16 (m, 1H), 7.25-7.28 (m, 1H), 7.60-7.61 (m, 2H), 7.66-7.67 (m, 1H), 8.13 (s, 1H), 8.19-8.21 (m, 2H), 8.99 (s, 1H), 9.12 (m, 1H), 13.1 (s, 1H)


5-Ethoxy-2-{4-[3-(4-methyl-1H-benzotriazol-5-yl)-ureido]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.28 (m, 3H), 1.32-1.35 (m, 3H), 2.38 (s, 3H), 3.75-3.76 (m, 4H), 3.82-3.83 (m, 4H), 3.89-3.92 (m, 2H), 4.36-4.40 (m, 2H), 7.24-7.26 (m, 1H), 7.60-7.62 (m, 2H), 7.67-7.69 (m, 1H), 8.17-8.18 (m, 2H), 8.57 (s, 1H), 9.44 (s, 1H)


74

5-Ethoxy-6-morpholin-4-yl-2-{4-[3-(2-oxo-2,3-dihydro-benzooxazol-5-yl)-ureido]-phenyl}-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.75-3.76 (m, 4H), 3.82-3.83 (m, 4H), 3.89-3.92 (m, 2H), 4.36-4.40 (m, 2H), 6.90-6.93 (m, 1H), 7.17-7.19 (m, 1H), 7.49 (s, 1H), 7.54 (m, 2H), 8.15-8.17 (m, 2H), 8.82 (s, 1H), 8.95 (s, 1H)

2-(4-Amino-3-fluoro-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ 1.30-1.33 (m, 3H), 1.39-1.42 (m, 3H), 3.79-3.80 (m, 4H), 3.83-3.84 (m, 4H), 3.92-3.94 (m, 4H), 4.42-4.44 (m, 2H), 6.75 (m, 1H), 7.93-7.94 (m, 2H)


5-Ethoxy-2-(3-fluoro-4-phenoxy-carbonylamino-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



75


¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.36 (m, 3H), 1.42-1.45 (m, 3H), 3.82-3.84 (m, 4H), 3.88-3.90 (m, 4H), 3.96-4.00 (m, 2H), 4.44-4.48 (m, 2H), 7.20-7.22 (m, 4H), 7.26-7.27 (m, 1H), 7.39-7.42 (m, 2H), 8.09-8.15 (m, 3H)

5-Ethoxy-2-(3-fluoro-4-{3-[4-(3-oxo-morpholin-4-yl)-phenyl]-ureido}-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.70 (m, 2H), 3.75-3.76 (m, 4H), 3.82-3.83 (m, 4H), 3.89-3.92 (m, 2H), 3.93-3.96 (m, 2H), 4.18 (s, 2H), 4.36-4.40 (m, 2H), 7.30-7.32 (m, 2H), 7.48-7.49 (m, 2H), 7.98-8.04 (m, 2H), 8.29-8.32 (m, 1H), 8.80 (s, 1H), 9.22 (s, 1H)

2-(4-Amino-2-fluoro-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, CDCl₃-d₁): δ 1.32-1.35 (m, 3H), 1.39-1.42 (m, 3H), 3.79-3.80 (m, 4H), 3.85-3.87 (m, 4H), 3.94-3.99 (m, 4H), 4.41-4.46 (m, 2H), 6.37-6.39 (m, 1H), 6.45-6.47 (m, 1H), 7.89-7.92 (m, 1H)

5-Ethoxy-2-(2-fluoro-4-phenoxy-carbonylamino-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester

**76**


¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.36 (m, 3H), 1.42-1.45 (m, 3H), 3.82-3.84 (m, 4H), 3.88-3.90 (m, 4H), 3.96-4.00 (m, 2H), 4.44-4.48 (m, 2H), 7.13-7.14 (m, 2H), 7.18-7.20 (m, 2H), 7.39-7.48 (m, 3H), 8.04-8.07 (m, 1H)

5-Ethoxy-2-(2-fluoro-4-{3-[4-(3-oxo-morpholin-4-yl)-phenyl]-ureido}-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.28 (m, 3H), 1.30-1.33 (m, 3H), 3.69-3.80 (m, 10H), 3.89-3.97 (m, 4H), 4.18 (s, 2H), 4.34-4.39 (m, 2H), 7.19-7.21 (m, 1H), 7.29-7.31 (m, 2H), 7.47-7.49 (m, 2H), 7.55-7.58 (m, 1H), 7.91 (m, 1H), 8.88 (s, 1H), 9.13 (s, 1H)

2-{4-[3-(4-Carbamoyl-phenyl)-ureido]-3-fluoro-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.75-3.76 (m, 4H), 3.81-3.82 (m, 4H), 3.89-3.93 (m, 2H), 4.36-4.40 (m, 2H), 7.21 (m, 1H), 7.51-7.53 (m, 2H), 7.85-7.85 (m, 3H), 7.99-8.05 (m, 3H), 8.29-8.32 (m, 1H), 8.87 (s, 1H), 9.39 (s, 1H)


2-{4-[3-(4-Carbamoyl-phenyl)-ureido]-2-fluoro-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



77


¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.28 (m, 3H), 1.30-1.33 (m, 3H), 3.73-3.74 (m, 10H), 3.79-3.80 (m, 4H), 3.90-3.94 (m, 2H), 4.34-4.38 (m, 2H), 7.21-7.22 (m, 2H), 7.59 (m, 3H), 7.82-7.84 (m, 3H), 7.93-7.96 (m, 1H), 9.05 (s, 1H), 9.20 (s, 1H)

2-[4-(3-Benzo[1,3]dioxol-5-yl-ureido)-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.75 (m, 4H), 3.82 (m, 4H), 3.90-3.91 (m, 4H), 4.37-4.38 (m, 2H), 5.98 (s, 2H), 6.77 (m, 1H), 6.83 (m, 1H), 7.21 (s, 1H), 7.52-7.53 (m, 2H), 8.15-8.17 (m, 2H), 8.60 (s, 1H), 8.87 (s, 1H)

5-Ethoxy-2-{4-[3-(4-ethoxycarbonyl-phenyl)-ureido]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.28 (m, 3H), 1.29-1.33 (m, 6H), 3.75 (m, 4H), 3.82 (m, 4H), 3.90-3.91 (m, 4H), 4.37-4.38 (m, 2H), 5.98 (s, 2H), 7.55-7.60 (m, 4H), 7.88-7.90 (m, 2H), 8.17-8.19 (m, 2H), 9.06 (s, 1H), 9.15 (s, 1H)

2-(4-{3-[4-(2 Dimethylamino-ethoxy)-phenyl]-ureido}-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.28 (m, 3H), 1.29-1.33 (m, 3H), 2.20 (s, 6H), 2.59 (m, 2H), 3.75 (m, 4H), 3.82 (m, 4H), 3.90-3.91 (m, 4H), 3.99-4.03 (m, 2H), 4.37-

78


4.38 (m, 2H), 6.86 (m, 2H), 7.35-7.37 (m, 0.86 (m, 2H), 7.35-7.37 (m, 2H), 7.53-7.55 (m, 2H), 8.14-8.16 (m, 2H), 8.71 (s, 1H), 9.04 (s, 1H)

4-(3-{4-[5-Ethoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-ureido)-benzamide




¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.28 (m, 3H), 3.28 (m, 2H), 3.56 (m, 2H), 3.67 (m, 4H), 3.75 (m, 4H), 3.82 (m, 4H), 3.88-3.92 (m, 2H), 7.19 (m, 1H), 7.51-7.56 (m, 4H), 7.81-7.83 (m, 3H), 8.17-8.19 (m, 2H), 8.98 (s, 1H), 9.02 (s, 1H)

5-Ethoxy-2-{4-[3-(4-methanesulfonyl-phenyl)-ureido]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.28 (m, 3H), 1.32-1.35 (m, 3H), 3.16 (s, 3H), 3.75-3.76 (m, 4H), 3.82-3.83 (m, 4H), 3.90-3.93 (m, 2H), 4.35-4.39 (m, 2H), 7.56-7.58 (m, 2H), 7.69-7.71 (m, 2H), 7.82 (m, 2H), 8.18-8.19 (m, 2H), 9.10 (m, 1H), 9.25 (s, 1H)


5-Ethoxy-2-[4-(3-isopropyl-ureido)-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



79


¹H NMR (500 MHz, CDCl₃-d₁): δ 1.14-1.15 (m, 6H), 1.32-1.35 (m, 3H), 1.41-1.43 (m, 3H), 3.81-3.82 (m, 4H), 3.87 (m, 4H), 3.94-3.99 (m, 2H), 4.44-4.47 (m, 2H), 4.83-4.85 (m, 1H), 6.77 (s, 1H), 7.34-7.36 (m, 2H), 8.22-8.24 (m, 2H)

2-[4-(3-tert-Butyl-ureido)-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, CDCl₃-d₁): δ 1.32-1.36 (m, 12H), 1.41-1.44 (m, 3H), 3.81-3.82 (m, 4H), 3.86-3.87 (m, 4H), 3.94-3.99 (m, 2H), 4.44-4.47 (m, 2H), 4.77 (m, 1H), 6.46 (m, 1H), 7.31-7.33 (m, 2H), 8.22-8.24 (m, 2H)

5-Ethoxy-2-{4-[3-(4-methanesulfonylamino-phenyl)-ureido]-phenyl}-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.26-1.27 (m, 3H), 1.32-1.34 (m, 3H), 2.92 (s, 3H), 3.75-3.76 (m, 4H), 3.81-3.82 (m, 4H), 3.88-3.93 (m, 2H), 4.35-4.39 (m, 2H), 7.14-7.16 (m, 2H), 7.42-7.43 (m, 2H), 7.53-7.55 (m, 2H), 8.15-8.17 (m, 2H), 8.71 (s, 1H), 8.90 (s, 1H), 9.46 (s, 1H)

2-{4-[3-(4-Acetylamino-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.27 (m, 3H), 1.32-1.34 (m, 3H), 2.01 (s, 3H), 3.75-3.76 (m, 4H), 3.81-3.82

80


(m, 4H), 3.88-3.93 (m, 2H), 4.35-4.39 (m, 2H), 7.36-7.78 (m, 2H), 7.48-7.49 (m, 2H), 7.53-7.55 (m, 2H), 8.15-8.16 (m, 2H), 8.66 (s, 1H), 8.91 (s, 1H), 9.83 (m, 1H)

5-Ethoxy-6-morpholin-4-yl-2'-{3-[4-(3-oxo-morpholin-4-yl)-phenyl]-ureido}-[2,5']bipyrimidinyl-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, DMSO-d₆): δ 1.19-1.22 (m, 3H), 1.26-1.29 (m, 3H), 3.69-3.73 (m, 6H), 3.86-3.96 (m, 8H), 4.17 (s, 2H), 4.34-4.38 (m, 2H), 7.33-7.34 (m, 2H), 7.66-7.68 (m, 2H), 9.34 (s, 2H), 10.50 (s, 1H), 11.58 (s, 1H)

2'-[3-(4-Carbamoyl-phenyl)-ureido]-5-ethoxy-6-morpholin-4-yl-[2,5']bipyrimidinyl-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.29 (m, 3H), 1.32-1.35 (m, 3H), 3.76-3.77 (m, 4H), 3.88-3.95 (m, 6H), 4.36-4.41 (m, 2H), 7.26 (s, 1H), 7.73-7.75 (m, 2H), 7.86-7.90 (m, 3H), 9.38 (s, 2H), 10.62 (s, 1H), 11.81 (s, 1H)


2'-[3-(4-Acetylamino-phenyl)-ureido]-5-ethoxy-6-morpholin-4-yl-[2,5']bipyrimidinyl-4-carboxylic acid ethyl ester



81


¹H NMR (500 MHz, DMSO-d₆): δ 1.24-1.29 (m, 3H), 1.32-1.35 (m, 3H), 2.05 (s, 3H), 3.76-3.77 (m, 4H), 3.88-3.95 (m, 6H), 4.36-4.41 (m, 2H), 7.55 (m, 4H), 9.35 (s, 2H), 9.89 (s, 1H), 10.46 (s, 1H), 11.45 (s, 1H)

5-Ethoxy-2-(4-{3-[4-(4-methanesulfonyl-piperazin-1-yl)-phenyl]-ureido}-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.27 (m, 3H), 1.32-1.34 (m, 3H), 2.92 (s, 3H), 3.15-3.16 (m, 4H), 3.23-3.25 (m, 4H), 3.75-3.76 (m, 4H), 3.81-3.82 (m, 4H), 3.88-3.93 (m, 2H), 4.35-4.39 (m, 2H), 6.92-6.94 (m, 2H), 7.33-7.35 (m, 2H), 7.53-7.54 (m, 2H), 8.14-8.16 (m, 2H), 8.53 (s, 1H), 8.86 (s, 1H)

5-Ethoxy-6-morpholin-4-yl-2-{4-[3-(2-oxo-2,3-dihydro-benzooxazol-6-yl)-ureido]-phenyl}-pyrimidine-4-carboxylic acid ethyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.25-1.27 (m, 3H), 1.32-1.34 (m, 3H), 3.75 (m, 4H), 3.83 (m, 4H), 3.90-3.91 (m, 2H), 4.35-4.39 (m, 2H), 6.98-6.99 (m, 1H), 7.05-7.07 (m, 1H), 7.54-7.56 (m, 3H), 8.15-8.17 (m, 2H), 8.93 (s, 1H), 9.09 (s, 1H)

2-{4-[3-(6-Acetylamino-pyridin-3-yl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester




82

¹H NMR (500 MHz, DMSO-d₆): δ 1.20-1.23 (m, 3H), 1.27-1.30 (m, 3H), 2.02 (s, 3H), 3.71-3.72 (m, 4H), 3.78 (m, 4H), 3.84-3.87 (m, 2H), 4.32-4.34 (m, 2H), 7.50-7.52 (m, 2H), 7.77-7.79 (m, 1H), 7.96 (m, 1H), 8.11-8.13 (m, 2H), 8.38 (s, 1H), 8.75 (s, 1H), 9.01 (s, 1H)

Example 2

Preparation of Compounds of Formula (I) in Scheme 5


[2-(4-Amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidin-4-yl]-methanol



NaBH₄ (192 mg, 5 eq.) was added to a stirred solution of 2-(4-amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (378 mg, 1 eq.) in EtOH (12 ml) and the mixture reacted to reflux overnight. The mixture was quenched with H₂O, the solvent removed in vacuo, and the residue extracted with EA and washed with brine. The crude was purified by chromatography to give a product (267 mg, 80%).

¹H NMR (500 MHz, CDCl₃-d): δ 1.33-1.36 (m, 3H), 3.80-3.90 (m, 10H), 4.30 (s, 1H), 4.71 (s, 2H), 6.71-6.73 (m, 2H), 8.17-8.19 (m, 2H)

(2-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidin-4-yl)-methanol




NaBH₄ (0.64 g, 4 eq.) was added to a stirred solution of 2-chloro-5-ethoxy-6-morpholine-4-yl-pyrimidine-4-carboxylic acid ethyl ester (1.34 g, 1 eq.) in EtOH (20 ml) and the mixture reacted to reflux for 2 h. The mixture was quenched with H₂O, the solvent was removed in vacuo, and the residue was extracted with EA, washed with brine, and concentrated to give a product (1.13 g, 97.3%).

83

¹H NMR (500 MHz, CDCl₃-d₁): δ 1.34-1.37 (m, 3H), 3.37 (s, 1H), 3.77-3.84 (m, 10H), 4.63-4.64 (m, 2H)


(2-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carbaldehyde



To a suspension of Dess-Martin periodinane in CH₂Cl₂ (15%, 11.2 ml, 1.1 eq) was added to a stirred solution of (2-chloro-5-ethoxy-6-morpholin-4-yl-pyrimidin-4-yl)-methanol (1.13 g, 1 eq.) in CH₂Cl₂ (20 ml) and the mixture was reacted at r.t. for 2 h. The mixture was quenched with NaHCO₃(sat), extracted with EA, and washed with brine. The crude was purified by chromatography to give a product (1.08 g, 80.8%).

¹H NMR (500 MHz, CDCl₃-d₁) δ 1.38-1.43 (m, 3H), 3.77-4.01 (m, 10H), 10.06 (s, 1H)

4-[2-Chloro-5-ethoxy-6-(4-methanesulfonyl-piperazin-1-ylmethyl)-pyrimidin-4-yl]-morpholine




A mixture of 2-chloro-5-ethoxy-6-morpholine-4-yl-pyrimidine-4-carboxylic acid ethyl ester (271.7 mg, 1 eq.), 1-methanesulfonyl-1-piperazine (197 mg, 1.2 eq.) and trimethylorthoformate (0.33 ml, 3 eq.) was stirred in 1,2-dichloroethane (6 ml) for 6 h at room temperature. To this was added sodium triacetoxyborohydride (530 mg, 2.5 eq.) and the reaction mixture was stirred overnight at room temperature. The mixture was then quenched with brine, extracted with CH₂Cl₂, and dried, and the solvent was removed in vacuo. The residue was triturated with EA/Hex to yield 4-[2-chloro-5-ethoxy-6-(4-methanesulfonyl-piperazin-1-ylmethyl)-pyrimidin-4-yl]-morpholine as a white solid (336 mg, 80%).

¹H NMR (500 MHz, CDCl₃-d₁): δ 1.35-1.38 (m, 3H), 2.69-2.70 (m, 4H), 2.77 (s, 3H), 3.26 (m, 4H), 3.59 (s, 2H), 3.79-3.86 (m, 10H)


84

3-[5-Ethoxy-4-(4-methanesulfonyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenol




¹H NMR (500 MHz, DMSO-d₆): δ 1.31-1.34 (m, 3H), 2.66 (m, 4H), 2.86 (s, 3H), 3.10 (m, 4H), 3.61 (s, 2H), 3.74 (m, 8H), 3.90-3.93 (m, 2H), 6.82-6.84 (m, 1H), 7.23-7.26 (m, 1H), 7.71 (m, 2H), 9.50 (s, 1H)

4-[2-Chloro-5-ethoxy-6-(4-methyl-piperazin-1-ylmethyl)-pyrimidin-4-yl]-morpholine



¹H NMR (500 MHz, CDCl₃-d₁): δ 1.34-1.37 (m, 3H), 2.75 (s, 3H), 2.98 (m, 4H), 3.18 (m, 4H), 3.65 (s, 2H), 3.79-3.82 (m, 10H)

4-[2-Chloro-5-ethoxy-6-(4-morpholin-1-ylmethyl)-pyrimidin-4-yl]-morpholine




A mixture of 2-chloro-5-ethoxy-6-morpholine-4-yl-pyrimidine-4-carboxylic acid ethyl ester (878 mg, 1 eq.), morpholine (0.34 ml, 1.2 eq.) and trimethylorthoformate (1.06 ml, 3 eq.) was stirred in 1,2-dichloroethane (20 ml) for 6 hrs at room temperature. To this was added sodium triacetoxyborohydride (1.71 g, 2.5 eq.) and the reaction mixture was

85

stirred overnight at room temperature. The mixture was then quenched with brine, exacted with CH_2Cl_2 , and dried, and the solvent was removed in vacuo. The crude was purified by chromatography to give a product (1.0 g, 91%).


^1H NMR (500 MHz, $\text{CDCl}_3\text{-d}_1$): δ 1.33-1.36 (m, 3H), 2.56 (m, 4H), 3.50 (s, 2H), 3.68-3.69 (m, 4H), 3.77 (s, 8H), 3.86-3.90 (m, 2H)

4-[5-Ethoxy-4-(4-methanesulfonyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-1H-indazole




^1H NMR (500 MHz, $\text{CDCl}_3\text{-d}_1$): δ 1.39-1.42 (m, 3H), 2.79 (s, 3H), 2.83-2.84 (m, 4H), 3.30 (m, 4H), 3.81 (s, 2H), 3.86-3.88 (m, 8H), 3.95-3.99 (m, 2H), 7.46-7.49 (m, 1H), 7.57-7.58 (m, 1H), 8.20-8.21 (m, 1H), 9.01 (s, 1H)

4-[5-Ethoxy-4-(4-methanesulfonyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenylamine




^1H NMR (500 MHz, $\text{CDCl}_3\text{-d}_1$): δ 1.34-1.39 (m, 3H), 2.78 (s, 3H), 2.79-2.83 (m, 4H), 3.28 (m, 4H), 3.71 (s, 2H), 3.78-3.83 (m, 8H), 3.89-3.92 (m, 2H), 6.70-6.72 (m, 2H), 8.13-8.15 (m, 2H)

4-[5-Ethoxy-4-(4-methyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-1H-indazole

**86**


^1H NMR (500 MHz, $\text{CDCl}_3\text{-d}_1$): δ 1.38-1.41 (m, 3H), 2.33 (s, 3H), 2.57 (m, 4H), 2.79 (m, 4H), 3.77 (s, 2H), 3.85-3.88 (m, 8H), 3.97-4.01 (m, 2H), 7.44-7.47 (m, 1H), 7.55-7.56 (m, 1H), 8.20-8.21 (m, 1H), 9.02 (s, 1H)

4-[5-Ethoxy-4-(4-methyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenylamine




^1H NMR (500 MHz, $\text{CDCl}_3\text{-d}_1$): δ 1.34-1.36 (m, 3H), 2.31 (s, 3H), 2.52 (m, 4H), 2.73 (m, 4H), 3.66 (s, 2H), 3.77-3.83 (m, 8H), 3.89-3.93 (m, 2H), 6.69-6.71 (m, 2H), 8.15-8.17 (m, 2H)

4-(5-Ethoxy-4-morpholin-4-yl-6-morpholin-4-ylmethyl-pyrimidin-2-yl)-1H-indazole



^1H NMR (500 MHz, $\text{CDCl}_3\text{-d}_1$): δ 1.38-1.41 (m, 3H), 2.72 (m, 4H), 3.75-3.76 (m, 6H), 3.85-3.88 (m, 8H), 3.99-4.03 (m, 2H), 7.46-7.47 (m, 1H), 7.54-7.56 (m, 1H), 8.21-8.22 (m, 1H), 9.03 (s, 1H)


4-(5-Ethoxy-4-morpholin-4-yl-6-morpholin-4-ylmethyl-pyrimidin-2-yl)-phenylamine



87


¹H NMR (500 MHz, CDCl₃-d₁): δ 1.34-1.39 (m, 3H), 2.68 (m, 4H), 3.64 (s, 2H), 3.72-3.73 (m, 4H), 3.78-3.83 (m, 8H), 3.91-3.95 (m, 2H), 6.69-6.70 (m, 2H), 8.15-8.16 (m, 2H)

1-[4-[5-Ethoxy-4-(4-methanesulfonyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl]-3-phenyl-urea




¹H NMR (500 MHz, CDCl₃-d₁): δ 1.35-1.38 (m, 3H), 2.73 (s, 3H), 2.77 (m, 4H), 3.24 (m, 4H), 3.71 (s, 2H), 3.81-3.82 (m, 8H), 3.87-3.92 (m, 2H), 7.05 (m, 1H), 7.32-7.45 (m, 6H), 8.21-8.22 (m, 2H)

1-[4-[5-Ethoxy-4-(4-methyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl]-3-phenyl-urea




¹H NMR (500 MHz, DMSO-d₆): δ 1.30-1.33 (m, 3H), 2.16 (s, 3H), 2.36-2.63 (m, 8H), 3.53 (s, 2H), 3.74 (m, 8H), 3.91-3.93 (m, 2H), 6.98-6.99 (m, 1H), 7.27-7.30 (m, 2H), 7.46-7.47 (m, 2H), 7.53-7.53 (m, 2H), 8.18-8.20 (m, 2H), 8.83 (s, 1H), 9.00 (s, 1H)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-morpholin-4-ylmethyl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea

**88**


¹H NMR (500 MHz, CDCl₃-d₁): δ 1.35-1.38 (m, 3H), 2.65 (s, 4H), 3.66 (s, 2H), 3.69-3.71 (m, 4H), 3.80-3.82 (m, 8H), 3.90-3.95 (m, 2H), 7.02-7.05 (m, 1H), 7.23-7.37 (m, 6H), 8.21-8.23 (m, 2H)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-morpholin-4-ylmethyl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea hydrochloride




¹H NMR (500 MHz, DMSO-d₆): δ 1.33-1.36 (m, 3H), 3.48-3.96 (m, 18H), 4.58 (s, 2H), 6.97-6.99 (m, 1H), 7.27-7.30 (m, 2H), 7.46-7.48 (m, 2H), 7.58-7.60 (m, 2H), 8.29-8.31 (m, 2H), 9.40 (s, 1H), 9.64 (s, 1H)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-piperidin-1-ylmethyl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea




¹H NMR (500 MHz, CDCl₃-d₁): δ 1.33-1.40 (m, 5H), 1.56-1.58 (m, 4H), 2.60 (m, 4H), 3.64 (s, 2H), 3.77-3.80 (m, 8H), 3.86-3.90 (m, 2H), 6.93-6.96 (m, 1H), 7.16-7.19 (m, 2H), 7.26-7.28 (m, 2H), 7.31-7.33 (m, 2H), 7.92 (s, 1H), 8.03 (s, 1H), 8.14-8.16 (m, 2H)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-piperidin-1-ylmethyl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea hydrochloride




89

¹H NMR (500 MHz, DMSO-d₆): δ 1.33-1.36 (m, 3H), 1.86 (m, 4H), 2.08 (m, 2H), 3.14 (m, 2H), 3.58 (m, 2H), 3.76-3.81 (m, 8H), 3.88-3.91 (m, 2H), 4.48 (s, 2H), 6.95-6.98 (m, 1H), 7.26-7.29 (m, 2H), 7.46-7.47 (m, 2H), 7.58-7.60 (m, 2H), 8.29-8.31 (m, 2H), 9.51 (s, 1H), 9.77 (s, 1H)

Example 3**Preparation of Compounds of Formula (I) in Scheme 1****5-Methoxy-pyrimidine-2,4-diol**


A mixture of methyl methoxylacetate (15 ml, 1.0 eq) and ethyl formate (12.18 ml, 1.0 eq) was added dropwise to slurry of sodium methoxide (8.17 g, 1.0 eq) in toluene (100 ml) at ice both. After dropping at room temperature and stirring overnight, the resulting solution was dried in vacuo. Then a mixture of residue, urea (9.09 g, 1.0 eq) and NaOMe (4.10 g, 0.5 eq) in EtOH (100 ml) was refluxed at 110° C. for 4 hrs. After the solvent was dried in vacuo, water and conc. HCl solution (5<pH<4) were added. After formation of white precipitate, the mixture was filtered and dried solid in vacuo. A product was obtained as a white solid (5.14 g, 23.88%)

¹H NMR (500 MHz, DMSO-d₆): δ 7.01 (s, 1H), 3.54 (s, 3H) 2,4-Dichloro-5-methoxy-pyrimidine



In ice bath, dimethyl-phenyl-amine (6.58 ml, 51.65 mmol, 0.5 eq) was added dropwise to a slurry of 5-methoxy-pyrimidine-2,4-diol (14.68 g, 1.0 eq) in POCl₃ (47.42 ml, 5.0 eq) and refluxed overnight. Excess POCl₃ was evaporated in vacuo and the residue was poured into ice-water. After formation of pink precipitates, the mixture was filtered and dried solid in vacuo. A product was obtained as a white solid (3.89 g, 20.83%)


¹H NMR (500 MHz, CDCl₃-d₁): δ 8.18 (s, 1H), 4.01 (s, 3H)

4-(2-Chloro-5-methoxy-pyrimidin-4-yl)-morpholine

2,4-Dichloro-5-methoxy-pyrimidin (3.2 g, 1.0 eq) was stirred in toluene (20 ml) and a solution of morpholine (2.34 ml, 1.5 eq) in toluene (20 ml) was added dropwise at -10-0° C. After stirring the resulting solution overnight at r.t., NH₄Cl (aq) was added and the solution was extracted with EA. The combined organic layers were washed with brine, dried and evaporated in vacuo. A product was obtained as a white solid (3.80 g, 93.08%)


90

¹H NMR (500 MHz, CDCl₃-d₁): δ 7.71 (s, 1H), 3.84 (s, 3H), 3.82 (t, 4H), 3.76 (t, 4H)


4-[5-Methoxy-2-(3-nitro-phenyl)-pyrimidin-4-yl]-morpholine

A mixture of 4-(2-chloro-5-methoxy-pyrimidin-4-yl)-morpholine (0.2 g, 1.0 eq), 3-nitrophenyl boronic acid (0.17 g, 1.2 eq), (PPh₃)₄Pd (0.10 g, 0.1 eq) and 2 M Na₂CO₃ (1.3 ml, 3.0 eq) was heated at 80° C. and refluxed overnight. The solvent was removed in vacuo, and the residue was extracted with EA and water. The combined organic layers were washed with brine, dried and evaporated in vacuo. The crude was purified by chromatography to give a yellow solid (0.11 g, 39.97%)

¹H NMR (500 MHz, CDCl₃-d₁): δ 9.13 (s, 1H), 8.65 (d, 1H), 8.24 (d, 1H), 8.03 (s, 1H), 7.60 (t, 1H), 3.93 (s, 3H), 3.88 (d, 4H), 3.85 (d, 4H)

4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine


¹H NMR (500 MHz, CDCl₃-d₁): δ 8.12 (d, 2H), 7.98 (s, 1H), 6.70 (d, 2H), 3.86 (s, 3H), 3.82 (s, 8H)

4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenol

¹H NMR (500 MHz, CDCl₃-d₁): δ 8.20 (d, 2H), 7.96 (s, 1H), 6.91 (d, 2H), 3.89 (s, 3H), 3.84 (m, 4H), 3.83 (m, 4H)

91


Phenyl-carbamic acid 4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester



Phenyl isocyanate (0.03 ml, 2.0 eq) was added to a solution of 4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenol (0.05 g, 1.0 eq) in dioxane (6 ml) and heated at 80° C. for 20 hrs. The solvent was removed in vacuo, and the residue was extracted with EA and water. The combined organic layers were washed with brine, dried and evaporated in vacuo. The crude was purified by chromatography to give a white solid (0.07 g, 24.03%).


¹H NMR (500 MHz, CDCl₃-d₁): δ 8.34 (d, 2H), 8.02 (s, 1H), 7.45 (d, 2H), 7.33 (t, 2H), 7.27 (s, 1H), 7.12 (m, 1H), 6.90 (s, 1H), 3.90 (s, 3H), 3.86 (m, 4H), 3.84 (m, 4H), 3.70 (s, 1H)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea



¹H NMR (500 MHz, CDCl₃-d₁): δ 8.19 (d, 2H), 7.95 (s, 1H), 7.37 (d, 2H), 7.33 (d, 2H), 7.27 (m, 2H), 7.23 (s, 1H), 7.19 (s, 1H), 7.05 (m, 1H), 3.86 (s, 3H), 3.83 (m, 4H), 3.83 (m, 4H)


1-Ethyl-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea



92


¹H NMR (500 MHz, CDCl₃-d₁): δ 8.24 (d, 2H), 8.00 (s, 1H), 7.34 (d, 2H), 6.35 (s, 1H), 4.72 (s, 1H), 3.89 (s, 3H), 3.83 (s, 4H), 3.30 (m, 2H), 1.15 (m, 3H)

5 1-(4-Fluoro-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea




¹H NMR (500 MHz, CDCl₃-d₁): δ 8.25 (d, 2H), 7.99 (s, 1H), 7.39 (d, 2H), 7.32 (m, 2H), 7.03 (m, 2H), 6.69 (s, 1H), 6.67 (s, 1H), 3.89 (s, 3H), 3.84 (s, 8H)

25 1-(3-Fluoro-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea



¹H NMR (500 MHz, CDCl₃-d₁): δ 8.27 (d, 2H), 8.00 (s, 1H), 7.39 (d, 2H), 7.30 (m, 1H), 7.24 (m, 1H), 7.00 (m, 1H), 6.80 (s, 1H), 6.78 (m, 1H), 6.60 (s, 1H), 3.89 (s, 3H), 3.84 (s, 8H)


50 1-(3,4-Difluoro-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea



93


¹H NMR (500 MHz, CDCl₃-d₁): δ8.27 (d, 2H), 8.00 (s, 1H), 7.42 (m, 1H), 7.39 (d, 2H), 7.00 (m, 1H), 6.97 (m, 1H), 6.75 (s, 1H), 6.68 (s, 1H), 3.89 (s, 3H), 3.84 (s, 8H)

1-(4-Fluoro-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thiourea




¹H NMR (500 MHz, CDCl₃-d₁): δ8.35 (d, 2H), 8.01 (s, 1H), 7.82 (m, 1H), 7.70 (m, 1H), 7.40 (d, 2H), 7.37 (d, 2H), 7.0 (d, 2H), 3.91 (s, 3H), 3.84 (s, 8H)

1-(3-Fluoro-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thiourea




¹H NMR (500 MHz, CDCl₃-d₁): δ8.36 (d, 2H), 8.02 (s, 1H), 7.41 (d, 2H), 7.33 (m, 2H), 7.26 (s, 1H), 7.11 (m, 1H), 6.94 (m, 1H), 3.91 (s, 3H), 3.84 (s, 8H)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea

**94**


¹H NMR (500 MHz, DMSO-d₆): δ9.18 (s, 1H), 9.09 (s, 1H), 8.62 (d, 1H), 8.17-8.19 (t, 3H), 8.13 (s, 1H), 7.95-7.97 (d, 1H), 7.54-7.56 (d, 2H), 7.30-7.33 (m, 1H), 3.87 (s, 1H), 3.73-3.74 (m, 8H)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-methyl-urea




¹H NMR (500 MHz, DMSO-d₆): δ9.35 (s, 1H), 8.19-8.21 (d, 2H), 7.71-7.73 (d, 3H), 6.42 (s, 1H), 4.22 (s, 4H), 3.94 (s, 3H), 3.87 (s, 4H), 2.84 (s, 3H)

1-(3,4-Dimethoxy-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ9.18 (s, 1H), 8.89 (s, 1H), 8.13-8.17 (m, 3H), 7.52-7.54 (d, 2H), 7.23 (s, 1H), 6.87-6.89 (m, 2H), 3.87 (s, 3H), 3.70-3.74 (m, 14H)


1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3-trifluoromethyl-phenyl)-urea



95


¹H NMR (500 MHz, DMSO-d₆): δ9.10 (s, 1H), 9.01 (s, 1H), 8.18-8.19 (d, 2H), 8.14 (s, 1H), 8.02 (s, 1H), 7.50-7.53 (m, 4H), 7.31-7.32 (d, 1H), 3.87 (s, 3H), 3.73 (m, 8H)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ9.11 (s, 1H), 8.98 (s, 1H), 8.17-8.18 (d, 2H), 8.14 (s, 1H), 7.53-7.54 (d, 2H), 7.48-7.50 (d, 2H), 7.28-7.29 (d, 2H), 4.18 (s, 2H), 3.96 (s, 2H), 3.87 (s, 3H), 3.69-3.74 (m, 10H)

1-(3-Fluoro-4-morpholin-4-yl-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ9.11 (s, 1H), 8.98 (s, 1H), 8.16-8.17 (d, 2H), 8.13 (s, 1H), 7.51-7.53 (d, 2H), 7.47-7.44 (s, 1H), 7.08-7.09 (m, 1H), 6.95-6.99 (m, 1H), 3.87 (s, 3H), 3.73-3.74 (m, 12H), 2.92-2.94 (m, 4H)


1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-morpholin-4-yl-ethyl)-urea



96


¹H NMR (500 MHz, DMSO-d₆): δ8.91 (s, 1H), 8.10-8.12 (d, 3H), 7.45-7.46 (d, 2H), 6.25 (s, 1H), 3.86 (s, 3H), 3.72 (s, 8H), 3.58-3.60 (m, 4H), 3.19-3.22 (m, 2H), 2.36-2.39 (m, 6H)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-methyl-piperazin-1-yl)-urea




¹H NMR (500 MHz, DMSO-d₆): δ8.60 (s, 1H), 8.12-8.14 (d, 2H), 8.12 (s, 1H), 7.77 (s, 1H), 7.60-7.62 (d, 2H), 3.86 (s, 3H), 3.73 (s, 8H), 2.58-2.74 (m, 8H), 2.19 (s, 3H)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-morpholin-4-yl)-urea



¹H NMR (500 MHz, DMSO-d₆): δ8.85 (s, 1H), 8.56 (s, 1H), 8.15-8.17 (d, 2H), 8.13 (s, 1H), 7.50-7.52 (d, 2H), 7.31-7.33 (d, 2H), 6.88-6.90 (d, 2H), 3.87 (s, 3H), 3.72-3.74 (m, 12H), 3.01-3.03 (m, 4H)


1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-thiomorpholin-4-yl)-urea



¹H NMR (500 MHz, DMSO-d₆): δ8.89 (s, 1H), 8.59 (s, 1H), 8.15-8.17 (d, 2H), 8.13 (s, 1H), 7.50-7.52 (d, 2H), 7.31-7.32 (d, 2H), 6.87-6.89 (d, 2H), 3.87 (s, 3H), 3.73-3.74 (m, 8H), 3.38-3.39 (m, 4H), 2.67-2.69 (m, 4H)

97


[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester



A mixture of 4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (1.50 g, 1.0 eq.) and sodium bicarbonate solution (6 ml) in ethyl acetate (6 ml) stirring 5 min. After dropping phenyl chloroformate (0.98 ml, 1.2 eq) for 4 hrs at room temperature. The resultant mixture extracted with EA and the combined organic layers were washed with brine and dried in vacuo to give a light brown solid (1.80 g, 40.5%) as the product.


¹H NMR (500 MHz, CDCl₃-d₁): δ8.36-8.38 (d, 2H), 7.95 (s, 1H), 7.61-7.63 (d, 2H), 7.39-7.42 (m, 3H), 7.26 (s, 1H), 7.19-7.25 (m, 2H), 4.02 (s, 4H), 3.90 (s, 3H), 3.85-3.86 (m, 4H)

4-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide



¹H NMR (500 MHz, DMSO-d₆): δ9.40-9.60 (m, 2H), 8.17-8.19 (d, 2H), 8.14 (s, 1H), 7.80-7.82 (d, 3H), 7.53-7.56 (m, 4H), 7.18 (s, 1H), 3.87 (s, 3H), 3.73-3.74 (m, 8H).


4-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzenesulfonamide



98


¹H NMR (500 MHz, DMSO-d₆): δ9.30 (s, 1H), 9.15 (s, 1H), 8.18-8.20 (d, 2H), 8.14 (s, 1H), 7.72-7.74 (d, 2H), 7.62-7.64 (d, 2H), 7.54-7.56 (d, 2H), 7.21 (s, 2H), 3.87 (s, 3H), 3.73-3.74 (m, 8H)

5 2-Nitro-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenylamine



¹H NMR (500 MHz, CDCl₃-d₁): δ8.59 (s, 1H), 7.71-7.73 (d, 1H), 6.76-6.78 (d, 1H), 6.20-6.30 (m, 2H), 1.33 (s, 12H).


4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-2-nitro-phenylamine



A mixture of 4-(2-chloro-5-methoxy-pyrimidin-4-yl)-morpholine (1.0 g, 1 eq.), 2-nitro-4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenylamine (1.26 g, 1.1 eq.), dioxane (20 ml), Pd(PPh₃)₄ (0.05 g, 0.1 eq.) and 2M sodium carbonate (6.53 ml, 3.0 eq.) was heated to reflux overnight. The solvent was removed in vacuo, the residue extracted with EA, and the organic layer washed with brine and dried. The crude was purified by chromatography to give a brown solid (2.08 g, 47.83%).

¹H NMR (500 MHz, CDCl₃-d₁): δ9.06 (s, 1H), 8.32-8.34 (d, 1H), 7.98 (s, 1H), 6.84-6.86 (d, 1H), 6.20 (s, 2H), 3.86 (s, 3H), 3.84 (s, 8H)

1-(2-Amino-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thiourea




99

A mixture of di-imidazol-1-yl-methanethione (0.5 g, 1.50 eq.), imidazole (0.03 g, 0.3 eq.) and CH₃CN (7 ml) was added dropwise to 4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (0.5 g, 1.0 eq.) in CH₂Cl₂ (5 ml) at 0° C. When S.M. is disappear in TLC, dropping benzene-1,2-diamine (0.37 g, 3.0 eq) to the solution for 3 hrs at 50° C. The solvent was removed in vacuo. The resulting solution was crystallized in a EA solution, and filtered to given a light yellow solid, which was then dried in vacuo to give of a product (0.53 g, 70.18%).

¹H NMR (500 MHz, DMSO-d₆): δ 9.80 (s, 1H), 9.20 (s, 1H), 8.18-8.19 (d, 2H), 8.15 (s, 1H), 7.63-7.64 (d, 2H), 7.09-7.10 (m, 1H), 6.95-6.96 (m, 1H), 6.74-6.75 (m, 1H), 6.55-6.58 (m, 1H), 4.92 (s, 2H), 3.88 (s, 3H), 3.72-3.75 (m, 8H).


4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-benzene-1,2-diamine



A mixture of 4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-2-nitro-phenylamine (0.69 g, 1.0 eq.) and Pd/C (Cat.) in MeOH/EA (10 ml/10 ml) was prepared. The mixture was sharked and H₂ gas was injected to the mixture for 4-6 hrs at room temperature. The mixture was filtered with Celite 454 and dried in vacuo to give a black sold (0.65 g, 99.9%) as a product.

¹H NMR (500 MHz, CDCl₃-d₁): δ 8.03 (s, 1H), 7.49 (s, 1H), 7.36-7.38 (m, 1H), 6.50-6.51 (d, 1H), 4.77 (s, 2H), 4.49 (s, 2H), 3.82 (s, 3H), 3.70 (s, 8H).

(1H-Benzoimidazol-2-yl)-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-amine




A mixture of 1-(2-Amino-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thiourea (0.3 g, 1.0 eq.) and DCC (0.18 g, 1.3 eq.) in THF (10 ml) for 9 hrs at 80-90° C. The mixture was dried in vacuo, crystallized in a EA solution, and then filtered to give a white solid (0.18 g, 68.36%), which was then dried in vacuo to give of a product.

¹H NMR (500 MHz, DMSO-d₆): δ 11.2 (s, 1H), 9.87 (s, 1H), 8.20-8.22 (d, 2H), 8.14 (s, 1H), 7.82-7.84 (d, 2H), 7.28-7.36 (d, 2H), 7.00 (s, 2H), 3.87 (s, 3H), 3.74 (s, 8H).

100


5-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-1H-benzoimidazol-2-ylamine



A mixture of 4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-benzene-1,2-diamine (0.58 g, 1.0 eq.) and ACN/H₂O (5 ml/1 ml). In ice bath, cyanogen bromide (0.3 g, 1.5 eq) was added to mixture solution dropwisely. The solution was stirred overnight at room temperature, quenched with a NaHCO₃(sat.) solution, and then filtered to give a light brown solid (0.32 g, 51.78%), which was dried in vacuo to give a product.


¹H NMR (500 MHz, DMSO-d₆): δ 8.12 (s, 1H), 8.06 (s, 1H), 7.93-7.95 (d, 1H), 7.14-7.5 (d, 1H), 6.63 (s, 2H), 3.86 (s, 3H), 3.74 (s, 8H).

1-(4-Dimethylaminomethoxy-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ 8.86 (s, 1H), 8.60 (s, 1H), 8.15-8.17 (d, 2H), 8.13 (s, 1H), 7.50-7.52 (d, 2H), 7.34-7.36 (d, 2H), 6.87-6.89 (d, 2H), 4.00-4.02 (m, 2H), 3.87 (s, 3H), 3.73-3.74 (m, 8H), 2.63-2.66 (m, 2H), 1.90 (s, 6H)


3-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide



101


¹H NMR (500 MHz, DMSO-d₆): δ8.92 (s, 1H), 8.86 (s, 1H), 8.17-8.19 (d, 2H), 8.14 (s, 1H), 7.94 (s, 1H), 7.89 (s, 1H), 7.53-7.55 (d, 2H), 7.46-7.47 (d, 2H), 7.33-7.37 (m, 2H), 3.87 (s, 3H), 3.73-3.74 (m, 8H)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-thiazol-2-yl-urea




¹H NMR (500 MHz, DMSO-d₆): δ9.15 (s, 1H), 8.20-8.22 (d, 2H), 8.14 (s, 1H), 7.55-7.56 (d, 2H), 7.39 (s, 1H), 7.20 (s, 1H), 3.87 (s, 3H), 3.73-3.74 (m, 8H)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-methyl-piperazin-1-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ9.22 (s, 1H), 8.18-8.20 (d, 2H), 8.14 (s, 1H), 7.97 (s, 1H), 7.56-7.58 (d, 2H), 7.47 (s, 2H), 3.87 (s, 3H), 3.73-3.74 (m, 8H), 3.17 (m, 4H), 3.06 (m, 4H), 1.16-1.19 (m, 3H).


1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-thio-urea



102


¹H NMR (500 MHz, DMSO-d₆): δ10.00-10.20 (m, 2H), 8.19-8.21 (d, 2H), 8.16 (s, 1H), 7.59-7.60 (d, 2H), 7.51 (s, 2H), 7.33-7.34 (d, 2H), 4.19 (s, 2H), 3.96-3.98 (m, 2H), 3.88 (s, 3H), 3.72-3.75 (m, 8H).

4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ8.70 (s, 1H), 8.12 (s, 1H), 8.12-8.10 (d, 2H), 7.45 (d, 2H), 5.90 (s, 2H), 3.86 (s, 3H), 3.72-3.73 (m, 8H).

[1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-methyl-piperazin-1-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ8.95 (s, 1H), 8.75 (s, 1H), 8.15-8.17 (d, 2H), 8.13 (s, 1H), 7.50-7.52 (d, 2H), 7.34-7.36 (d, 2H), 6.93-6.95 (d, 2H), 3.87 (s, 3H), 3.73-3.74 (m, 8H).


1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-urea



103


¹H NMR (500 MHz, DMSO-d₆): δ8.90 (s, 1H), 8.51 (s, 1H), 8.15-8.17 (d, 2H), 8.13 (s, 1H), 7.70-7.72 (m, 1H), 7.50-7.52 (d, 2H), 6.84-6.86 (d, 2H), 3.87 (s, 3H), 3.73-3.74 (m, 8H), 3.30-3.44 (m, 8H), 2.54 (s, 3H).

1-Isopropyl-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ8.49 (s, 1H), 8.10-8.11 (m, 3H), 7.42-7.44 (d, 2H), 6.05 (s, 1H), 3.86 (s, 3H), 3.72-3.76 (m, 8H), 1.09-1.10 (m, 6H).

1-(4-Amino-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ8.89 (s, 1H), 8.40 (s, 1H), 8.13-8.15 (m, 3H), 7.49-7.51 (d, 2H), 7.08-7.09 (d, 2H), 6.50-6.51 (d, 2H), 4.77 (s, 2H), 3.86 (s, 3H), 3.73-3.74 (m, 8H).


4-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzoic acid



104


¹H NMR (500 MHz, DMSO-d₆): δ9.38 (s, 1H), 9.27 (s, 1H), 8.18-8.20 (d, 2H), 8.14 (s, 1H), 7.87-7.88 (d, 2H), 7.55-7.59 (m, 4H), 3.87 (s, 3H), 3.73-3.74 (m, 8H)

5 1-(6-Bromo-pyridin-3-yl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ9.05-9.06 (d, 2H), 8.47 (s, 1H), 8.18-8.20 (d, 2H), 8.14 (s, 1H), 7.90-7.92 (d, 1H), 7.52-7.56 (m, 3H), 3.87 (s, 3H), 3.73-3.74 (m, 8H).

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(6-morpholin-4-yl-pyridin-3-yl)-urea



¹H NMR (500 MHz, DMSO-d₆): δ8.88 (s, 1H), 8.50 (s, 1H), 8.13-8.18 (m, 4H), 7.72-7.74 (m, 2H), 7.50-7.52 (d, 2H), 6.81-6.83 (d, 1H), 3.87 (s, 3H), 3.69-3.74 (m, 12H), 3.30-3.36 (m, 4H).


1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-carbonyl)-phenyl]-urea



105


¹H NMR (500 MHz, DMSO-d₆): δ8.95-8.97 (d, 2H), 8.18-8.19 (d, 2H), 8.14 (s, 1H), 7.52-7.54 (m, 4H), 7.36-7.37 (d, 2H), 3.87 (s, 3H), 3.73-3.74 (m, 8H), 3.60 (s, 4H), 3.43-3.48 (m, 4H).

2-Fluoro-4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine




¹H NMR (500 MHz, DMSO-d₆): δ7.91-7.97 (m, 3H), 6.78-6.81 (m, 1H), 3.88 (s, 3H), 3.80-3.84 (m, 8H).

[2-Fluoro-4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ8.12 (s, 1H), 8.06-8.09 (m, 2H), 7.40-7.43 (m, 3H), 7.27-7.29 (m, 2H), 7.21-7.22 (m, 2H), 3.90 (s, 3H), 3.84 (s, 8H).


3-Fluoro-4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine



106


¹H NMR (500 MHz, DMSO-d₆): δ8.02 (s, 1H), 7.84-7.88 (m, 1H), 6.47-6.49 (m, 1H), 6.40-6.43 (m, 1H), 3.88-3.89 (m, 5H), 3.81-3.88 (m, 8H).

[3-Fluoro-4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester




¹H NMR (500 MHz, CDCl₃-d₁): δ7.99-8.04 (m, 2H), 7.39-7.46 (m, 3H), 7.15-7.27 (m, 5H), 3.90 (s, 3H), 3.82 (s, 8H).

4-{3-[2-Fluoro-4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide



¹H NMR (500 MHz, DMSO-d₆): δ9.56 (s, 1H), 8.99 (s, 1H), 8.22-8.25 (m, 1H), 8.15 (s, 1H), 7.99-8.05 (m, 2H), 7.82-7.84 (m, 3H), 7.52-7.54 (d, 2H), 7.20 (s, 1H), 3.88 (s, 3H), 3.72-3.75 (m, 8H).


1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-2-yl-urea



¹H NMR (500 MHz, DMSO-d₆): δ10.65 (s, 1H), 9.52 (s, 1H), 8.29 (s, 1H), 8.20-8.21 (d, 2H), 8.14 (s, 1H), 7.74-7.76 (m, 1H), 7.59-7.61 (d, 2H), 7.53-7.54 (m, 1H), 7.01-7.03 (m, 1H), 3.87 (s, 3H), 3.73-3.74 (m, 8H).


107

1-[2-Fluoro-4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.50 (s, 1H), 9.01 (s, 1H), 8.21-8.25 (m, 1H), 8.15 (s, 1H), 7.98-8.05 (m, 2H), 7.49-7.51 (d, 2H), 7.29-7.31 (d, 2H), 4.18 (s, 2H), 3.95-3.97 (m, 2H), 3.88 (s, 3H), 3.69-3.76 (m, 10H).

1-[3-Fluoro-4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ 9.31 (s, 1H), 9.10 (s, 1H), 8.14 (s, 1H), 7.89-7.92 (m, 1H), 7.47-7.54 (m, 3H), 7.26-7.28 (d, 2H), 7.18 (s, 1H), 4.16 (s, 2H), 3.93-3.95 (m, 2H), 3.86 (s, 3H), 3.68-3.70 (m, 10H).


4-{3-[3-Fluoro-4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide



108


¹H NMR (500 MHz, DMSO-d₆): δ 8.16 (s, 1H), 7.92 (s, 1H), 7.80-7.82 (m, 3H), 7.56-7.58 (m, 3H), 7.19-7.30 (m, 2H), 3.88 (s, 3H), 3.71-3.72 (m, 8H), 3.16 (s, 2H).

5 1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-thiourea




¹H NMR (500 MHz, DMSO-d₆): δ 10.30-10.10 (m, 2H), 8.60 (s, 1H), 8.31-8.32 (d, 1H), 8.20-8.22 (m, 2H), 8.16 (s, 1H), 7.94 (s, 1H), 7.58-7.61 (m, 2H), 7.36-7.37 (d, 1H), 3.88 (s, 3H), 3.72-3.76 (m, 8H).

N-(3-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-acetamide



¹H NMR (500 MHz, DMSO-d₆): δ 9.92 (s, 1H), 8.92 (s, 1H), 8.75-8.78 (d, 1H), 8.18-8.20 (m, 2H), 8.17 (s, 1H), 7.77 (s, 1H), 7.51-7.55 (m, 2H), 7.18-7.19 (m, 2H), 3.87 (s, 3H), 3.73-3.75 (m, 8H), 1.90 (s, 3H).


N-(3-Dimethylamino-propyl)-4-{3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide



109


¹H NMR (500 MHz, DMSO-d₆): δ9.08-9.10 (d, 2H), 8.36 (s, 1H), 8.18-8.19 (d, 2H), 8.14 (s, 1H), 7.77-7.78 (d, 2H), 7.52-7.55 (m, 4H), 3.87 (s, 3H), 3.73-3.74 (m, 8H), 2.26-2.29 (s, 2H), 2.15 (s, 6H), 1.63-1.66 (s, 2H).

5-Methoxy-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-ylamine




¹H NMR (500 MHz, CDCl₃-d₁): δ9.14 (s, 2H), 7.96 (s, 1H), 5.20 (s, 2H), 3.89 (s, 3H), 3.82 (s, 8H).

(5-Methoxy-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-yl)-carbamic acid phenyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ9.47 (s, 2H), 8.51 (s, 1H), 7.99 (s, 1H), 7.39-7.42 (m, 2H), 7.26-7.29 (m, 3H), 3.91 (s, 3H), 3.80 (s, 8H).


4-{3-[3-Fluoro-4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide



110


¹H NMR (500 MHz, DMSO-d₆): δ9.39 (s, 2H), 8.19 (s, 1H), 7.86-7.88 (m, 2H), 7.72-7.74 (m, 2H), 7.56-7.57 (m, 1H), 7.25 (s, 1H), 6.50-6.51 (d, 1H), 5.58 (s, 1H), 3.90 (s, 3H), 3.82 (s, 4H), 3.73 (s, 4H).

5 N-(4-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-methanesulfonamide




¹H NMR (500 MHz, DMSO-d₆): δ9.09 (s, 1H), 8.91 (s, 1H), 8.16-8.17 (d, 2H), 8.13 (s, 1H), 7.52-7.53 (d, 2H), 7.42-7.44 (d, 2H), 7.13-7.14 (d, 2H), 3.87 (s, 3H), 3.73-3.74 (m, 8H), 2.91 (s, 3H).

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzoxazol-6-yl)-urea



¹H NMR (500 MHz, DMSO-d₆): δ8.99 (s, 1H), 8.92 (s, 1H), 8.16-8.18 (d, 2H), 8.13 (s, 1H), 7.52-7.53 (d, 2H), 7.47 (s, 1H), 7.15-7.16 (d, 1H), 6.90-6.92 (d, 1H), 3.87 (s, 3H), 3.73-3.74 (m, 8H).


1-(5-Methoxy-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-yl)-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea



111

¹H NMR (500 MHz, DMSO-d₆): δ 11.59 (s, 1H), 10.47 (s, 1H), 9.37 (s, 2H), 8.18 (s, 1H), 7.67-7.59 (d, 2H), 7.35-7.37 (d, 2H), 4.19 (s, 2H), 3.96-3.98 (m, 2H), 3.90 (s, 3H), 3.81-3.82 (m, 2H), 3.72 (s, 8H).


2-Chloro-4-morpholin-4-yl-pyrimidin-5-ol



A mixture of 4-(2-chloro-5-methoxy-pyrimidin-4-yl)-morpholine (5.0 g, 1.0 eq.) and CH₂Cl₂ (20 ml) was added dropwise to a slurry of aluminum chloride (23.22 g, 8.0 eq.) in CH₂Cl₂ (30 ml) in ice bath. After dropping, the resulting solution was stirred overnight at room temperature. The resultant mixture was extracted with EA and the combined organic layers were washed with brine and dried in vacuo to give a light yellow solid (4.18 g, 89.23%) as a product.

¹H NMR (500 MHz, DMSO-d₆): δ 10.34 (s, 1H), 7.67 (s, 1H), 3.72-3.73 (m, 4H), 3.65-3.66 (m, 4H).

4-(2-Chloro-5-(2-Morpholin-4-ethoxy)-pyrimidin-4-yl)-morpholine




A mixture of 2-chloro-4-morpholin-4-yl-pyrimidin-5-ol (0.1 g, 1.0 eq.), 4-(2-chloro-ethyl)-morpholine hydrochloric acid (0.12 g, 1.5 eq.) and K₂CO₃ (0.19 g, 3.0 eq.) in 1,4-dioxane (5 ml) was heated at 90° C. and refluxed overnight. The resultant mixture was extracted with EA and the combined organic layers were washed with brine and dried in vacuo to give a yellow solid (0.18 g, 68.02%) as a product.

112


¹H NMR (500 MHz, CDCl₃-d₁): δ 7.73 (s, 1H), 4.07-4.09 (m, 2H), 3.86-3.88 (m, 4H), 3.76-3.78 (m, 4H), 3.69-3.71 (m, 4H), 2.75-2.77 (m, 2H), 2.50-2.53 (m, 4H).

4-(5-(2-morpholin-4-yl-ethoxy)-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine




¹H NMR (500 MHz, CDCl₃-d₁): δ 8.13-8.15 (d, 2H), 7.97 (s, 1H), 6.71-6.72 (d, 2H), 4.12-4.14 (m, 2H), 3.90-3.91 (m, 4H), 3.87-3.88 (m, 4H), 3.82-3.83 (m, 4H), 2.78-2.80 (m, 2H), 2.50-2.58 (m, 4H).

{4-[4-Morpholin-4-yl-5-(2-morpholin-4-yl-ethoxy)-pyrimidin-2-yl]-phenyl}-carbamic acid phenyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ 8.31-8.32 (d, 2H), 8.01 (s, 1H), 7.53-7.55 (d, 2H), 7.39-7.42 (m, 2H), 7.20-7.26 (m, 3H), 7.09 (s, 1H), 4.22-4.24 (m, 2H), 3.91-3.93 (m, 4H), 3.83-3.85 (m, 4H), 3.77-3.80 (m, 4H), 2.87-2.89 (m, 2H), 2.64-2.66 (m, 4H).


1-[4-(Morpholine-4-carbonyl)-phenyl]-3-{4-[4-morpholin-4-yl-5-(2-morpholin-4-yl-ethoxy)-pyrimidin-2-yl]-phenyl}-urea



113


¹H NMR (500 MHz, DMSO-d₆): δ9.19-9.21 (d, 2H), 8.17-8.18 (d, 2H), 8.14 (s, 1H), 7.53-7.55 (d, 4H), 7.35-7.37 (d, 2H), 4.17-4.19 (m, 2H), 3.79-3.81 (d, 4H), 3.73 (s, 4H), 3.59 (s, 4H), 3.56 (s, 4H), 3.49 (s, 4H), 2.70-2.71 (d, 2H), 2.36 (s, 4H).

1-{4-[4-Morpholin-4-yl-5-(2-morpholin-4-ylethoxy)-pyrimidin-2-yl]-phenyl}-3-[4-(3-oxomorpholin-4-yl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ9.20 (s, 1H), 9.19 (s, 1H), 8.14-8.18 (m, 3H), 7.53-7.54 (d, 2H), 7.48-7.50 (d, 2H), 7.27-7.29 (d, 2H), 4.18 (s, 4H), 3.95-3.97 (m, 2H), 3.79-3.80 (s, 4H), 3.73 (s, 4H), 3.68-3.70 (m, 2H), 3.55-3.56 (m, 4H), 2.69-2.71 (m, 2H), 2.45 (s, 4H).

4-(3-{4-[4-Morpholin-4-yl-5-(2-morpholin-4-ylethoxy)-pyrimidin-2-yl]-phenyl}-ureido)-benzamide



¹H NMR (500 MHz, DMSO-d₆): δ9.30-9.50 (m, 2H), 8.14-8.18 (m, 3H), 7.80-7.82 (m, 3H), 7.53-7.56 (m, 4H), 7.18 (s, 1H), 4.17-4.19 (m, 2H), 3.79-3.80 (m, 4H), 3.73-3.74 (m, 4H), 3.55-3.56 (m, 4H), 2.69-2.71 (m, 2H), 2.36 (s, 4H).

1-{4-[4-Morpholin-4-yl-5-(2-morpholin-4-ylethoxy)-pyrimidin-2-yl]-phenyl}-3-pyridin-4-yl-urea




¹H NMR (500 MHz, DMSO-d₆): δ9.68 (s, 1H), 9.45 (s, 1H), 8.34-8.35 (d, 2H), 8.17-8.19 (d, 2H), 8.15 (s, 1H), 7.54-

114


7.56 (d, 2H), 7.45-7.48 (d, 2H), 4.18 (s, 2H), 3.79-3.80 (m, 4H), 3.73-3.74 (m, 4H), 3.55-3.57 (m, 4H), 2.63-2.64 (m, 2H), 2.50-2.57 (m, 4H).

1-(4-Methanesulfonyl-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ9.72 (s, 1H), 9.45 (s, 1H), 8.18-8.20 (d, 2H), 8.14 (s, 1H), 7.81-7.82 (d, 2H), 7.71-7.73 (d, 2H), 7.56-7.58 (d, 2H), 3.87 (s, 3H), 3.73-3.74 (m, 8H), 3.15 (s, 3H).

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-sulfonyl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ9.52 (s, 1H), 9.23 (s, 1H), 8.19-8.20 (d, 2H), 8.14 (s, 1H), 7.72-7.74 (d, 2H), 7.64-7.66 (d, 2H), 7.55-7.57 (d, 2H), 3.87 (s, 3H), 3.73-3.74 (m, 8H), 3.62-3.64 (m, 4H), 2.84 (m, 4H).


Pyridin-4-yl-carbamic acid phenyl ester



115


¹H NMR (500 MHz, CDCl₃-d₁): δ8.50-8.51 (m, 2H), 7.88 (s, 1H), 7.39-7.45 (m, 4H), 7.27-7.28 (m, 1H), 7.17-7.22 (m, 2H).

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea




¹H NMR (500 MHz, DMSO-d₆): δ9.32 (s, 1H), 9.22 (s, 1H), 8.35-8.36 (d, 2H), 8.18-8.20 (d, 2H), 8.14 (s, 1H), 7.54-7.55 (d, 2H), 7.44-7.45 (d, 2H), 3.87 (s, 3H), 3.70-3.74 (m, 8H).

1-[4-(4-Methanesulfonyl-piperazin-1-yl)-phenyl]-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ8.85 (s, 1H), 8.57 (s, 1H), 8.13-8.17 (m, 3H), 7.50-7.52 (d, 2H), 7.33-7.34 (d, 2H), 6.92-6.94 (d, 2H), 3.87 (s, 3H), 3.72-3.73 (m, 8H), 3.22-3.24 (m, 4H), 3.15-3.16 (m, 4H), 2.92 (s, 3H).


N-(5-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]ureido}-pyridin-2-yl)-phenyl-urea



116

¹H NMR (500 MHz, DMSO-d₆): δ10.37 (s, 1H), 9.01 (s, 1H), 8.82 (s, 1H), 8.42 (d, 1H), 8.14-8.18 (d, 3H), 8.00-8.05 (m, 1H), 7.82-7.83 (m, 1H), 7.52-7.54 (d, 2H), 3.87 (s, 3H), 3.73-3.74 (m, 8H), 2.06 (s, 3H).


5-Ethoxy-pyrimidine-2,4-diol



A mixture of ethyl ethoxylacetate (20 g, 1.0 eq) and ethyl formate (11.21 ml, 1.0 eq) was added dropwise to a slurry of sodium ethoxide (10.3 g, 1.0 eq) in toluene (100 ml) in ice both. After dropping at room temperature and stirring overnight, the resulting solution was dried in vacuo. Then a mixture of residue, urea (9.09 g, 1.0 eq) and sodium ethoxide (10.3 g, 1.0 eq) in EtOH (100 ml) was refluxed at 110° C. for 6 hrs. After the solvent was dried in vacuo, water and conc. HCl solution (5<pH<4) were added. After formation of white precipitates, the mixture was filtered and dried solid in vacuo. A product was obtained as a white solid (7.76 g, 32%)

¹H NMR (500 MHz, DMSO-d₆): δ1.20-1.23 (m, 3H), 3.75-3.80 (m, 2H), 5.50 (s, 2H), 7.05 (s, 1H)


2,4-Dichloro-5-ethoxy-pyrimidine



In ice bath, dimethyl-phenyl-amine (5.21 ml, 1.0 eq) was added dropwise to slurry of 5-ethoxy-pyrimidine-2,4-diol (6.45 g, 1.0 eq) in POCl₃ (20 ml, 5.0 eq) and refluxed overnight. Excess POCl₃ was evaporated in vacuo and the residue was poured into ice-water. After formation of precipitates, the mixture was filtered and dried solid in vacuo. A product was obtained as a white solid (2.84 g, 35.57%)

¹H NMR (500 MHz, CDCl₃-d₁): δ1.47-1.50 (m, 3H), 4.16-4.20 (m, 2H), 8.13 (s, 1H)

4-(2-Chloro-5-ethoxy-pyrimidin-4-yl)-morpholine




2,4-dichloro-5-ethoxy-pyrimidin (2.83 g, 1.0 eq) was stirred in toluene (20 ml) and a solution of morpholine (1.93 ml, 1.5 eq) in toluene (20 ml) was added dropwise at -10-0° C. After stirring the resulting solution overnight at r.t., NH₄Cl (aq) was added to the solution and the solution was extracted

117

with EA. The combined organic layers were washed with brine, dried and evaporated in vacuo. A product was obtained as a white solid (2.7 g, 73.5%).


¹H NMR (500 MHz, CDCl₃-d₁): δ 1.41-1.44 (m, 3H), 3.75-3.77 (m, 4H), 3.83-3.85 (m, 4H), 4.01-4.05 (m, 2H), 7.69 (s, 1H)

4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine




¹H NMR (500 MHz, CDCl₃-d₁): δ 1.41-1.44 (m, 3H), 3.83 (m, 8H), 4.05-4.09 (m, 2H), 6.70-6.72 (m, 2H), 7.97 (s, 1H), 8.11-8.12 (m, 2H)

[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.34-1.37 (m, 3H), 3.72-3.78 (m, 8H), 4.09-4.13 (m, 2H), 7.24-7.29 (m, 3H), 7.42-7.46 (m, 2H), 7.58-7.60 (m, 2H), 8.12 (s, 1H), 8.21-8.23 (m, 2H), 10.40 (s, 1H)


4-{3-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide



118


¹H NMR (500 MHz, DMSO-d₆): δ 1.34-1.37 (m, 3H), 3.73-3.78 (m, 8H), 4.10-4.14 (m, 2H), 7.18 (m, 1H), 7.51-7.55 (m, 4H), 7.81-7.83 (m, 3H), 8.12 (s, 1H), 8.18-8.20 (m, 2H), 8.94 (s, 1H), 8.97 (s, 1H)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-thiomorpholin-4-yl-phenyl)-urea




¹H NMR (500 MHz, DMSO-d₆): δ 1.34-1.37 (m, 3H), 2.65-2.67 (m, 4H), 3.36-3.37 (m, 4H), 3.71-3.76 (m, 8H), 4.08-4.12 (m, 2H), 6.85-6.87 (m, 2H), 7.30-7.31 (m, 2H), 7.49-7.51 (m, 2H), 8.10-8.14 (m, 3H), 8.57 (s, 1H), 8.85 (s, 1H)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ 1.34-1.37 (m, 3H), 3.68-3.70 (m, 2H), 3.72-3.73 (m, 4H), 3.76-3.78 (m, 4H), 3.95-3.96 (m, 2H), 3.97-4.14 (m, 2H), 4.19 (s, 2H), 7.28-7.30 (m, 2H), 7.47-7.53 (m, 4H), 8.12 (s, 1H), 8.17-8.19 (m, 2H), 8.79 (s, 1H), 8.87 (s, 1H)


4-{3-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzenesulfonamide



119


¹H NMR (500 MHz, DMSO-d₆): δ 1.34-1.37 (m, 3H), 3.73-3.78 (m, 8H), 4.10-4.14 (m, 2H), 7.18 (m, 2H), 7.49-7.50 (m, 4H), 7.57-7.59 (m, 2H), 8.08 (s, 1H), 8.14-8.16 (m, 2H), 8.93 (s, 1H), 9.06 (s, 1H)

N-(4-{3-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-acetamide




¹H NMR (500 MHz, DMSO-d₆): δ 1.34-1.37 (m, 3H), 2.01 (s, 3H), 3.72-3.77 (m, 8H), 4.09-4.14 (m, 2H), 7.36-7.38 (m, 2H), 7.47-7.52 (m, 4H), 8.12 (s, 1H), 8.16-8.18 (m, 2H), 8.64 (s, 1H), 8.83 (s, 1H), 9.82 (s, 1H)

N-(4-{3-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-methanesulfonamide



¹H NMR (500 MHz, DMSO-d₆): δ 1.34-1.37 (m, 3H), 2.92 (s, 3H), 3.72-3.77 (m, 8H), 4.09-4.14 (m, 2H), 7.36-7.38 (m, 2H), 7.14-7.16 (m, 2H), 7.41-7.43 (m, 2H), 7.50-7.52 (m, 2H), 8.12 (s, 1H), 8.16-8.18 (m, 2H), 8.70 (s, 1H), 8.82 (s, 1H), 9.45 (s, 1H)


1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzooxazol-5-yl)-urea



120


¹H NMR (500 MHz, DMSO-d₆): δ 1.34-1.37 (m, 3H), 3.72-3.77 (m, 8H), 4.09-4.14 (m, 2H), 6.92 (m, 1H), 7.16-7.17 (m, 1H), 7.49-7.53 (m, 2H), 8.12 (s, 1H), 8.17-8.18 (m, 2H), 8.83 (s, 1H), 8.89 (s, 1H)

5 1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-methanesulfonyl-piperazin-1-yl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ 1.34-1.37 (m, 3H), 2.92 (s, 1H), 3.15-3.16 (m, 4H), 3.23-3.24 (m, 4H), 3.72-3.73 (m, 4H), 3.76-3.77 (m, 4H), 4.09-4.13 (m, 4H), 6.92-6.94 (m, 2H), 7.33-7.35 (m, 2H), 7.50-7.52 (m, 2H), 8.11 (s, 1H), 8.15-8.17 (m, 2H), 8.57 (s, 1H), 8.84 (s, 1H)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-carbonyl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ 1.34-1.37 (m, 3H), 3.42-3.43 (m, 4H), 3.49 (m, 4H), 3.72-3.73 (m, 4H), 3.76-3.77 (m, 4H), 4.09-4.13 (m, 2H), 7.35-7.37 (m, 2H), 7.51-7.54 (m, 4H), 8.12 (s, 1H), 8.17-8.19 (m, 2H), 8.92-8.96 (m, 2H)


1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzooxazol-6-yl)-urea



121


¹H NMR (500 MHz, DMSO-d₆): δ 1.34-1.37 (m, 3H), 3.73-3.74 (m, 4H), 3.76-3.77 (m, 4H), 4.10-4.14 (m, 2H), 6.99-7.01 (m, 1H), 7.06-7.08 (m, 1H), 7.51-7.53 (m, 2H), 7.58 (m, 1H), 8.12 (s, 1H), 8.16-8.18 (m, 2H), 8.73 (s, 1H), 8.73 (s, 1H), 11.46 (s, 1H)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-methanesulfonyl-phenyl)-urea




¹H NMR (500 MHz, DMSO-d₆): δ 1.34-1.37 (m, 3H), 3.16 (s, 3H), 3.73-3.74 (m, 4H), 3.77-3.78 (m, 4H), 4.10-4.14 (m, 2H), 7.54-7.56 (m, 2H), 7.69-7.71 (m, 2H), 7.82-7.84 (m, 2H), 8.12 (s, 1H), 8.19-8.20 (m, 2H), 9.03 (s, 1H), 9.25 (s, 1H)

5-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-1,3-dihydro-benzoimidazol-2-one



¹H NMR (500 MHz, DMSO-d₆): δ 1.34-1.37 (m, 3H), 3.16 (s, 3H), 3.73-3.74 (m, 4H), 3.77-3.78 (m, 4H), 4.10-4.14 (m, 2H), 6.95-6.97 (m, 1H), 7.83 (m, 1H), 7.93-7.94 (m, 1H), 8.11 (s, 1H)


N-(5-{3-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-pyridin-2-yl)-acetamide



122


¹H NMR (500 MHz, DMSO-d₆): δ 1.34-1.37 (m, 3H), 2.09 (s, 3H), 3.73-3.74 (m, 4H), 3.77-3.78 (m, 4H), 4.10-4.14 (m, 2H), 7.52-7.54 (m, 2H), 7.81-7.83 (m, 1H), 8.00-8.02 (m, 1H), 8.12 (s, 1H), 8.17-8.18 (m, 2H), 8.42-8.43 (m, 1H), 8.75 (s, 1H), 8.94 (s, 1H), 10.38 (s, 1H)

N-[3-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-methanesulfonamide




¹H NMR (500 MHz, DMSO-d₆): δ 1.35-1.37 (m, 3H), 2.99 (s, 3H), 3.73-3.74 (m, 4H), 3.77-3.78 (m, 4H), 4.13-4.14 (m, 2H), 7.26-7.28 (m, 1H), 7.38-7.41 (m, 1H), 7.97-7.99 (m, 1H), 8.16-8.18 (m, 2H), 9.83 (s, 1H)

N-[3-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-2,4-difluoro-benzenesulfonamide



¹H NMR (500 MHz, DMSO-d₆): δ 1.33-1.37 (m, 3H), 3.71-3.75 (m, 8H), 4.09-4.13 (m, 2H), 7.14-7.15 (m, 1H), 7.22-7.26 (m, 1H), 7.29-7.32 (m, 1H), 7.49-7.53 (m, 1H), 7.88-7.92 (m, 2H), 8.04 (s, 1H), 8.12 (s, 1H), 10.74 (s, 1H)


1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea



123


¹H NMR (500 MHz, DMSO-d₆): δ1.33-1.37 (m, 3H), 3.71-3.72 (m, 4H), 3.76-3.77 (m, 4H), 4.08-4.12 (m, 2H), 7.44-7.45 (m, 2H), 7.52-7.54 (m, 2H), 8.11 (s, 1H), 8.18-8.19 (m, 2H), 8.35 (m, 1H), 9.04 (s, 1H), 9.15 (s, 1H)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea hydrochloride



¹H NMR (500 MHz, DMSO-d₆): δ1.33-1.37 (m, 3H), 3.71-3.72 (m, 4H), 3.76-3.77 (m, 4H), 4.08-4.12 (m, 2H), 7.44-7.45 (m, 2H), 7.52-7.54 (m, 2H), 8.11 (s, 1H), 8.18-8.19 (m, 2H), 8.35 (m, 1H), 9.04 (s, 1H), 9.15 (s, 1H)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-sulfonyl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ1.33-1.37 (m, 3H), 2.84 (m, 4H), 3.63 (m, 4H), 3.73-3.74 (m, 4H), 3.77-3.78 (m, 4H), 4.10-4.14 (m, 2H), 7.54 (m, 2H), 7.64-7.66 (m, 2H), 7.71-7.73 (m, 2H), 8.13 (s, 1H), 8.19-8.20 (m, 2H), 9.05 (s, 1H), 9.29 (s, 1H)

Example 4

Preparation of Compounds of Formula (I) in Scheme 2

2,4,6-Trichloro-5-methylsulfanyl-pyrimidine



Barbituric acid (5.2 g, 40.6 mmol.), 3.5 ml of DMSO, 20 ml of acetic acid, and 6.0 ml of acetic anhydride were heated progressively to 90-100° C. This temperature was maintained

124


for 4 hr and then 130 ml of water were added to the mixture. After cooling and filtering, the precipitate was washed with acetone. The dimethylsulfonium-substituted barbituric acid weighed 6.1 g (80.1%).

¹H NMR (500 MHz, DMSO-d₆): δ10.14 (s, 2H), 2.96 (s, 6H)

The above product (5.2 g, 27.5 mmol.) was reacted with 30 ml of phosphorus oxychloride and 5.0 ml of dimethylaniline. After heating for 24 hr under reflux, the reaction mixture was subjected to hydrolysis in iced salt water. The reaction mixture was subjected to filtration and the resulting solid was washed with water. After drying in vacuo, a product of 4.5 g (60.1%) was obtained.

¹H NMR (500 MHz, CDCl₃-d₁): δ2.80 (s, 3H)


4-(2,6-Dichloro-5-methylsulfanyl-pyrimidin-4-yl)-morpholine



A mixture of 2,4,6-trichloro-5-methylsulfanyl-pyrimidine (4.0 g, 17.4 mmol.), morpholine (2.3 g, 26.1 mmol, 1.5 eq.) and THF (20 ml) was stirred at room temperature for 2.0 hr. The reaction mixture was dried and the solvent was evaporated. Thereafter, a 0.1 N HCl washing free morpholine was added to the solution. The solution was extracted with EA and the EA in the solution was removed by evaporation. A crude product was obtained. Purification of the crude by flash chromatography on silica gel (Hexane/EtOAc 6:1) gave a gray solid of 2.34 g (48.2%).

¹H NMR (500 MHz, CDCl₃-d₁): δ3.89 (t, 4H), 3.80 (t, 4H), 2.32 (s, 3H)

4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenylamine




A mixture of 4-(2,6-dichloro-5-methylsulfanyl-pyrimidin-4-yl)-morpholine (200 mg, 0.71 mmol.), 4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-aniline (172.0 mg, 0.781 mmol.), (PPh₃)₄Pd (437 mg, 0.07 mmol.), 2M Na₂CO₃ (1.0 ml, 2.13 mmol, 3.0 eq.) and dioxane (15.0 ml) was added to the reaction vessel which was flushed with argon. The reac-

125

tion mixture was stirred at 90° C. for 12 hrs and then dried so that the solvent was evaporated. The reaction mixture was partitioned between EA and water. The organic layers were collected, washed with brine, dried over MgSO₄, filtered and evaporated in vacuo. The resulting residue was purified by


¹H NMR (500 MHz, DMSO-d₆): δ8.19 (d, 2H), 6.72 (d, 2H), 3.85 (s, 8H), 2.35 (s, 3H).

4-[6-Chloro-2-(3-methoxy-phenyl)-5-methylsulfa-
nyl-pyrimidin-4-yl]-morpholine




¹H NMR (500 MHz, CDCl₃-d₁): δ7.96 (d, 1H), 7.90 (d, 1H), 7.36 (t, 1H), 7.01 (d, 1H), 3.87 (m, 11H), 2.36 (s, 3H)

4-(4-Chloro-5-methylsulfonyl-6-morpholin-4-yl-
pyrimidin-2-yl)-phenol



¹H NMR (500 MHz, CDCl₃-d₁): δ8.27 (d, 2H), 6.88 (d, 2H), 5.08 (s, 1H), 3.86 (s, 8H), 2.35 (s, 3H)


3-(4-Chloro-5-methylsulfonyl-6-morpholin-4-yl-
pyrimidin-2-yl)-phenol



126


¹H NMR (500 MHz, CDCl₃-d₁): δ7.93 (d, 1H), 7.82 (dd, 1H), 7.32 (t, 1H), 6.96 (m, 1H), 5.31 (b, 1H), 3.87 (q, 8H), 2.36 (s, 3H)

4-(4-Chloro-5-methylsulfonyl-6-morpholin-4-yl-
pyrimidin-2-yl)-benzenesulfonamide




¹H NMR (500 MHz, Acetone-d₆): δ8.52 (d, 2H), 7.99 (d, 2H), 6.71 (s, 2H), 3.83 (t, 8H), 2.41 (s, 3H)

5-(4-Chloro-5-methylsulfonyl-6-morpholin-4-yl-
pyrimidin-2-yl)-pyridin-2-ylamine



¹H NMR (500 MHz, CDCl₃-d₁): δ9.05 (d, 1H), 8.36 (dd, 1H), 6.53 (s, 1H), 4.87 (b, 2H), 3.85 (t, 8H), 2.35 (s, 3H)


5-(4-Chloro-5-methylsulfonyl-6-morpholin-4-yl-
pyrimidin-2-yl)-1H-indole




¹H NMR (500 MHz, CDCl₃-d₁): δ8.72 (d, 1H), 8.28-8.24 (dd, 2H), 7.43-7.41 (d, 1H), 7.25 (dd, 1H), 7.25-7.24 (d, 1H), 3.89 (m, 8H), 2.37 (s, 3H)

127

4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-1H-indazole

¹H NMR (500 MHz, CDCl₃-d₁): δ8.98 (s, 1H), 8.30 (d, 1H), 7.64 (d, 1H), 7.49 (s, 1H), 3.91 (s, 8H), 2.41 (s, 1H)


N-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-methanesulfonamide




A mixture of 4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (30 mg, 0.091 mmol.), methanesulfonyl chloride (15.5 mg, 0.136 mmol, 1.5 eq.) and CH₂Cl₂ was added to the reaction vessel and flushed with argon. The reaction mixture was stirred at r.t. for 2 hrs. The reaction mixture was partitioned between CH₂Cl₂ and NaHCO₃ (aq), the organic layer was washed with brine, dried over MgSO₄, filtered and evaporated in vacuo. A white solid of 30.3 mg (82.1%) was obtained as the product.

¹H NMR (500 MHz, CDCl₃-d₁): δ8.34 (d, 2H), 7.26 (d, 1H), 6.72 (s, 1H), 3.87 (d, 8H), 3.06 (s, 3H), 2.35 (t, 3H)


N-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-acetamide

**128**¹H NMR (500 MHz, CDCl₃-d₁): δ10.25 (s, 1H), 8.21 (d, 2H), 7.72 (d, 2H), 3.84 (t, 4H), 3.84 (t, 4H), 3.75 (t, 4H), 2.33 (s, 3H), 2.08 (s, 3H)


N-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-benzenesulfonamide

¹H NMR (500 MHz, CDCl₃-d₁): δ8.24 (d, 2H), 7.80 (t, 2H), 7.53 (t, 1H), 7.44 (t, 2H), 7.14 (d, 2H), 6.75 (s, 1H), 3.84 (d, 8H), 2.34 (d, 3H)

Methanesulfonic acid-4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester


¹H NMR (500 MHz, CDCl₃-d₁): δ8.42 (m, 2H), 7.36 (q, 2H), 3.86 (m, 8H), 3.17 (s, 3H), 2.37 (s, 3H)

Benzenesulfonic acid-4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester

¹H NMR (500 MHz, CDCl₃-d₁): δ8.29-8.27 (d, 2H), 7.84-7.83 (d, 2H), 7.68-7.65 (d, 1H), 7.53-7.50 (d, 2H), 7.05-7.04 (s, 1H), 3.86 (t, 8H), 3.18 (s, 3H), 2.36 (s, 3H)


129

Methanesulfonic acid-3-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ 8.33 (d, 1H), 8.23 (d, 1H), 7.49 (d, 1H), 7.42 (d, 1H), 3.86 (t, 8H), 3.18 (s, 3H), 2.36 (s, 3H)


1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-chloro-3-trifluoromethyl-phenyl)-urea



A mixture of 4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (40 mg, 0.118 mmol.), 2-chloro-4-isocyanato-1-trifluoromethyl-benzene (39 mg, 0.178 mmol.), CH₂Cl₂ (1.0 ml), and toluene (2.0 ml) was added to a reaction vessel which was flushed with argon. The reaction mixture was stirred at 90° C. for 16 hrs. After cooling and filtering, the precipitate was washed with a minor portion of CH₂Cl₂ again to give a white solid of 26.7 mg (40.3%).

¹H NMR (500 MHz, Acetone-d₆): δ 8.66 (s, 1H), 8.59 (s, 1H), 8.30 (dd, 2H), 8.16 (d, 1H), 7.77 (dd, 1H), 7.66 (dd, 2H), 7.56 (d, 1H), 3.90 (t, 4H), 3.82 (t, 4H), 2.39 (s, 3H)


1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-chloro-pyridin-4-yl)-urea



130


¹H NMR (500 MHz, Acetone-d₆): δ 8.72 (s, 1H), 8.61 (s, 1H), 8.50 (d, 2H), 8.32 (s, 1H), 7.67 (d, 2H), 7.38 (d, 1H), 3.91 (t, 4H), 3.82 (t, 4H), 2.39 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea




¹H NMR (500 MHz, CDCl₃-d₁): δ 8.27 (d, 2H), 7.43 (d, 2H), 7.34 (t, 4H), 7.12 (s, 1H), 7.05 (s, 1H), 6.90 (s, 1H), 3.96 (d, 8H), 2.35 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-dimethylamino-phenyl)-urea



¹H NMR (500 MHz, CDCl₃-d₁): δ 8.31 (b, 1H), 8.29 (s, 1H), 7.89 (b, 1H), 8.27 (s, 1H), 7.91 (s, 1H), 7.66 (d, 2H), 7.36 (d, 2H), 6.73 (d, 2H), 3.90 (t, 4H), 3.82 (t, 4H), 2.80 (d, 6H), 2.39 (s, 3H)


1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-fluoro-phenyl)-urea



131


¹H NMR (500 MHz, DMSO-d₆): δ 9.04 (s, 1H), 8.82 (s, 1H), 8.21-8.19 (d, 2H), 7.58 (d, 2H), 7.49-7.46 (dd, 2H), 7.15-7.11 (t, 2H), 3.84 (d, 4H), 3.76 (d, 4H), 2.36 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3-fluoro-phenyl)-urea




¹H NMR (500 MHz, Acetone-d₆): δ 8.56 (s, 1H), 8.52 (s, 1H), 8.31-8.30 (d, 2H), 7.68-7.66 (d, 2H), 7.62-7.60 (d, 1H), 7.30-7.29 (d, 2H), 7.20 (s, 1H), 3.71 (t, 4H), 3.82 (t, 4H), 2.39 (s, 3H).

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3,4-difluoro-phenyl)-urea



¹H NMR (500 MHz, Acetone-d₆): δ 8.77 (s, 1H), 8.73 (s, 2H), 8.30 (d, 2H), 7.82 (m, 1H), 7.66 (d, 2H), 7.21 (m, 1H), 3.91 (t, 4H), 3.82 (t, 4H), 2.39 (s, 3H)

[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea




Trimethylsilyl isocyanate (136 mg, 5 eq.) was added to a stirred solution of 4-(4-Chloro-5-methylsulfanyl-6-morpho-

132

lin-4-yl-pyrimidin-2-yl)-phenylamine (80 mg, 0.237 mmol.) in THF (2.0 ml) and the mixture was reacted to reflux for 16 hrs. The reaction mixture was extracted with EA and washed with brine. The crude was purified by Chromatography (Hexane/EtOAc 1:1) to give a product 28.1 mg (31.2%).


¹H NMR (500 MHz, CDCl₃-d₁): δ 8.58 (s, 1H), 8.33 (s, 2H), 7.66 (s, 2H), 6.42 (b, 2H), 3.92 (t, 4H), 3.82 (t, 4H), 2.39 (s, 3H)

1-(2-Chloro-ethyl)-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea




¹H NMR (500 MHz, CDCl₃-d₁): δ 8.26 (d, 2H), 7.40 (d, 2H), 7.12 (s, 1H), 5.58 (t, 1H), 3.86 (d, 8H), 3.62 (m, 4H), 2.35 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-ethyl-urea



¹H NMR (500 MHz, CDCl₃-d₁): δ 8.26 (d, 2H), 7.39 (d, 2H), 6.97 (s, 1H), 5.05 (b, 1H), 3.85 (d, 8H), 3.29 (d, 2H), 2.35 (s, 3H), 1.15 (t, 3H)


1-[5-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-pyridin-2-yl]-3-phenyl-urea



133


¹H NMR (500 MHz, DMSO-d₆): δ10.4 (s, 1H), 9.76 (s, 1H), 9.14 (s, 1H), 8.50 (dd, 1H), 7.66 (d, 1H), 7.55 (d, 2H), 7.32 (t, 2H), 7.04 (s, 1H), 3.87 (t, 4H), 3.76 (t, 4H), 2.36 (d, 3H)

1-[5-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-pyridin-2-yl]-3-(3-fluoro-phenyl)-urea




¹H NMR (500 MHz, CDCl₃-d₁): δ12.1 (s, 1H), 9.25 (s, 1H), 8.57 (dd, 1H), 8.44 (s, 1H), 7.59 (d, 1H), 7.28 (dd, 2H), 6.90 (s, 1H), 6.80 (s, 1H), 3.88 (t, 8H), 2.38 (s, 3H)

1-[5-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-pyridin-2-yl]-3-ethyl-urea



¹H NMR (500 MHz, DMSO-d₆): δ9.52 (s, 1H), 9.04 (s, 1H), 8.41 (dd, 1H), 8.00 (b, 1H), 7.51 (d, 1H), 3.85 (t, 4H), 3.75 (t, 4H), 3.20 (t, 2H), 2.34 (s, 3H), 1.10 (t, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-imidazolidin-2-one




A mixture of 1-(2-chloro-ethyl)-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (20

134

mg, 0.045 mmol.), 10% NaOH (0.5 ml), THF (2.0 ml), and H₂O (1.0 ml) was added to a vessel. The reaction mixture was stirred at 80° C. for 8 hr and then dried, and the solvent was evaporated. The reaction mixture was partitioned between EA and 0.1N HCl, the organic layer was washed with brine, dried over MgSO₄, filtered and evaporated in vacuo. The resulting residue was purified by flash chromatography on silica gel (Hexane/EtOAc 3:1) to give a gray solid of 11.4 mg (62.3%).

¹H NMR (500 MHz, CDCl₃-d₁): δ8.33 (d, 2H), 7.64 (d, 2H), 4.88 (s, 1H), 4.01 (m, 2H), 3.87 (t, 8H), 3.66 (m, 2H), 2.36 (s, 3H)


(2-Chloro-ethyl)-carbamic acid 4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester



A mixture of 4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenol (80.0 mg, 0.236 mmol), 1-chloro-2-isocyanato-ethane (43.1 mg, 0.354 mmol, eq=1.5), toluene and CH₂Cl₂ was added to the reaction vessel and flushed with argon. The reaction mixture was stirred at 80° C. for 16 hrs and then dried, and the solvent was evaporated. The reaction mixture was partitioned between EA and water, the organic layer was washed with brine, dried over MgSO₄, filtered and evaporated in vacuo. The resulting residue was purified by flash chromatography on silica gel (Hexane/EtOAc 2:1) to give a gray solid of 58.9 mg (58.1%)

¹H NMR (500 MHz, CDCl₃-d₁): δ8.36 (d, 2H), 7.23 (d, 2H), 5.50 (s, 1H), 3.87 (d, 8H), 3.70 (t, 2H), 3.64 (dd, 2H), 2.36 (s, 3H)

[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester




A mixture of 4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (915 mg, 2.717 mmol.) in EA (20 ml) and NaHCO₃ (20 ml) the reaction was stirred at r.t. then added dropwise to slurry of phenyl chloroformate (851 mg, 0.543 mmol, 2.0 eq) at reaction for 2 hr. The reaction

135

mixture was partitioned between EA/NaHCO₃, the organic layer was washed with brine, dried over MgSO₄, filtered and evaporated in vacuo. The residue was recrystallized by (Hexane/EtOAc) to give a pale yellow product of 1056 mg (85.1%).


¹H NMR (500 MHz, CDCl₃-d₁): δ8.35-8.33 (d, 2H), 7.54-7.52 (d, 2H), 7.41-7.38 (dd, 2H), 7.26-7.18 (m, 4H), 3.87 (s, 8H), 2.36 (s, 3H)

[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenyl]-carbamic acid phenyl ester




¹H NMR (500 MHz, CDCl₃-d₁): δ8.18 (b, 1H), 8.14-8.11 (m, 2H), 7.43-7.40 (dd, 2H), 7.35 (s, 1H), 7.28-7.26 (s, 1H), 7.22-7.20 (s, 2H), 3.88-3.87 (s, 4H), 2.36 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-methyl-urea



¹H NMR (500 MHz, DMSO-d₆): δ8.28 (d, 2H), 7.40 (d, 2H), 6.82 (b, 1H), 5.91 (b, 1H), 3.85 (d, 8H), 2.86 (d, 3H), 2.35 (s, 3H)


1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-morpholin-4-yl-urea



136


¹H NMR (500 MHz, CDCl₃-d₁): δ8.32-8.31 (d, 2H), 8.23 (s, 1H), 7.59-7.57 (d, 2H), 5.87 (s, 1H), 4.01 (b, 2H), 3.87-3.87 (d, 8H), 3.72 (b, 2H), 3.09 (b, 2H), 2.77 (b, 2H), 2.36 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea




¹H NMR (500 MHz, DMSO-d₆): δ9.31 (s, 1H), 9.10 (s, 1H), 8.63 (d, 1H), 8.20 (t, 3H), 7.95 (dd, 1H), 7.55 (d, 2H), 7.32 (s, 1H), 3.84 (d, 4H), 3.77 (d, 4H), 2.36 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea



¹H NMR (500 MHz, DMSO-d₆): δ9.27-9.25 (d, 2H), 8.38-8.36 (d, 2H), 8.23-8.22 (d, 2H), 7.61-7.59 (d, 2H), 7.46-7.45 (d, 2H), 3.85-3.84 (d, 4H), 3.77-3.75 (d, 4H), 2.34 (s, 3H)


1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-morpholin-4-yl-ethyl)-urea



137


¹H NMR (500 MHz, CDCl₃-d₁): δ8.25 (d, 2H), 7.89 (b, 1H), 7.44 (d, 2H), 5.88 (b, 1H), 3.85 (d, 8H), 3.65 (t, 4H), 3.55 (dd, 2H), 2.51 (d, 2H), 2.47 (s, 4H), 2.34 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(2-di methylamino-ethoxy)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ9.63 (s, 1H), 9.31 (s, 1H), 8.19-8.18 (d, 2H), 7.59-7.57 (d, 2H), 7.44-7.42 (d, 2H), 6.97-6.95 (d, 2H), 4.45 (s, 2H), 3.90 (s, 2H), 3.84-3.38 (dd, 8H), 3.24 (s, 6H), 2.34 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-thiazol-2-yl-urea




¹H NMR (500 MHz, DMSO-d₆): δ11.01 (b, 1H), 9.72 (b, 1H), 8.23 (d, 2H), 7.65 (d, 2H), 7.35 (s, 1H), 7.11 (s, 1H), 3.84 (d, 4H), 3.77 (d, 4H), 2.36 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-urea

**138**


¹H NMR (500 MHz, DMSO-d₆): δ8.32-8.30 (d, 2H), 8.25 (d, 1H), 7.58-7.56 (d, 2H), 5.54 (b, 1H), 3.87-3.85 (d, 8H), 3.09 (b, 2H), 2.89 (b, 2H), 2.65 (b, 2H), 2.35 (s, 3H), 2.34 (b, 2H) 2.17 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3-fluoro-4-morpholin-4-yl-phenyl)-urea




¹H NMR (500 MHz, DMSO-d₆): δ9.16 (s, 1H), 8.93 (s, 1H), 8.20-8.19 (d, 2H), 7.59-7.57 (d, 2H), 7.47-7.44 (d, 1H), 7.10-7.08 (d, 1H), 7.00 (d, 1H), 3.84-3.83 (d, 4H), 3.76-3.72 (dd, 8H), 2.94-2.93 (d, 4H), 2.34 (s, 3H)

4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzoic acid ethyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ8.17 (t, 2H), 8.09 (s, 1H), 8.01 (s, 1H), 7.90 (d, 2H), 7.43 (t, 4H), 4.33 (dd, 2H), 3.86 (s, 4H), 3.83 (s, 4H), 2.34 (s, 3H), 1.36 (dd, 3H)


1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea



139


¹H NMR (500 MHz, DMSO-d₆): δ 9.18 (s, 1H), 8.98 (d, 1H), 8.21-8.19 (d, 2H), 7.61-7.59 (d, 2H), 7.51-7.49 (d, 2H), 7.30-7.28 (d, 2H), 4.18 (s, 2H), 3.97-3.95 (dd, 2H), 3.84-3.76 (d, 4H), 3.76-3.75 (d, 4H), 3.70-3.68 (dd, 2H), 2.36 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3-trifluoromethyl-phenyl)-urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.19-9.18 (d, 2H), 8.23-8.21 (d, 2H), 8.03 (s, 1H), 7.62-7.58 (t, 3H), 7.54-7.53 (d, 1H), 7.34-7.33 (d, 1H), 3.84-3.83 (d, 4H), 3.77-3.75 (d, 4H), 2.34 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-methyl-piperazin-1-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ 9.02 (s, 1H), 8.60 (s, 1H), 8.19-8.18 (d, 2H), 7.58-7.56 (d, 2H), 7.31-7.30 (d, 2H), 6.89-6.87 (d, 2H), 3.84-3.83 (d, 4H), 3.76-3.75 (d, 4H), 3.05 (s, 4H), 2.45-2.44 (d, 4H), 2.36-2.34 (d, 3H), 2.34 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-thiomorpholin-4-yl-phenyl)-urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.03 (s, 1H), 8.61 (s, 1H), 8.20-8.18 (d, 2H), 7.58-7.56 (d, 2H), 7.33-7.31 (d, 2H),

140


6.89-6.87 (d, 2H), 3.84-3.83 (d, 4H), 3.76-3.75 (d, 4H), 3.40-3.38 (s, 4H), 2.69-2.67 (d, 4H), 2.32 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-morpholin-4-yl-phenyl)-urea




¹H NMR (500 MHz, DMSO-d₆): δ 8.96 (s, 1H), 8.54 (s, 1H), 8.16-8.14 (d, 2H), 7.54-7.52 (d, 2H), 7.29-7.28 (d, 2H), 6.86-6.84 (d, 2H), 3.80-3.79 (d, 4H), 3.72-3.69 (d, 8H), 2.99-2.97 (d, 4H), 2.32 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ 9.12 (s, 1H), 8.60 (s, 1H), 8.20-8.18 (d, 2H), 8.16 (d, 1H), 7.70-7.60 (d, 1H), 7.58-7.56 (d, 2H), 6.82-6.81 (d, 1H), 3.84-3.83 (d, 4H), 3.76-3.75 (d, 4H), 3.43-3.39 (d, 4H), 2.40-2.39 (d, 4H), 2.38-2.36 (d, 3H), 2.30 (s, 3H)


1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(6-morpholin-4-yl-pyridin-3-yl)-urea



141


¹H NMR (500 MHz, DMSO-d₆): δ 9.21 (s, 1H), 8.72 (s, 1H), 8.20-8.18 (d, 3H), 7.74-7.72 (d, 1H), 7.59-7.57 (d, 2H), 6.84-6.82 (d, 1H), 3.84-3.83 (d, 4H), 3.76-3.75 (d, 4H), 3.71-3.69 (d, 4H), 3.32-3.30 (d, 4H), 2.34 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-2-yl-urea




¹H NMR (500 MHz, DMSO-d₆): δ 10.73 (s, 1H), 9.53 (s, 1H), 8.30-8.29 (d, 1H), 8.24-8.22 (d, 2H), 7.78-7.75 (dd, 1H), 7.67-7.65 (d, 2H), 7.56-7.55 (d, 1H), 7.05-7.02 (d, 1H), 3.85-3.84 (d, 4H), 3.77-3.76 (d, 4H), 2.36-2.34 (s, 3H)

2-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-thiophene-3-carboxylic acid methyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ 9.72 (s, 1H), 8.32-8.30 (d, 2H), 8.06-8.05 (d, 1H), 7.57-7.55 (dd, 2H), 7.48-7.46 (d, 1H), 3.91 (s, 4H), 3.90-3.88 (d, 3H), 3.87-3.86 (d, 4H), 2.36 (s, 3H)


1-Benzo[1,3]dioxol-5-yl-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea



142


¹H NMR (500 MHz, DMSO-d₆): δ 9.13 (s, 1H), 8.82 (s, 1H), 8.20-8.18 (d, 2H), 7.58-7.56 (d, 2H), 7.21 (s, 1H), 6.84-6.78 (dd, 2H), 5.97 (s, 2H), 3.84-3.83 (d, 4H), 3.76-3.76 (d, 4H), 2.36-2.34 (s, 3H)

5 4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide




¹H NMR (500 MHz, DMSO-d₆): δ 9.13 (s, 1H), 9.04 (s, 1H), 8.23-8.21 (d, 2H), 7.84-7.82 (d, 3H), 7.61-7.59 (d, 2H), 7.54-7.52 (d, 2H), 7.20 (b, 1H), 3.85-3.84 (d, 4H), 3.77-3.76 (d, 4H), 2.34 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3,4-dimethoxy-phenyl)-urea



¹H NMR (500 MHz, DMSO-d₆): δ 8.94 (s, 1H), 8.60 (s, 1H), 8.21-8.19 (d, 2H), 7.59-7.57 (d, 2H), 7.21 (s, 1H), 6.88-6.87 (s, 2H), 3.84 (s, 3H), 3.75 (s, 8H), 3.71 (s, 3H), 2.36 (s, 3H)


1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[5-(4-methyl-piperazin-1-yl)-pyridin-2-yl]-urea



143


¹H NMR (500 MHz, DMSO-d₆): δ10.82 (b, 1H), 9.26 (s, 1H), 8.22-8.20 (d, 2H), 7.95 (s, 1H), 7.64-7.63 (d, 2H), 7.46-7.44 (t, 2H), 3.84-3.77 (d, 4H), 3.76-3.75 (d, 4H), 3.10-3.09 (t, 4H), 2.52-2.49 (d, 4H), 2.46-2.45 (s, 3H), 2.34-2.21 (s, 3H)

N-(3-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-2,2-trifluoro-acetamide




¹H NMR (500 MHz, DMSO-d₆): δ9.33 (3, 1H), 9.29 (b, 1H), 8.58 (s, 1H), 8.23-8.21 (d, 2H), 7.86-7.85 (d, 1H), 7.79-7.76 (d, 1H), 7.63-7.58 (m, 3H), 3.84-3.76 (d, 4H), 3.75-3.75 (d, 4H), 2.36-2.34 (s, 3H)

N-(3-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-acetamide



¹H NMR (500 MHz, DMSO-d₆): δ9.93 (s, 1H), 9.05 (s, 1H), 8.91 (s, 1H), 8.21-8.19 (d, 2H), 7.78 (s, 1H), 7.59-7.58 (d, 2H), 7.21-7.17 (dd, 3H), 3.84-3.83 (d, 4H), 3.77-3.76 (d, 4H), 2.36-2.34 (s, 3H), 2.03-1.98 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-methyl-piperazine-1-carbonyl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ9.21 (s, 1H), 9.13 (s, 1H), 8.22-8.21 (d, 2H), 7.60-7.59 (d, 2H), 7.54-7.53 (d, 2H),

144

7.37-7.36 (d, 2H), 3.84 (d, 4H), 3.76 (d, 4H), 3.39-3.35 (d, 4H), 2.50-2.46 (d, 4H), 2.36-2.34 (s, 3H), 2.03-1.98 (s, 3H)

1-(6-Bromo-pyridin-3-yl)-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea

5

10

15

20

25

30

35

40


45

50

55


60

65




¹H NMR (500 MHz, DMSO-d₆): δ9.71 (b, 2H), 8.51-8.50 (d, 1H), 8.22-8.20 (d, 2H), 7.93-7.91 (dd, 1H), 7.62-7.60 (d, 2H), 7.56-7.54 (d, 1H), 3.84-3.76 (d, 4H), 3.76 (s, 4H), 2.36-2.34 (s, 3H)

4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzoic acid



¹H NMR (500 MHz, DMSO-d₆): δ9.39-9.38 (d, 1H), 9.33-9.32 (d, 1H), 8.22-8.20 (d, 2H), 7.90-7.88 (dd, 2H), 7.61-7.59 (dd, 4H), 3.84-3.81 (d, 4H), 3.76-3.74 (d, 4H), 2.33 (s, 3H)

1-(2-Amino-phenyl)-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thio-urea




145

A mixture of bis-(2,5-dihydro-imidazol-1-yl)-methanethione (29.2 mg, 0.164 mmol), imidazole (3.7 mg, 0.05 mmol, 0.5 eq) in CH₃CN (5 ml) was stirred at 0° C., then added dropwise to a slurry of 4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (50 mg, 0.109 mmol.). The resulting mixture was reacted 3.0 hrs at r.t. and then benzene-1,2-diamine (23.6 mg, 0.218 mmol, 2.0 eq) was added thereto. The reaction resulted in a white solid, which was filtered, collected and evaporated in vacuo to give 49.5 mg of a white compound (68.5%).


¹H NMR (500 MHz, DMSO-d₆): δ 10.01 (b, 1H), 9.65 (b, 1H), 8.22-8.20 (d, 2H), 7.75-7.74 (d, 2H), 7.10-7.09 (d, 1H), 6.97-6.95 (d, 1H), 6.75-6.74 (d, 1H), 6.58-6.55 (d, 1H), 4.95 (b, 2H), 3.85-3.84 (d, 4H), 3.76-3.75 (d, 4H), 2.36-2.34 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-thiourea



¹H NMR (500 MHz, DMSO-d₆): δ 8.63-8.63 (d, 1H), 8.44-8.43 (d, 1H), 8.26-8.24 (d, 2H), 7.98-7.97 (d, 1H), 7.70-7.66 (m, 4H), 7.12-7.04 (d, 1H), 3.86-3.85 (d, 4H), 3.76-3.75 (d, 4H), 2.35 (s, 3H)


1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-thiourea



¹H NMR (500 MHz, DMSO-d₆): δ 10.10 (b, 1H), 10.01 (b, 1H), 8.23-8.22 (d, 2H), 7.70-7.68 (d, 2H), 7.53-7.50 (d, 2H), 7.36-7.34 (d, 2H), 4.20-3.98 (d, 2H), 3.97-3.96 (d, 2H), 3.85-3.75 (dd, 4H), 3.74-3.71 (dd, 6H), 2.36-2.34 (s, 3H)

146


(1H-Benzoimidazol-2-yl)-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-amine



A mixture of 1-(2-amino-phenyl)-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thiourea (120 mg, 1.0 eq.) and DCC (0.18 g, 0.271 mmol, 1.1 eq.) in THF (8 ml) was reacted for 9 hrs at 80-90° C. The resulting mixture was dried in vacuo, crystallized in EA solution, filtered to give a yellow solid (69.5 mg, 62.3%), which was then dried in vacuo to give a product.


¹H NMR (500 MHz, DMSO-d₆): δ 10.78 (b, 1H), 9.98 (s, 1H), 8.25-8.23 (d, 2H), 7.90-7.89 (d, 2H), 7.24-7.20 (d, 2H), 7.02 (s, 2H), 3.85-3.84 (d, 4H), 3.77-3.75 (d, 4H), 2.34 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-isopropyl-urea



¹H NMR (500 MHz, DMSO-d₆): δ 8.67 (s, 1H), 8.15-8.13 (d, 2H), 7.50-7.49 (d, 2H), 6.02 (s, 1H), 3.84-3.83 (d, 4H), 3.76-3.75 (d, 4H), 2.33 (s, 3H), 1.11-1.09 (s, 6)


1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-carbonyl)-phenyl]-urea



147


¹H NMR (500 MHz, DMSO-d₆): δ9.34-9.24 (d, 2H), 8.22-8.20 (d, 2H), 7.61-7.60 (d, 2H), 7.55-7.53 (d, 2H), 7.37-7.36 (d, 2H), 3.84-3.83 (d, 4H), 3.77-3.76 (d, 4H), 3.60-3.39 (m, 8H), 2.36-2.34 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(5-morpholin-4-yl-pyridin-2-yl)-urea




¹H NMR (500 MHz, DMSO-d₆): δ10.31 (b, 1H), 9.27 (s, 1H), 8.23-8.21 (d, 2H), 7.97 (s, 1H), 7.65-7.63 (d, 2H), 7.47 (s, 2H), 3.85-3.83 (d, 4H), 3.77-3.74 (dd, 8H), 3.09-3.07 (d, 4H), 2.49-2.34 (s, 3H)

1-(4-Amino-phenyl)-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ9.09 (s, 1H), 8.48 (s, 1H), 8.18-8.16 (d, 2H), 7.57-7.55 (d, 2H), 7.10-7.08 (d, 2H), 6.52-6.50 (d, 2H), 4.79 (s, 2H), 3.83-3.83 (d, 4H), 3.76-3.75 (d, 4H), 2.49-2.46 (s, 3H)

1-tert-Butyl-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea

**148**


¹H NMR (500 MHz, DMSO-d₆): δ8.60 (s, 1H), 8.14-8.13 (d, 2H), 7.48-7.46 (d, 2H), 6.12 (s, 1H), 3.83-3.81 (d, 4H), 3.76-3.74 (d, 4H), 2.33 (s, 3H), 1.29-1.26 (d, 9H)

4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenylamine




¹H NMR (500 MHz, CDCl₃-d₁): δ8.11-8.06 (dd, 2H), 7.15-7.14 (m, 1H), 3.87-3.86 (d, 8H), 2.36 (s, 3H)

4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-3-fluoro-phenylamine



¹H NMR (500 MHz, CDCl₃-d₁): δ8.04-8.01 (dd, 1H), 6.74-6.72 (d, 1H), 6.67-6.65 (d, 1H), 3.89 (s, 4H), 3.84-3.83 (s, 4H), 2.36 (s, 3H)


6-Chloro-5-methylsulfanyl-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-ylamine



149


¹H NMR (500 MHz, DMSO-d₆): δ8.99 (s, 2H), 7.35 (s, 2H), 3.84-3.83 (d, 4H), 3.74-3.73 (d, 4H), 2.32 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(1H-indazol-4-yl)-urea




¹H NMR (500 MHz, DMSO-d₆): δ10.45 (s, 1H), 8.55 (s, 1H), 8.29-8.26 (d, 2H), 7.98-7.95 (d, 2H), 7.67 (s, 1H), 7.43-7.41 (d, 1H), 7.26-7.23 (d, 1H), 6.42-6.41 (d, 1H), 6.12 (s, 1H), 3.87-3.85 (d, 4H), 3.77-3.76 (d, 4H), 2.50-2.50 (s, 3H)

N-(4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-methanesulfonamide




¹H NMR (500 MHz, DMSO-d₆): δ9.87 (b, 1H), 9.03 (s, 1H), 8.78 (s, 1H), 8.21-8.19 (d, 2H), 7.59-7.57 (d, 2H), 7.44-7.42 (d, 2H), 7.16-7.14 (d, 2H), 3.85-3.83 (d, 4H), 3.77-3.75 (d, 4H), 2.92 (s, 3H), 2.34 (s, 3H)

[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenyl]-urea

**150**


¹H NMR (500 MHz, DMSO-d₆): δ8.64 (s, 1H), 8.34-8.31 (s, 1H), 8.01-7.96 (dd, 2H), 6.34 (s, 2H), 3.84-3.83 (d, 4H), 3.76-3.75 (d, 4H), 2.34 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ9.31 (s, 1H), 8.92 (s, 1H), 8.37-8.34 (s, 1H), 8.08-8.01 (dd, 2H), 7.50-7.48 (d, 2H), 7.32-7.31 (d, 2H), 4.19 (s, 2H), 3.97-3.95 (s, 2H), 3.85-3.85 (d, 4H), 3.77-3.75 (d, 4H), 3.71-3.69 (s, 2H), 2.34 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-3-fluoro-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ9.34 (s, 1H), 9.03 (s, 1H), 8.05-8.01 (d, 1H), 7.62-7.59 (d, 1H), 7.50-7.46 (d, 2H), 7.31-7.23 (m, 3H), 4.18 (s, 2H), 3.97-3.95 (s, 2H), 3.83-3.82 (d, 4H), 3.75-3.74 (d, 4H), 3.71-3.69 (d, 2H), 2.36-2.34 (s, 3H)


4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzenesulfonamide



151


¹H NMR (500 MHz, DMSO-d₆): δ9.14 (s, 2H), 8.23-8.21 (s, 2H), 7.75-7.73 (s, 2H), 7.63-7.62 (d, 2H), 7.61-7.59 (d, 2H), 7.22 (s, 2H), 3.85 (d, 4H), 3.77-3.76 (d, 4H), 2.36 (s, 3H)

N-(4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-acetamide




¹H NMR (500 MHz, DMSO-d₆): δ9.83 (s, 1H), 9.03 (s, 1H), 8.72 (s, 1H), 8.20-8.19 (d, 2H), 7.59-7.57 (d, 2H), 7.49-7.48 (d, 2H), 7.38-7.36 (d, 2H), 3.84-3.83 (d, 4H), 3.77-3.76 (d, 4H), 2.36 (s, 3H), 2.01 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzoxazol-5-yl)-urea



¹H NMR (500 MHz, DMSO-d₆): δ9.16 (s, 1H), 8.95 (s, 1H), 8.21-8.19 (d, 2H), 7.60-7.58 (d, 2H), 7.43 (s, 1H), 7.14-7.12 (s, 1H), 6.92-6.90 (m, 1H), 3.85-3.83 (d, 4H), 3.77-3.76 (d, 4H), 2.36-2.34 (s, 3H)


1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-isoxazol-3-yl-urea



152


¹H NMR (500 MHz, DMSO-d₆): δ9.74 (s, 1H), 9.22 (s, 1H), 8.76-8.76 (d, 1H), 8.23-8.21 (d, 2H), 7.63-7.59 (d, 2H), 6.87-6.87 (d, 1H), 3.84-3.83 (d, 4H), 3.77-3.76 (s, 4H), 2.31 (s, 3H)

5 1-(6-Chloro-5-methylsulfanyl-4-morpholin-4-yl-[2, 5']bipyrimidinyl-2'-yl)-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ11.57 (s, 1H), 10.24 (b, 1H), 9.39 (d, 2H), 7.70-7.68 (d, 2H), 7.37-7.35 (d, 2H), 4.19-4.18 (d, 2H), 3.98-3.90 (d, 2H), 3.90 (s, 4H), 3.77-3.71 (m, 6H), 2.35 (s, 3H)

25 4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-3-fluoro-phenyl]-ureido}-benzamide



¹H NMR (500 MHz, DMSO-d₆): δ9.35 (s, 1H), 9.16 (s, 1H), 8.04-8.02 (d, 1H), 7.84-7.82 (d, 3H), 7.62-7.52 (m, 3H), 7.26-7.24 (m, 2H), 3.84-3.82 (d, 4H), 3.75-3.73 (d, 4H), 2.35 (s, 3H)


50 4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenyl]-ureido}-benzamide



153


¹H NMR (500 MHz, DMSO-d₆): δ9.45 (s, 1H), 8.97 (s, 1H), 8.37-8.33 (d, 1H), 8.09-8.01 (dd, 2H), 7.85-7.83 (d, 3H), 7.54-7.52 (d, 2H), 7.21 (s, 1H), 3.86-3.85 (d, 4H), 3.77-3.75 (d, 4H), 2.34 (s, 3H)

4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenyl]-ureido}-benzenesulfonamide




¹H NMR (500 MHz, DMSO-d₆): δ9.62 (s, 1H), 9.01 (s, 1H), 8.33-8.32 (d, 1H), 8.09-8.02 (dd, 2H), 7.76-7.75 (d, 2H), 7.64-7.63 (d, 1H), 7.23 (s, 1H), 3.86-3.85 (d, 4H), 3.77-3.75 (d, 4H), 2.32 (s, 3H)

4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-3-fluoro-phenyl]-ureido}-benzenesulfonamide



¹H NMR (500 MHz, DMSO-d₆): δ9.37 (s, 1H), 9.28 (s, 1H), 8.05-8.02 (d, 1H), 7.75-7.73 (d, 2H), 7.63-7.58 (m, 3H), 7.26-7.22 (m, 3H), 3.82-3.81 (d, 4H), 3.73-3.72 (d, 4H), 2.31 (s, 3H)


N-(5-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-pyridin-2-yl)-acetamide



154


¹H NMR (500 MHz, DMSO-d₆): δ10.38 (s, 1H), 9.20 (s, 1H), 8.90 (s, 1H), 8.43-8.42 (d, 1H), 8.21-8.20 (d, 2H), 7.91-7.89 (d, 1H), 7.84-7.82 (dd, 1H), 7.60-7.59 (d, 2H), 3.84-3.83 (d, 4H), 3.77-3.75 (d, 4H), 2.34 (s, 3H), 2.08-2.06 (d, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-methanesulfonyl-piperazin-1-yl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ9.01 (s, 1H), 8.61 (s, 1H), 8.20-8.18 (d, 2H), 7.58-7.56 (d, 2H), 7.35-7.33 (d, 2H), 6.94-6.93 (d, 2H), 3.85-3.83 (d, 4H), 3.77-3.75 (d, 4H), 3.25-3.22 (d, 4H), 3.17-3.15 (d, 4H), 2.92 (s, 3H), 2.34 (s, 3H)

4-[3-(6-Chloro-5-methylsulfanyl-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-yl)-ureido]-benzenesulfonamide



¹H NMR (500 MHz, DMSO-d₆): δ11.85 (s, 1H), 10.75 (s, 1H), 9.41 (s, 2H), 7.87-7.85 (d, 2H), 7.80-7.78 (d, 2H), 7.27 (s, 2H), 3.91-3.90 (d, 4H), 3.77-3.75 (d, 4H), 2.35 (s, 3H)


1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzooxazol-6-yl)-urea



155


¹H NMR (500 MHz, DMSO-d₆): δ 11.49 (s, 1H), 9.07 (s, 1H), 8.87 (s, 1H), 8.21-8.20 (d, 2H), 7.59-7.58 (d, 3H), 7.09-7.07 (d, 1H), 7.02-7.00 (d, 1H), 3.84 (d, 4H), 3.76 (d, 4H), 2.34 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-sulfonyl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ 9.36 (s, 1H), 9.23 (s, 1H), 8.23-8.22 (d, 2H), 7.74-7.72 (d, 2H), 7.67-7.65 (d, 2H), 7.62-7.61 (d, 2H), 3.84 (d, 4H), 3.76 (d, 4H), 3.63 (s, 4H), 2.84 (s, 4H), 2.34 (s, 3H)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-dimethylamino-piperidine-1-carbonyl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.24 (s, 1H), 9.14 (s, 1H), 8.22-8.20 (d, 2H), 7.61-7.59 (d, 2H), 7.54-7.52 (d, 2H), 7.36-7.34 (d, 2H), 3.84-3.83 (d, 4H), 3.77-3.75 (d, 4H), 2.72 (b, 4H), 2.38 (s, 6H), 2.36-2.34 (m, 4H), 1.84 (b, 2H), 1.43-1.41 (b, 2H)

Example 5

Preparation of Compounds of Formula (I) in Scheme 3

5-(Methylthio)pyrimidine-2,4-diol




5-iodouracil (476 mg, 2 mmole) was placed in a dry and argon-flushed Schlenk-tube equipped with a magnetic stirring bar and a septum. With vigorous stirring, the substrate

156

was dried for 15 min in high vacuum to exclude the presence of water in the hygroscopic substrate. Then, a solution of LiCl in THF (0.5M; 8 ml; 4 mmole, 2.0 eq) was added and after stirring for some minutes at r.t., the substrate dissolved to give a clear and colourless solution. The solution was cooled to -20° C. and MeMgCl (3M in THF; 1.33 ml, 4 mmole, 2 eq) was added dropwise. After completion of the addition, the resulting clear solution was stirred at -20° C. for 20 min. Afterward, i-PrMgCl (1.32 M in THF; 1.82 ml, 2.40 mmole, 1.20 eq) was added slowly and the resulting mixture was allowed to warm up to room temperature. After one hour, a thick, grayish slurry had formed and the mixture was cooled to -20° C. and S-methyl methanesulfonylthioate (328 mg, 2.6 mmole, 1.3 eq) was obtained. Afterwards, the mixture was warmed up to r.t. and stirred at that temperature until TLC indicated completion of the reaction. After quenching with MeOH, the mixture was transferred to a separation funnel containing water (40 ml). By careful addition of 2.0 M HCl, the pH of the mixture was adjusted to ca. 5-6. The aqueous layer was extracted with EA, the combined organic layers were dried (Na₂SO₄) and evaporated. Recrystallization from MeOH afforded the product as a colorless, crystalline solid which was dried in high vacuum (64%).

¹H NMR (500 MHz, DMSO-d₆): δ 7.41 (s, 1H), 2.49 (s, 3H)


2,4-Dichloro-5-(methylthio)pyrimidine



5-(methylthio)pyrimidine-2,4-diol (0.5 g, 3.16 mmole), POCl₃ (3 ml) and dimethylaniline (0.5 ml) were heated to reflux for 12 hrs. The mixture was cooled to 0° C. and carefully poured over ice. The white solid formed was separated by filtration and dried under vacuum to afford the product 2,4-dichloro-5-(methylthio)pyrimidine (0.49 g, 80%).

¹H NMR (500 MHz, CDCl₃-d₁): δ 8.29 (s, 1H), 2.55 (s, 3H)

4-(2-Chloro-5-(methylthio)pyrimidin-4-yl)morpholine




A solution of the 2,4-dichloro-5-(methylthio)pyrimidine (1.16 g, 6 mmole) in acetone (10 ml) at 0° C. containing Na₂CO₃ (1.1 eq, 6.6 mmole, 0.7 g) was treated dropwise with a solution of morpholine (1.0 eq, 6.6 mmole, 0.6 ml) in acetone (2 ml) and stirred for 1 hr at 0° C. TLC (EA:Hex=1:2, R_f=0.6) showed completion of the reaction to give two prod-

157

ucts. The acetone was removed in vacuo, the residue was partitioned between water and EA, and the organic layer was dried (MgSO₄) and evaporated in vacuo. Column chromatography (EA/Hex=1/6) gave the product (700 mg, 47%).

¹H NMR (500 MHz, CDCl₃-d₁): δ 8.13 (s, 1H), 3.79~3.82 (m, 8H), 2.39 (s, 3H)


4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)benzenamine



A mixture of 4-(2-chloro-5-(methylthio)pyrimidin-4-yl) morpholine (0.3 g, 1.22 mmole), 4-aminophenylboronic acid pinacol ester (0.58 g, 2.68 mmole), sodium bicarbonate (0.28 g, 2.68 mmole), Pd(PPh₃)₄ (0.14 g, 0.122 mmole), dioxane (15 ml) and H₂O (3.5 ml) was heated to reflux. After 12 hrs, the reaction mixture was cooled, diluted with EA, washed with brine, and dried (MgSO₄), and the solvent was removed in vacuo. Purification on silica gel (EA/Hex=1/2) yielded the desired compound (180 mg, 49%).

¹H NMR (500 MHz, DMSO-d₆): δ 8.33 (s, 1H), 8.01~8.03 (d, 2H), 6.58~6.60 (d, 2H), 5.59 (s, 2H), 3.73~3.74 (d, 4H), 3.64~3.66 (d, 4H), 2.42 (s, 3H)


3-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenol



¹H NMR (500 MHz, CDCl₃-d₁): δ 8.37 (s, 1H), 7.92~7.94 (d, 2H), 7.85 (s, 1H), 7.30~7.33 (m, 1H), 6.93~6.95 (m, 1H), 3.85~3.87 (m, 4H), 3.79~3.80 (m, 4H), 2.44 (s, 3H)


158

4-(2-(4-(Methylsulfonyl)phenyl)-5-(methylthio)pyrimidin-4-yl)morpholine



¹H NMR (500 MHz, DMSO-d₆): δ 8.46~8.48 (m, 3H), 7.91~7.93 (m, 2H), 7.42 (br, 2H), 3.75~3.77 (m, 4H), 3.68~3.69 (m, 4H), 2.49 (s, 3H)


1-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)-3-phenylurea



A mixture of 4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)benzeneamine (0.1 g, 0.33 mmole), co-solvent (5 ml, DMF/THF=6/40) and phenyl isocyanate (0.2 ml, 0.92 mmole) was heated to 80° C. After 12 hrs, the reaction mixture was cooled, and the THF removed in vacuo. Then the mixture was diluted with EA, washed with brine, and dried (MgSO₄). The solvent was removed in vacuo. Purification of the residue on silica gel (EA/Hex=1/2) yielded the desired compound (42.2 mg, 30.4%).

¹H NMR (500 MHz, DMSO-d₆): δ 9.01 (s, 1H), 8.82 (s, 1H), 8.41 (s, 1H), 8.24~8.26 (d, 2H), 7.56~7.57 (d, 2H), 7.46~7.47 (d, 2H), 7.27~7.30 (m, 2H), 6.98 (m, 1H), 3.74 (s, 4H), 3.68 (s, 4H), 2.47 (s, 3H)


1-Ethyl-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea



159


¹H NMR (500 MHz, DMSO-d₆): δ 8.71 (s, 1H), 8.39 (s, 1H), 8.17~8.19 (d, 2H), 7.47~7.49 (d, 2H), 6.20~6.22 (m, 2H), 3.73~3.75 (m, 4H), 3.65~3.67 (m, 4H), 3.10~3.12 (m, 2H), 2.36 (s, 3H), 1.04~1.06 (m, 3H)

1-(3,4-Difluorophenyl)-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.05 (s, 1H), 9.01 (s, 1H), 8.41 (s, 1H), 8.24~8.26 (d, 2H), 7.67~7.70 (m, 1H), 7.57~7.66 (d, 2H), 7.34~7.36 (m, 1H), 7.13~7.15 (br, 1H), 3.74 (s, 4H), 3.68 (s, 4H), 2.48 (s, 3H)

1-(4-Fluorophenyl)-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea



¹H NMR (500 MHz, DMSO-d₆): δ 8.41 (s, 3H), 8.21~8.23 (d, 2H), 7.57~7.59 (d, 2H), 7.48~7.50 (m, 2H), 7.08~7.12 (m, 2H), 3.75 (s, 4H), 3.67 (s, 4H), 2.47 (s, 3H).

Phenyl 4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenylcarbamate




160

The phenyl chloroformate (0.84 ml, 6.68 mmole, 2.5 eq) was added to a mixture of 4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)benzeneamine (0.84 g, 2.18 mmole, 1 eq), NaHCO₃(aq) (1M, 6.3 ml, 6.3 mmole 2.2 eq) and ACN (13 ml) at 0°C. The reaction mixture was stirred for 8 hrs and then dried. The reaction mixture was partitioned between EA and water, the organic layer was washed with brine, dried over MgSO₄, filtered and evaporated in vacuo. The resulting residue was purified by flash chromatography on silica gel (Hexane/EtOAc 2:1) to give a yellow solid (80%).


¹H NMR (500 MHz, CDCl₃-d₁): δ 8.38 (s, 1H), 8.35~8.36 (d, 2H), 7.53~7.55 (d, 2H), 7.39~7.42 (m, 2H), 7.1~7.2 (m, 3H), 3.79~3.87 (d, 8H), 2.44 (s, 3H)

1-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)-3-morpholinourea



¹H NMR (500 MHz, DMSO-d₆): δ 8.78 (s, 1H), 8.41 (s, 1H), 8.20~8.22 (d, 2H), 7.91 (s, 1H), 7.67~7.69 (d, 2H), 3.66~3.76 (m, 13H), 2.49 (s, 3H)


1-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)-3-(pyridin-3-yl)urea



¹H NMR (500 MHz, CD₃OD-d₄): δ 8.63 (s, 1H), 8.35 (s, 1H), 8.26~8.28 (d, 2H), 8.19~8.20 (d, 1H), 8.03~8.05 (d, 1H), 7.56~7.57 (d, 2H), 7.37~7.40 (m, 1H), 3.85 (s, 8H), 2.46 (s, 3H)


161

1-(3-Fluoro-4-morpholinophenyl)-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.07 (s, 1H), 8.91 (s, 1H), 8.41 (s, 1H), 8.23~8.25 (d, 2H), 7.55~7.56 (d, 2H), 7.43~7.44 (d, 1H), 7.07 (d, 1H), 6.98~7.0 (t, 1H), 3.67~3.76 (m, 12H), 2.93~2.94 (s, 4H), 2.48 (s, 3H)

1-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)-3-(4-(3-oxomorpholino)phenyl)urea



¹H NMR (500 MHz, DMSO-d₆): δ 9.09 (s, 1H), 8.96 (s, 1H), 8.42 (s, 1H), 8.24~8.26 (d, 2H), 7.56~7.58 (d, 2H), 7.48~7.50 (d, 2H), 7.28~7.30 (d, 2H), 4.18 (s, 2H), 3.95~3.97 (t, 2H), 3.68~3.76 (m, 8H), 2.48 (s, 3H)

1-(4-(4-Methylpiperazin-1-yl)phenyl)-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea




¹H NMR (500 MHz, DMSO-d₆): δ 8.94 (s, 1H), 8.58 (s, 1H), 8.41 (s, 1H), 8.22~8.24 (d, 2H), 7.54~7.56 (d, 2H),

162


7.29~7.31 (d, 2H), 6.86~6.88 (d, 2H), 3.74~3.76 (d, 4H), 3.67~3.68 (d, 4H), 3.03~3.04 (s, 4H), 2.43~2.44 (s, 3H), 2.36 (s, 4H), 2.21 (s, 3H)

1-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)-3-(4-thiomorpholinophenyl)urea




¹H NMR (500 MHz, DMSO-d₆): δ 8.99 (s, 1H), 8.65 (s, 1H), 8.41 (s, 1H), 8.22~8.23 (d, 2H), 7.55~7.57 (d, 2H), 7.31~7.33 (d, 2H), 6.86~6.88 (d, 2H), 3.75~3.76 (d, 4H), 3.67~3.38 (d, 4H), 3.37~3.39 (d, 4H), 2.67~2.68 (d, 4H), 2.47 (s, 3H)

1-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)-3-(4-morpholinophenyl)urea



¹H NMR (500 MHz, DMSO-d₆): δ 9.01 (s, 1H), 8.67 (s, 1H), 8.38 (s, 1H), 8.22~8.24 (d, 2H), 7.54~7.56 (d, 2H), 7.28~7.34 (d, 2H), 6.85~6.89 (d, 2H), 3.68~3.76 (m, 8H), 3.67~3.68 (d, 4H), 3.02~3.03 (d, 4H), 2.49 (s, 3H)


1-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)-3-(thiazol-2-yl)urea



163


¹H NMR (500 MHz, DMSO-d₆): δ 8.42 (s, 1H), 8.27~8.28 (d, 2H), 7.59~7.67 (d, 2H), 7.38 (s, 1H), 6.85~6.89 (s, 1H), 3.76 (s, 4H), 3.68 (s, 4H), 2.49 (s, 3H)

1-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)-3-(3-(trifluoromethyl)phenyl)urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.27 (s, 1H), 9.20 (s, 1H), 8.42 (s, 1H), 8.25~8.26 (d, 2H), 8.03 (s, 1H), 7.58~7.59 (d, 3H), 7.50~7.53 (t, 1H), 7.31~7.32 (d, 1H), 3.75 (s, 4H), 3.68 (s, 4H), 2.48 (s, 3H)

Ethyl 4-(3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)ureido)benzoate



¹H NMR (500 MHz, DMSO-d₆): δ 9.29 (s, 1H), 9.18 (s, 1H), 8.42 (s, 1H), 8.25~8.27 (d, 2H), 7.88~7.90 (d, 2H), 7.57~7.61 (t, 4H), 4.25~4.30 (q, 2H), 3.75~3.76 (d, 4H), 3.67~3.69 (d, 4H), 2.48 (s, 3H), 1.25~1.32 (t, 3H)

1-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)-3-(6-morpholinopyridin-3-yl)urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.01 (s, 1H), 8.67 (s, 1H), 8.41 (s, 1H), 8.19~8.22 (m, 3H), 7.77 (m, 1H),

164


7.55~7.56 (d, 2H), 6.85~6.89 (d, 1H), 3.75~3.76 (m, 4H), 3.67~3.70 (m, 4H), 2.48 (s, 3H)

1-(6-(4-Methylpiperazin-1-yl)pyridin-3-yl)-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.01 (s, 1H), 8.67 (s, 1H), 8.41 (s, 1H), 8.22~8.24 (d, 2H), 8.17 (s, 1H), 7.7 (d, 2H), 7.54~7.56 (d, 2H), 6.85~6.89 (d, 1H), 3.74~3.76 (d, 4H), 3.67~3.68 (d, 4H), 3.37~3.38 (d, 4H), 2.47 (s, 3H), 2.38~2.40 (d, 4H), 2.20 (s, 3H)

4-(3-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)ureido)benzamide



¹H NMR (500 MHz, DMSO-d₆): δ 9.10 (s, 1H), 9.08 (s, 1H), 8.42 (s, 1H), 8.25~8.27 (d, 2H), 7.21~7.83 (m, 3H), 7.57~7.59 (d, 2H), 7.51~7.53 (d, 2H), 7.19 (s, 1H), 3.75~3.76 (d, 4H), 3.67~3.69 (d, 4H), 2.47 (s, 3H)


1-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea



165

¹H NMR (500 MHz, DMSO-d₆): δ 8.81 (s, 1H), 8.40 (s, 1H), 8.18~8.20 (d, 2H), 7.49~7.50 (d, 2H), 5.96 (d, 2H), 3.74 (s, 4H), 3.67 (s, 4H), 2.46 (s, 3H)


1-(2-Aminophenyl)-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)thiourea



To a stirred solution of 1,1'-thiocarbonyldiimidazole (0.35 g, 2 mmole), imidazole (27 mg, 0.4 mmole), and CH₃CN (15 ml) at 0° C. was added 4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)benzenamine (0.4 g, 1.32 mmole) dissolved in CH₃CN (30 ml) dropwise over 10 minutes. After 10 minutes, the cooling bath was removed. After 3 hours, 1,2-phenylenediamine (0.285 g, 2.64 mmole) was added to the reaction mixture and the reaction mixture was heated to 50° C. for 3 hours and then stirred at ambient temperature for 16 hours. The solvent was removed by evaporation. The residue was chromatographed to give thiourea (10%) as a white solid.

¹H NMR (500 MHz, DMSO-d₆): δ 9.85 (s, 1H), 9.20 (s, 1H), 8.43 (s, 1H), 8.25~8.27 (d, 2H), 7.68~7.70 (d, 2H), 7.09~7.10 (d, 1H), 6.95~6.98 (t, 1H), 6.74~6.76 (d, 1H), 6.55~6.58 (t, 1H), 4.94 (br, 2H), 3.74~3.76 (d, 4H), 3.67~3.69 (d, 4H), 2.49 (s, 3H)

N-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)-1H-benzod[*d*]imidazol-2-amine




A suspension of 1-(2-aminophenyl)-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)thiourea (60 mg, 0.132 mmole) and dicyclohexylcarbodiimide (31 mg, 0.15 mmole) in THF (5 ml) was heated at 64° C. for 20 hr. The cooled mixture was purified by flash chromatography to give a product (40 mg, 73%).

¹H NMR (500 MHz, DMSO-d₆): δ 11.13 (s, 1H), 9.85 (s, 1H), 8.42 (s, 1H), 8.28~8.30 (d, 2H), 7.86~7.88 (d, 2H),

166


7.31~7.37 (br, 2H), 7.01 (br, 1H), 3.75~3.76 (d, 4H), 3.69~3.70 (d, 4H), 2.47 (s, 3H)

4-(3-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)ureido)benzenesulfonamide




¹H NMR (500 MHz, DMSO-d₆): δ 8.42 (s, 1H), 8.25~8.27 (d, 2H), 7.72~7.74 (d, 2H), 7.63~7.65 (d, 2H), 7.59~7.60 (d, 2H), 7.20 (s, 2H), 3.75~3.76 (d, 4H), 3.68~3.69 (d, 4H)

1-(3,4-Dimethoxyphenyl)-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea



¹H NMR (500 MHz, DMSO-d₆): δ 8.98 (s, 1H), 8.71 (s, 1H), 8.41 (s, 1H), 8.23~8.24 (d, 2H), 7.55~7.57 (d, 2H), 7.21 (s, 1H), 6.87~6.88 (t, 2H), 3.67~3.76 (m, 14H), 2.47 (s, 3H)


4-(3-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)ureido)benzoic acid



167


¹H NMR (500 MHz, DMSO-d₆): δ 9.47 (br, 2H), 8.35 (s, 1H), 8.24~8.25 (d, 2H), 7.87~7.88 (d, 2H), 7.63~7.64 (d, 2H), 7.57~7.59 (d, 2H), 3.87 (s, 5H), 3.78 (s, 6H),

1-(Benzo[d][1,3]dioxol-5-yl)-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.05 (s, 1H), 8.81 (s, 1H), 8.41 (s, 1H), 8.22~8.24 (d, 2H), 7.54~7.56 (d, 2H), 7.21 (s, 1H), 6.82~6.84 (d, 1H), 6.77~6.79 (d, 1H), 5.97 (s, 2H), 3.74~3.75 (d, 4H), 3.67~3.68 (d, 4H), 2.47 (s, 3H)

1-(5-(4-Methylpiperazin-1-yl)pyridin-2-yl)-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea




¹H NMR (500 MHz, DMSO-d₆): δ 10.57 (br, 1H), 9.29 (s, 1H), 8.42 (s, 1H), 8.25~8.27 (d, 2H), 7.95 (s, 1H), 7.61~7.62 (d, 2H), 7.45~7.46 (t, 2H), 3.75~3.76 (d, 4H), 3.67~3.68 (d, 4H), 3.56 (d, 4H), 3.08 (s, 4H), 2.47 (s, 3H), 2.21 (s, 3H)

1-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)-3-(4-(3-oxomorpholino)phenyl)thiourea

**168**


¹H NMR (500 MHz, DMSO-d₆): δ 8.44 (s, 1H), 8.26~8.28 (d, 2H), 7.64~7.66 (d, 2H), 7.50~7.54 (t, 3H), 7.33~7.35 (t, 2H), 4.19 (s, 2H), 3.97~3.98 (t, 2H), 3.67~3.76 (m, 7H), 2.51 (s, 3H)

1-(6-Bromopyridin-3-yl)-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.20~9.25 (d, 2H), 8.48 (s, 1H), 8.41 (s, 1H), 8.25~8.26 (d, 2H), 7.90~7.92 (d, 1H), 7.54~7.58 (t, 3H), 3.68~3.75 (d, 8H), 2.47 (s, 3H)

1-Isopropyl-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea



¹H NMR (500 MHz, DMSO-d₆): δ 8.61 (s, 1H), 8.39 (s, 1H), 8.17~8.19 (d, 2H), 7.46~7.48 (d, 2H), 6.14~6.15 (d, 1H), 3.75~3.78 (d, 5H), 3.66~3.67 (d, 4H), 2.46 (s, 3H), 1.09~1.10 (d, 6H)


1-Tert-butyl-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea



169


¹H NMR (500 MHz, DMSO-d₆): δ 8.56 (br, 1H), 8.39 (s, 1H), 8.17~8.18 (d, 2H), 7.44~7.46 (d, 2H), 6.14 (br, 1H), 3.74~3.76 (d, 4H), 3.66~3.67 (d, 4H), 2.46 (s, 3H), 1.29 (s, 9H)

1-(4-(2-(Dimethylamino)ethoxy)phenyl)-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea




¹H NMR (500 MHz, DMSO-d₆): δ 8.97 (s, 1H), 8.62 (s, 1H), 8.41 (s, 1H), 8.22~8.24 (d, 2H), 7.54~7.56 (d, 2H), 7.34~7.36 (d, 2H), 6.86~6.88 (d, 2H), 3.98~4.00 (t, 2H), 3.74~3.76 (d, 4H), 3.67~3.68 (d, 4H), 2.59~2.60 (d, 2H), 2.49 (s, 3H), 2.20 (s, 6H)

1,3-Bis(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea



¹H NMR (500 MHz, DMSO-d₆): δ 9.24 (s, 1H), 8.42 (s, 1H), 8.25~8.27 (d, 2H), 7.58~7.60 (d, 2H), 3.75~3.76 (d, 4H), 3.68~3.69 (d, 4H)


3-Fluoro-4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)benzenamine



170


¹H NMR (500 MHz, CDCl₃-d₁): δ 8.38 (s, 1H), 7.94~7.98 (t, 2H), 7.53~7.55 (d, 2H), 7.39~7.42 (m, 2H), 7.1~7.2 (m, 3H), 3.79~3.87 (d, 8H), 2.44 (s, 3H)

N-(4-(3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)ureido)phenyl)methanesulfonamide



¹H NMR (500 MHz, DMSO-d₆): δ 9.45 (s, 1H), 9.16 (s, 1H), 8.98 (s, 1H), 8.41 (s, 1H), 8.23~8.25 (d, 2H), 7.56~7.57 (d, 2H), 7.42~7.44 (d, 2H), 7.13~7.15 (d, 2H), 3.74~3.75 (d, 4H), 3.67~3.68 (d, 4H), 2.91 (s, 3H), 2.48 (s, 3H)

Phenyl 4-(5-(methylsulfinyl)-4-morpholinopyrimidin-2-yl)phenylcarbamate




The H₂O₂ (0.32 ml, 3.13 mmole) solution was added to a stirred solution of phenyl 4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenylcarbamate (0.54 g, 1.28 mmole), CH₂Cl₂ (7 ml), Ac₂O (0.13 ml), and silica gel (230-400 mesh, 200 mg). The solution was stirred at room temperature for 2-3 hrs, and quenched with NaHSO_{3(aq)} (70 ml), and NaHCO_{3(aq)} (70 ml), the solvent was removed, the residue was chromatographed to give the product (71%) as a white solid.

¹H NMR (500 MHz, CDCl₃-d₁): δ 8.97 (s, 1H), 8.43 (s, 1H), 8.41 (s, 1H), 7.56~7.58 (t, 2H), 7.39~7.42 (t, 2H), 7.17~7.27 (m, 3H), 3.81~3.87 (m, 6H), 3.48~3.68 (m, 2H), 2.79 (s, 3H)


171

1-(4-(5-(Methylsulfinyl)-4-morpholinopyrimidin-2-yl)phenyl)-3-(4-(3-oxomorpholino)phenyl)urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.09 (s, 1H), 8.92 (s, 1H), 8.76 (s, 1H), 8.31~8.32 (d, 2H), 7.59~7.61 (d, 2H), 7.48~7.50 (d, 2H), 7.28~7.30 (d, 2H), 4.18 (s, 2H), 3.96 (d, 4H), 3.62~3.73 (m, 10H), 2.79 (s, 3H)

1-(4-(Methylsulfonyl)phenyl)-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea



¹H NMR (500 MHz, DMSO-d₆): δ 9.61 (br, 1H), 9.33 (br, 1H), 8.42 (s, 1H), 8.25~8.27 (d, 2H), 7.81~7.83 (d, 2H), 7.71~7.72 (d, 2H), 7.59~7.61 (d, 2H), 3.75~3.76 (d, 4H), 3.68~3.69 (d, 4H), 3.15 (s, 3H)

1-(4-(4-(Methylsulfonyl)piperazin-1-yl)phenyl)-3-(4-(5-(methylthio)-4-morpholinopyrimidin-2-yl)phenyl)urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.38 (br, 1H), 9.02 (br, 1H), 8.41 (s, 1H), 8.21~8.23 (d, 2H), 7.56~7.58 (d, 2H),

172


7.35~7.37 (d, 2H), 6.91~6.93 (d, 2H), 3.74~3.76 (d, 4H), 3.56~3.68 (d, 4H), 3.24~3.25 (d, 4H), 3.15~3.16 (d, 4H), 2.92 (s, 3H), 2.47 (s, 3H)

5-(Methylthio)-4-morpholino-2,5'-bipyrimidin-2'-amine




¹H NMR (500 MHz, DMSO-d₆): δ 9.03 (s, 2H), 8.36 (s, 1H), 7.14 (s, 2H), 3.72~3.73 (d, 4H), 3.67~3.68 (d, 4H), 2.49 (s, 3H)

1-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)-3-(2-oxo-2,3-dihydrobenzo[d]oxazol-6-yl)urea



¹H NMR (500 MHz, DMSO-d₆): δ 9.00 (s, 1H), 8.58 (s, 1H), 8.41 (s, 1H), 8.23~8.25 (d, 2H), 7.55~7.57 (d, 3H), 7.05~7.07 (d, 1H), 6.97~6.99 (d, 1H), 3.74~3.76 (d, 4H), 3.67~3.69 (d, 4H), 2.48 (s, 3H)


1-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)-3-(4-(morpholine-4-carbonyl)phenyl)urea



¹H NMR (500 MHz, DMSO-d₆): δ 8.41 (s, 1H), 8.20~8.22 (d, 2H), 7.60~7.66 (m, 4H), 7.31~7.33 (d, 2H), 3.74~3.75 (d, 4H), 3.68 (s, 4H), 3.59 (s, 4H), 3.50 (s, 4H), 2.48 (s, 3H)

173

1-(4-(5-(Methylthio)-4-morpholinopyrimidin-2-yl)phenyl)-3-(pyridin-4-yl)urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.35 (s, 1H), 9.28 (s, 1H), 8.42 (s, 1H), 8.36~8.37 (d, 2H), 8.26~8.27 (d, 2H), 7.57~7.58 (d, 2H), 7.44~7.45 (d, 2H), 3.75~3.76 (d, 4H), 3.67~3.68 (d, 4H), 2.48 (s, 3H)

Example 6

Preparation of Compounds of Formula (I) in Scheme 6


4-Chloro-2,6-dimethoxy-5-methylsulfanyl-pyrimidine



A mixture of 2,4,6-trichloro-5-methylsulfanyl-pyrimidine (5.00 g, 21.73 mmol.) was added dropwise to a slurry of sodium methoxide (2.47 g, 45.65 mmol, 2.1 eq) in MeOH (80 ml) in ice bath at 0° C., then the reaction mixture was stirred at r.t. for 2 hrs and dried, and the solvent was evaporated. The reaction mixture was partitioned between EA and 0.1 N HCl. The organic layer was washed with brine, dried over MgSO₄, filtered and evaporated in vacuo. The resulting residue was purified by flash chromatography on silica gel (Hexane/EtOAc 9:1) to give a gray solid 3.36 g (70.2%).

¹H NMR (500 MHz, CDCl₃-d₁): δ 4.07 (s, 3H), 4.00 (s, 3H), 2.33 (s, 3H)

2-(2,6-Dimethoxy-5-methylsulfanyl-pyrimidin-4-yl)-malonic acid dimethyl ester




174

A mixture of 4-chloro-2,6-dimethoxy-5-methylsulfanyl-pyrimidine (3.30 g, 15.0 mmol.) and dimethyl malonate (11.88 g, 90.0 mmol, 6.0 eq.) was added dropwise to slurry of sodium hydride (60% w/w) (3.67 g, 90.0 mmol, 6.0 eq) in DMF (80 ml) in ice bath at 0° C., then the reaction mixture was stirred at 100° C. for 10 hrs and dried, and the solvent was evaporated. The reaction mixture was partitioned between EA and 0.1 N HCl. The organic layer was washed with brine, dried over MgSO₄, filtered and evaporated in vacuo. The resulting residue was purified by flash chromatography on silica gel (Hexane/EtOAc 6:1) to give a pale yellow oil product of 2.85 g (60.2%).


¹H NMR (500 MHz, CDCl₃-d₁): δ 5.40 (s, 1H), 4.07 (s, 3H), 3.95 (s, 3H), 3.80 (s, 6H), 2.23 (s, 3H)

(2,6-Dihydroxy-5-methylsulfanyl-pyrimidin-4-yl)-acetic acid methyl ester



A mixture of 2-(2,6-dimethoxy-5-methylsulfanyl-pyrimidin-4-yl)-malonic acid dimethyl ester (2.85 g, 9.0 mmol.) was added dropwise to slurry of HCl (37% w/w) (1.0 ml) in MeOH (50 ml). The reaction mixture was stirred at 100° C. for 24 hr and dried, and the solvent was evaporated in vacuo to give a crude as a pale yellow solid of 1.85 g (89.2%). ¹H NMR (500 MHz, DMSO-d₆): δ 11.35 (s, 1H), 11.23 (s, 1H), 3.81 (s, 2H), 3.66 (s, 3H), 2.11 (s, 3H)

(2,6-Dichloro-5-methylsulfanyl-pyrimidin-4-yl)-acetic acid methyl ester




A mixture of (2,6-dihydroxy-5-methylsulfanyl-pyrimidin-4-yl)-acetic acid methyl ester (1.85 g, 9.0 mmol.) was added dropwise to a slurry of POCl₃ (8.0 ml). After the reaction mixture was stirred at 90° C. for 4 hrs, hydrolysis was carried out in iced salt water. The product of the reaction was extracted with EA and then evaporated to remove the EA so as to give a crude. Purification of the crude by flash chromatography on silica gel (Hexane/EtOAc 3:1) gave a brown oil product of 866 mg (40.5%).

¹H NMR (500 MHz, CDCl₃-d₁): δ 4.15 (s, 2H), 3.74 (s, 3H), 2.39 (s, 3H)

175


(2-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-4-yl)-acetic acid methyl ester



A mixture of (2,6-dichloro-5-methylsulfanyl-pyrimidin-4-yl)-acetic acid methyl ester (860 mg, 3.22 mmol.), morpholine (336 mg, 3.86 mmol, 1.2 eq.) and THF (10 ml) was stirred at room temperature for 2.0 hrs. The reaction mixture was dried and the solvent was evaporated. The reaction mixture was subjected to addition of 0.1 N HCl washing free morpholine, extraction with EA and evaporation to remove the EA so as to give a crude. Purification of the crude by flash chromatography on silica gel (Hexane/EtOAc 2:1) gave a gray solid of 427 mg (42.7%).

¹H NMR (500 MHz, CDCl₃-d₁): δ 3.99 (s, 2H), 3.80 (dd, 4H), 3.73 (dd, 4H), 3.71 (s, 3H), 2.21 (s, 3H)

[2-(4-Amino-phenyl)-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-4-yl]-acetic acid methyl ester




A mixture of (2-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-4-yl)-acetic acid methyl ester (420 mg, 1.47 mmol.), 4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-aniline (357 mg, 1.62 mmol.), (PPh₃)₄Pd (170 mg, 0.07 mmol.), 2M Na₂CO₃ (2.2 ml, 4.14 mmol, 3.0 eq.) and dioxane (10 ml) was added a reaction vessel which was flushed with argon. The reaction mixture was stirred at 90° C. for 12 hr then dried, and the solvent was evaporated. The reaction mixture was partitioned between EA and water, the organic layer was washed with brine, dried over MgSO₄, filtered and evaporated in vacuo. The resulting residue was purified by flash chromatography on silica gel (Hexane/EtOAc 3:2) to give a pale yellow solid of 424 mg (43.8%).

¹H NMR (500 MHz, DMSO-d₆): δ 7.75 (dd, 2H), 6.70 (dd, 2H), 4.05 (s, 2H), 3.84 (s, 3H), 3.74 (d, 8H), 1.91 (s, 3H)

176


{5-Methylsulfanyl-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidin-4-yl}-acetic acid methyl ester



A mixture of [2-(4-amino-phenyl)-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-4-yl]acetic acid methyl ester (400 mg, 1.06 mmol.), phenyl isocyanate (254 mg, 2.13 mmol, 2.0 eq) and CH₂Cl₂ (1.0 ml), toluene (2.0 ml) was added a reaction vessel which was flushed with argon. The reaction mixture was stirred at 90° C. for 16 hrs. After cooling and filtering, the precipitate was washed with a minor portion of CH₂Cl₂ again to give a white solid 316 of mg (60.1%).

¹H NMR (500 MHz, DMSO-d₆): δ 8.89 (s, 1H), 8.74 (s, 1H), 7.73 (d, 2H), 7.54 (d, 2H), 7.47 (d, 2H), 7.29 (dd, 2H), 6.98 (d, 1H), 4.02 (s, 2H), 3.73 (d, 4H), 3.66 (d, 4H), 3.65 (s, 3H), 2.08 (s, 3H)

{5-Methylsulfanyl-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidin-4-yl}-acetic acid sodium salt




1N NaOH (1.0 ml, 6.0 eq.) was added to a stirred solution of {5-methyl sulfanyl-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidin-4-yl}-acetic acid methyl ester (80 mg, 0.16 mmol.) in MeOH (2.0 ml) and the mixture was refluxed for 4 hrs. The mixture was cooled to r.t. and filtered. The precipitate was washed with MeOH and dried in vacuo to give a white solid of 61 mg (75.2%).

¹H NMR (500 MHz, DMSO-d₆): δ 7.59 (d, 2H), 7.48 (d, 2H), 7.38 (d, 2H), 7.00 (t, 2H), 6.54 (t, 1H), 3.69 (dd, 8H), 1.96 (s, 3H)

177

5-Methylsulfanyl-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidin-4-yl}-acetic acid




1N NaOH (1.0 ml, 6.0 eq.) was added to a stirred solution of {5-methyl sulfanyl-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidin-4-yl}-acetic acid methyl ester (80 mg, 0.16 mmol.) in MeOH (2.0 ml) and the mixture was refluxed at 0° C. and carefully acidified with conc. HCl. The mixture was filtered and the precipitate was dried in vacuo to give a white solid 64 mg (84.2%).

¹H NMR (500 MHz, DMSO-d₆): δ 9.00 (s, 1H), 8.76 (s, 1H), 7.72 (d, 2H), 7.65-7.46 (m, 4H), 7.29 (t, 2H), 6.99 (t, 1H), 3.93 (s, 2H), 3.72 (d, 4H), 3.65 (d, 4H), 2.08 (s, 3H)

Example 7

Preparation of Formula (I) in Scheme 7


6-Hydroxy-5-methoxy-1H-pyrimidine-2,4-dione



The NaOCH₃ (8.5 g, 157 mmole), was added to a stirred solution of dimethyl 2-methoxymalonate (10.2 g, 62.91 mmole), urea (2.5 g, 41.7 mmole), and EtOH (80 ml). The reaction mixture was heated to 100° C. for 4 hrs, then the reaction was evaporated to remove the solvent. Water was added to the mixture and then concentrated HCl was added. The pH of the mixture was adjusted to ca. 3~4. The mixture was filtered and wash water and acetone. After drying in vacuo, it gave a product (99%).

¹H NMR (500 MHz, DMSO-d₆): δ 8.74 (s, 1H), 3.23 (s, 3H)


2,4,6-Trichloro-5-methoxypyrimidine



¹H NMR (500 MHz, CDCl₃-d₁): δ 3.97 (s, 3H)


178

4-(2,6-Dichloro-5-methoxypyrimidin-4-yl)morpholine




¹H NMR (500 MHz, CDCl₃-d₁): δ 3.87~3.89 (d, 4H), 3.76~3.78 (d, 4H), 3.73 (s, 3H)

4-(4-chloro-5-methoxy-6-morpholinopyrimidin-2-yl)benzenamine



¹H NMR (500 MHz, DMSO-d₆): δ 7.90~7.91 (d, 2H), 6.56~6.58 (d, 2H), 5.65 (br, 2H), 3.79 (s, 4H), 3.73 (s, 4H), 3.67 (s, 3H)


1-[4-(4-Chloro-5-methoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.28 (s, 1H), 9.23 (s, 1H), 8.36~8.37 (d, 2H), 8.15~8.17 (d, 2H), 7.56~7.58 (d, 2H), 7.44~7.46 (d, 2H), 3.84~3.86 (d, 4H), 3.74~3.75 (d, 4H), 3.71 (s, 3H)

179


5-Ethoxy-6-hydroxy-1H-pyrimidine-2,4-dione

¹H NMR (500 MHz, DMSO-d₆): δ 1.06-1.09 (m, 3H), 3.58-3.61 (m, 2H), 9.00 (s, 2H)


2,4,6-Trichloro-5-ethoxy-pyrimidine

¹H NMR (500 MHz, CDCl₃-d): δ 1.47-1.50 (m, 3H), 4.17-4.21 (m, 2H)


4-(2,6-Dichloro-5-ethoxy-pyrimidin-4-yl)-morpholine

¹H NMR (500 MHz, CDCl₃-d₁): δ 1.38-1.40 (m, 3H), 3.76-3.78 (m, 4H), 3.87-3.89 (m, 4H), 3.91-3.95 (m, 2H)


4-(4-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenylamine

**180**¹H NMR (500 MHz, CDCl₃-d₁): δ 1.38-1.41 (m, 3H), 3.81-3.83 (m, 4H), 3.87-3.89 (m, 4H), 3.94-3.98 (m, 2H), 6.72-6.73 (m, 2H), 8.12-8.14 (m, 2H)

1-[4-(4-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea

¹H NMR (500 MHz, DMSO-d₆): δ 1.31-1.34 (m, 3H), 3.73-3.75 (m, 4H), 3.84-3.86 (m, 4H), 3.92-3.96 (m, 4H), 7.45-7.46 (m, 2H), 7.57-7.59 (m, 2H), 8.16-8.18 (m, 2H), 8.37-8.38 (m, 2H), 9.17 (s, 1H), 9.20 (s, 1H)


1-[4-(4-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea

¹H NMR (500 MHz, DMSO-d₆): δ 1.31-1.34 (m, 3H), 3.73-3.75 (m, 4H), 3.84-3.86 (m, 4H), 3.92-3.96 (m, 2H), 7.33-7.34 (m, 1H), 7.56-7.58 (m, 2H), 7.95-7.97 (m, 1H), 8.15-8.17 (m, 2H), 8.21 (m, 1H), 8.62 (s, 1H), 8.92 (s, 1H), 9.10 (s, 1H)

Example 8

Preparation of Compounds of Formula (I) in Scheme 8


5-Methoxy-2-(4-nitro-phenyl)-3H-pyrimidin-4-one



181


A mixture of methyl methoxy acetate (5 mL, 1.0 eq) and ethyl formate (4.07 mL, 1.0 eq) was added dropwise to a slurry of sodium methoxide (5.45 g, 2.0 eq) in toluene (20 mL) in ice both. After dropping at room temperature and stirring overnight, the resulting solution was dried in vacuo. Then a mixture of residue, 4-nitrobenzamide (10.17 g, 1.0 eq) and sodium methoxide (2.72 g, 1.0 eq) in EtOH (80 mL) was refluxed at 110° C. for 6 hrs. After the solvent was dried in vacuo, water and conc. HCl solution (5<pH<4) were added. After formation of white precipitates, the mixture was filtered and dried solid in vacuo. A product was obtained as a white solid (3.21 g, 26%).

4-Chloro-5-methoxy-2-(4-nitro-phenyl)-pyrimidine



In ice bath, dimethyl-phenyl-amine (1.19 mL, 1.0 eq) was added dropwise to a slurry of 5-Methoxy-2-(4-nitro-phenyl)-3H-pyrimidin-4-one (2.3 g, 1.0 eq) in POCl₃ (4.4 mL, 5.0 eq) and the mixture was refluxed overnight. Excess POCl₃ was evaporated in vacuo and the residue was poured into ice-water. After formation of precipitates, the mixture was filtered and the residue was dried solid in vacuo. A product was obtained as a white solid (2.47 g, 61%).

4-[5-Methoxy-2-(4-nitro-phenyl)-pyrimidin-4-yl]-morpholine




4-chloro-5-methoxy-2-(4-nitro-phenyl)-pyrimidine (1.50 g, 1.0 eq) was stirred in dichloromethane (10 mL) and a solution of morpholine (0.75 mL, 1.5 eq) in dichloromethane (5 mL) was added dropwise to the mixture at 0° C. After stirring, the resulting solution was reacted overnight at r.t., NH₄Cl (aq) was added to the solution and the solution was extracted with dichloromethane. The combined organic layers were washed with brine, dried and evaporated in vacuo. A product was obtained as a white solid (0.73 g, 41%).

¹H NMR (500 MHz, DMSO-d₆): δ 3.74 (s, 4H), 3.80 (s, 4H), 3.93 (s, 3H), 8.24 (s, 1H), 8.30 (d, 2H), 8.48 (d, 2H)

182


4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine



4-[5-methoxy-2-(4-nitro-phenyl)-pyrimidin-4-yl]-morpholine (0.2 g, 1 eq) was stirred under hydrogen balloon with palladium on carbon (10 percent, 0.1 g, 0.15 eq) in a mixture of MeOH and EA (1:1, 8 mL) at room temperature overnight. The reaction mixture was then filtered through Celite, volatiles were removed in vacuo, and the residue was purified by flash chromatography to give a white solid (0.15 g, 83%).

¹H NMR (500 MHz, DMSO-d₆): δ 3.71 (s, 8H), 3.83 (s, 3H), 5.54 (s, 2H), 6.58 (d, 2H), 7.95 (d, 2H), 8.04 (s, 1H)


6-Hydroxy-5-methoxy-2-(4-nitro-phenyl)-3H-pyrimidin-4-one



The NaOCH₃ (7.8 g, 3 eq) was added to a stirred solution of 1-ethyl 3-methyl 2-methoxymalonate (11 g, 1.3 eq), 4-nitrobenzimidamide hydrochloride (9.6 g, 1 eq) and EtOH (240 ml) and the mixture was refluxed for 8 hrs. After 8 hrs, the solvent was evaporated and water (10–15 ml) was added to the mixture then the mixture was cooled to 0° C. and carefully acidified with conc. HCl. After overnight cooling, it led to the formation of brown precipitates. The mixture was filtered and the precipitates was dried in vacuo to give a brown solid (65%) as a product.

¹H NMR (500 MHz, DMSO-d₆): δ 3.71 (s, 3H), 8.28–8.35 (m, 4H)

4,6-Dichloro-5-methoxy-2-(4-nitro-phenyl)-pyrimidine




183

The POCl_3 (30 ml, 16.7 eq) was added to a stirred solution of 5-methoxy-2-(4-nitrophenyl)pyrimidine-4,6-diol (5.13 g, 1 eq) and *N,N*-dimethylaniline (3.2 ml, 1.27 eq) the mixture was refluxed overnight. Excess POCl_3 was evaporated in vacuo and the residue was poured into ice-water to give a solid. The solid was filtered and dried in vacuo to give a brown solid (81%) as a product.

^1H NMR (500 MHz, $\text{CDCl}_3\text{-d}_1$): δ 4.03 (s, 3H), 8.31~8.32 (d, 2H), 8.54~8.56 (d, 2H)


4-[6-Chloro-5-methoxy-2-(4-nitro-phenyl)-pyrimidin-4-yl]-morpholine



Morpholine (0.6 ml, 2.2 eq) was added to a stirred solution of 4,6-dichloro-5-methoxy-2-(4-nitrophenyl)pyrimidine (0.87 g, 1 eq) in THF, the mixture was stirred at r.t for 4 hrs, the solvent was evaporated, water was added to the solution and the solution was extracted with EA. The combined organic layers were washed with brine, dried and evaporated in vacuo, the product was obtained as a yellow solid (56%).

^1H NMR (500 MHz, $\text{CDCl}_3\text{-d}_1$): δ 3.08 (s, 3H), 3.84~3.86 (d, 4H), 3.93~3.94 (d, 4H), 8.26~8.27 (d, 2H), 8.46~8.47 (d, 2H)

4-(4-Chloro-5-methoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenylamine




To a stirring solution of 4-(6-chloro-5-methoxy-2-(4-nitro-phenyl)pyrimidin-4-yl) morpholine (200 mg, 1 eq), and acetic acid (1 ml) in a 25 ml single-necked round-bottomed flask equipped with a magnetic stirrer was added Zn dust (78 mg, 2.1 eq). The mixture was stirred at ambient temperature for 1 hr. The mixture was poured into a solution of saturated NaHCO_3 (34 ml). The mixture was filtered and partitioned between CH_2Cl_2 and water. The organic portion was dried (MgSO_4) and concentrated under reduced pressure. The residue was purified by silica gel to obtain a product (45%),

184


^1H NMR (500 MHz, DMSO-d_6): δ 3.67 (s, 3H), 3.73 (d, 4H), 3.79 (d, 4H), 5.65 (s, 2H), 6.56~6.58 (d, 2H), 7.90~7.91 (d, 2H)

5-Ethoxy-6-hydroxy-2-(4-nitro-phenyl)-3H-pyrimidin-4-one



A mixture of 4-nitrobenzamide (8.0 g, 1.0 eq) and 2-ethoxy-malonic acid diethyl ester (10.54 g, 1.3 eq) and sodium methoxide (6.43 g, 3.0 eq) in EtOH (140 mL) was refluxed at 80° C. for 6 hrs. After the solvent was dried in vacuo, water and conc. HCl solution ($5 < \text{pH} < 4$) were added. After formation of white precipitates, the mixture was filtered and dried solid in vacuo. A product was obtained as a white solid (1.85 g, 26%).


4,6-Dichloro-5-ethoxy-2-(4-nitro-phenyl)-pyrimidine



In ice bath, dimethyl-phenyl-amine (0.94 mL, 1.1 eq) was added dropwise to a slurry of 5-ethoxy-6-hydroxy-2-(4-nitro-phenyl)-3H-pyrimidin-4-one (1.85 g, 1.0 eq) in POCl_3 (6.5 mL, 10.32 eq) and the resulting solution was refluxed overnight. Excess POCl_3 was evaporated in vacuo and the residue was poured into ice-water. After formation of precipitates, the mixture was filtered and dried solid in vacuo. A product was obtained as a white solid (0.5 g, 24%).

^1H NMR (500 MHz, DMSO-d_6): δ 1.42 (t, 3H), 4.25 (q, 2H), 8.39 (d, 2H), 8.48 (d, 2H)

4-[6-Chloro-5-ethoxy-2-(4-nitro-phenyl)-pyrimidin-4-yl]-morpholine




185

4,6-dichloro-5-ethoxy-2-(4-nitro-phenyl)-pyrimidine (0.50 g, 1.0 eq) was stirred in dichloromethane (10 mL) and a solution of morpholine (0.21 mL, 1.5 eq) in dichloromethane (5 mL) was added dropwise at 0° C. After stirring the resulting solution overnight at r.t., NH₄Cl (aq) was added to the solution and the solution was extracted with dichloromethane. The combined organic layers were washed with brine, dried and evaporated in vacuo. A product was obtained as a white solid (0.57 g, 81%).

¹H NMR (500 MHz, DMSO-d₆): δ 1.34 (t, 3H), 3.76 (t, 4H), 3.89 (t, 4H), 3.98 (q, 2H), 8.32 (d, 2H), 8.45 (d, 2H)

4-(4-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenylamine




4-[6-chloro-5-ethoxy-2-(4-nitro-phenyl)-pyrimidin-4-yl]-morpholine (0.2 g, 1 eq) was stirred under hydrogen balloon with platinum on carbon (10 percent, 0.11 g, 0.1 eq) in a mixture of MeOH and EA (1:1, 8 mL) at room temperature overnight. The reaction mixture was then filtered through Celite, the volatiles in the mixture was removed in vacuo, and the residue was purified by flash chromatography to give a white solid (0.13 g, 83%).

¹H NMR (500 MHz, CDCl₃-d₁): δ 1.41 (t, 3H), 3.81 (t, 4H), 3.88 (t, 4H), 3.96 (q, 2H), 6.75 (d, 2H), 8.14 (d, 2H)

Example 9

Preparation of Compounds of Formula (I) in Scheme 9

2-Chloro-5-methoxy-4-pyridin-4-yl-pyrimidine




A mixture of 2,4-dichloro-5-methoxy-pyrimidine (3.0 g, 1.0 eq.), 4-pyridinylboronic acid (2.26 g, 1.1 eq.), Pd(PPh₃)₂Cl₂ (0.58 g, 0.05 eq.), K₃PO₄·H₂O (10.67 g, 3.0 eq.), H₂O (3 ml) and 1,4-Dioxane (20.0 ml) was heated at 90° C. and refluxed overnight. The solvent was removed in vacuo, and the residue was extracted with EA and water. The combined organic layers was wash with brine, dried and evaporated in

186

vacuo. The crude was purified by chromatography to give a yellow thick liquid (1.83 g, 49.26%).


¹H NMR (500 MHz, CDCl₃-d₁): δ 8.76-8.75 (d, 2H), 8.42 (s, 1H), 8.02-8.04 (d, 2H), 4.05 (s, 3H).

4-(5-Methoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenylamine




¹H NMR (500 MHz, CDCl₃-d₁): δ 8.75-8.76 (m, 2H), 8.55 (s, 1H), 8.26-8.28 (d, 2H), 8.15-8.17 (m, 2H), 6.76-6.78 (d, 2H), 4.00 (s, 3H).

[4-(5-Methoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester



¹H NMR (500 MHz, CDCl₃-d₁): δ 8.78-8.79 (d, 2H), 8.61 (s, 1H), 8.44-8.45 (d, 2H), 8.22-8.23 (d, 2H), 7.59-7.60 (d, 2H), 7.40-7.43 (m, 2H), 7.21-7.27 (m, 3H), 7.13 (s, 1H), 4.08 (s, 3H).


1-[4-(5-Methoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-carbonyl)-phenyl]-urea



187


¹H NMR (500 MHz, DMSO-d₆): δ 9.94-9.58 (m, 2H), 8.87 (s, 1H), 8.76-8.78 (m, 2H), 8.32-8.34 (m, 2H), 8.12-8.13 (m, 2H), 7.63-7.67 (m, 2H), 7.55-7.56 (m, 2H), 7.38-7.37 (m, 2H), 4.07 (s, 3H), 3.60 (s, 4H), 3.50 (s, 4H).

1-[4-(5-Methoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ 9.32 (s, 1H), 9.12 (s, 1H), 8.86-8.87 (m, 1H), 8.76-8.78 (m, 2H), 8.31-8.34 (m, 2H), 8.12-8.13 (m, 2H), 7.64-7.67 (m, 2H), 7.50-7.52 (m, 2H), 7.28-7.30 (m, 2H), 4.18 (s, 2H), 4.07 (s, 3H), 3.95-3.97 (m, 2H), 3.69-3.71 (m, 2H).

1-[4-(5-Methoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.72 (s, 1H), 9.45 (s, 1H), 8.87 (s, 1H), 8.77-8.78 (m, 2H), 8.33-8.37 (m, 4H), 8.12-8.13 (m, 2H), 7.63-7.65 (d, 2H), 7.46-7.48 (d, 2H), 4.07 (s, 3H).

Example 10

Preparation of Compounds of Formula (I) in Scheme 10


[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester



188


¹H NMR (500 MHz, CDCl₃-d₁): δ 8.76-8.73 (d, 2H), 8.45-8.43 (d, 2H), 7.71-7.68 (m, 2H), 7.57-7.55 (d, 2H), 7.42-7.39 (m, 2H), 7.27-7.20 (m, 3H), 3.91-3.90 (s, 8H), 1.62 (s, 3H)

5 4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenylamine




¹H NMR (500 MHz, CDCl₃-d₁): δ 8.73-8.72 (d, 2H), 8.28-8.26 (d, 2H), 7.69-7.67 (dd, 2H), 6.72-6.71 (s, 2H), 3.95 (s, 2H), 3.89 (s, 8H), 2.04 (s, 3H)

[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ 8.85 (s, 1H), 8.72-8.71 (d, 2H), 8.24-8.22 (d, 2H), 7.68-7.67 (d, 2H), 7.52-7.50 (d, 2H), 5.96 (s, 2H), 3.87-3.86 (d, 4H), 3.80-3.79 (d, 4H), 2.05 (s, 3H)


1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea



189


¹H NMR (500 MHz, DMSO-d₆): δ 9.24 (s, 1H), 9.06 (s, 1H), 8.72-8.71 (d, 2H), 8.29-8.28 (d, 2H), 7.69-7.68 (dd, 2H), 7.61-7.59 (d, 2H), 7.51-7.49 (d, 2H), 7.29-7.27 (d, 2H) 4.18 (s, 2H), 3.96-3.94 (d, 2H), 3.88-3.87 (d, 4H), 3.81-3.80 (d, 4H), 3.70-3.68 (d, 2H), 2.05 (s, 3H)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea




¹H NMR (500 MHz, DMSO-d₆): δ 9.34 (s, 1H), 9.29 (s, 1H), 8.73-8.72 (d, 2H), 8.38-8.37 (d, 2H), 8.33-8.31 (d, 2H), 7.69-7.68 (d, 2H), 7.61-7.59 (d, 2H), 7.48-7.46 (d, 2H), 3.88-3.87 (d, 4H), 3.81-3.80 (d, 4H), 2.06 (s, 3H)

4-{3-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzamide



¹H NMR (500 MHz, DMSO-d₆): δ 9.40 (b, 1H), 9.34 (b, 1H), 8.72-8.71 (d, 2H), 8.29-8.27 (d, 2H), 7.82-7.80 (d, 3H), 7.69-7.67 (d, 2H), 7.63-7.61 (d, 2H), 7.56-7.54 (d, 2H), 7.14 (s, 1H), 3.87-3.86 (d, 4H), 3.81-3.80 (d, 4H), 2.05 (s, 3H)


1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea



190


¹H NMR (500 MHz, DMSO-d₆): δ 9.07 (s, 1H), 8.83 (s, 1H), 8.73-8.72 (d, 2H), 8.30-8.29 (d, 2H), 7.69-7.68 (d, 2H), 7.59-7.58 (d, 2H), 7.47-7.46 (d, 2H), 7.30-7.27 (d, 2H), 6.98 (s, 1H), 3.88-3.87 (d, 4H), 3.81-3.80 (d, 4H), 2.05 (s, 3H)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-thiourea




¹H NMR (500 MHz, DMSO-d₆): δ 10.0 (b, 1H), 9.99 (b, 1H), 8.73-8.72 (d, 2H), 8.32-8.30 (d, 2H), 7.70-7.66 (m, 4H), 7.50-7.49 (d, 2H), 7.35-7.32 (d, 2H), 7.15-7.12 (d, 1H), 3.89-3.88 (d, 4H), 3.81-3.80 (d, 4H), 2.04 (s, 3H)

1-Ethyl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ 8.75 (s, 1H), 8.72-8.71 (d, 2H), 8.24-8.22 (d, 2H), 7.68-7.67 (d, 2H), 7.52-7.50 (d, 2H), 6.22-6.19 (t, 1H), 3.87-3.86 (d, 4H), 3.80-3.79 (d, 4H), 3.13-3.10 (dd, 2H), 2.05 (s, 3H), 1.07-1.04 (t, 3H)


1-Methyl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-urea



191


¹H NMR (500 MHz, DMSO-d₆): δ8.90 (s, 1H), 8.72-8.71 (d, 2H), 8.24-8.22 (d, 2H), 7.68-7.67 (d, 2H), 7.52-7.51 (d, 2H), 6.18-6.17 (d, 1H), 3.87-3.86 (d, 4H), 3.79 (d, 4H), 2.65 (s, 3H), 2.05 (s, 3H)

1-Isopropyl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ8.72-8.71 (d, 2H), 8.62 (s, 1H), 8.24-8.22 (d, 2H), 7.68-7.67 (d, 2H), 7.50-7.48 (d, 2H), 6.11-6.10 (d, 1H), 3.87-3.86 (d, 4H), 3.77-3.74 (d, 4H), 2.05 (s, 3H), 1.11-1.10 (t, 6H), 1.00-0.99 (d, 1H)

1-Isoxazol-3-yl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ9.74 (s, 1H), 9.22 (s, 1H), 8.75-8.72 (m, 3H), 8.32-8.29 (d, 2H), 7.69-7.68 (d, 2H), 7.62-7.58 (d, 2H), 6.87-6.87 (d, 1H), 3.88-3.87 (d, 4H), 3.81-3.80 (s, 4H), 2.06 (s, 3H)

4-(5-Methylsulfanyl-4,6-di-morpholin-4-yl-pyrimidin-2-yl)-phenylamine




192

A mixture of 4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (200 mg, 0.594 mmol.), morpholin (103 mg, 1.118 mmol, 2.0 eq), triethylamine (120 mg, 1.118 mmol, 2.0 eq) in dioxane (8 ml) and the mixture was reacted for 16 hrs at 90° C. The reaction mixture was dried and the solvent was removed by evaporation in vacuo. The reaction mixture was partitioned between EA and water. The organic layer was washed with brine, dried over MgSO₄, filtered and evaporated in vacuo. The resulting residue was purified using flash chromatography on silica gel (EA), and recrystallized by (Hexane/EtOAc) to give a gray white 106 mg (46.1%).

¹H NMR (500 MHz, CDCl₃-d₁): δ8.19-8.17 (d, 2H), 6.69-6.68 (d, 2H), 3.88 (s, 2H), 3.84-3.82 (d, 8H), 3.69-3.68 (d, 8H), 2.2 (s, 3H)


1-{4-[4-(4-Methanesulfonyl-piperazin-1-yl)-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-phenyl-urea



A mixture of 1-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea (120 mg, 0.26 mmol.), 1-methanesulfonyl-piperazine (65 mg, 0.39 mmol, 1.5 eq.), triethylamine (40 mg, 0.39 mmol, 1.5 eq.), and DMF (4.0 ml) was added to the reaction vessel and flushed with argon. The reaction mixture was stirred at 80° C. for 5 hrs and then dried, and the solvent was evaporated. The reaction mixture was partitioned between EA and NH₄Cl_(aq), the organic layer was washed with brine, dried over MgSO₄, filtered and evaporated in vacuo. The resulting residue was purified with flash chromatography on silica gel (Hexane/EtOAc 1:1) to give a pale yellow solid of 77.2 mg (50.3%).

¹H NMR (500 MHz, DMSO-d₆): δ8.87 (s, 1H), 8.73 (s, 1H), 7.57-7.50 (m, 5H), 7.29 (t, 2H), 6.98 (t, 1H), 3.85 (d, 4H), 3.79 (d, 8H), 3.20 (d, 4H), 2.90 (s, 3H), 1.93 (s, 3H)


1-{4-[4-(4-Methyl-piperazin-1-yl)-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-pyridin-3-yl-urea



193


¹H NMR (500 MHz, DMSO-d₆): δ 9.21 (s, 1H), 9.07 (s, 1H), 8.60 (s, 1H), 8.21-8.18 (m, 3H), 7.95 (s, 1H), 7.55-7.53 (d, 2H), 7.31 (d, 1H), 3.72 (b, 4H), 3.60 (s, 8H), 2.44 (b, 4H), 2.21 (s, 3H), 2.14 (s, 3H)

4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenylamine




¹H NMR (500 MHz, CDCl₃-d₁): δ 9.02-9.02 (d, 1H), 8.68-8.67 (dd, 1H), 8.29-8.28 (dd, 2H), 8.14-8.12 (dd, 1H), 7.41-7.38 (s, 1H), 6.73-6.71 (d, 2H), 3.94 (s, 2H), 3.90 (s, 8H), 2.04-2.03 (s, 3H)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea



¹H NMR (500 MHz, DMSO-d₆): δ 8.98 (s, 1H), 8.91-8.90 (d, 1H), 8.74 (s, 1H), 8.69-8.67 (dd, 1H), 8.33-8.31 (d, 2H), 8.16-8.14 (d, 1H), 7.60-7.58 (m, 3H), 7.56-7.53 (dd, 2H), 7.47-7.27 (dd, 2H), 6.98 (s, 1H), 3.88-3.87 (d, 4H), 3.81-3.80 (d, 4H), 2.04 (s, 3H)


1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-thiourea



194


¹H NMR (500 MHz, DMSO-d₆): δ 10.03 (s, 1H), 9.93 (s, 1H), 8.91-8.91 (d, 1H), 8.68-8.67 (dd, 1H), 8.34-8.32 (d, 2H), 8.16-8.154 (d, 1H), 7.68-7.66 (d, 2H), 7.55-7.49 (m, 3H), 7.36-7.33 (d, 2H), 7.15-7.14 (d, 1H), 3.89-3.88 (d, 4H), 3.81-3.80 (d, 4H), 2.04 (s, 3H)

[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester




¹H NMR (500 MHz, CDCl₃-d₁): δ 9.06-9.05 (d, 1H), 8.71-8.70 (d, 1H), 8.46-8.44 (d, 2H), 8.17-8.15 (dd, 1H), 7.58-7.56 (d, 2H), 7.43-7.41 (d, 3H), 7.41-7.39 (d, 1H), 7.27-7.20 (m, 3H), 3.92-3.92 (d, 8H), 2.05-2.04 (s, 3H)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ 9.05 (s, 1H), 8.90-8.87 (d, 2H), 8.68-8.67 (d, 1H), 8.33-8.31 (d, 2H), 8.16-8.14 (d, 1H), 7.60-7.56 (m, 3H), 7.55-7.48 (dd, 2H), 7.30-7.28 (d, 2H), 4.18 (s, 2H), 3.97-3.95 (d, 2H), 3.88-3.87 (d, 4H), 3.81-3.80 (d, 4H), 3.70-3.68 (d, 2H), 2.04 (s, 3H)


1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea



195


¹H NMR (500 MHz, DMSO-d₆): δ9.21 (s, 1H), 9.01 (s, 1H), 8.90 (s, 1H), 8.68-8.67 (d, 1H), 8.62 (d, 1H), 8.33-8.31 (d, 2H), 8.21-8.20 (d, 1H), 8.15-8.14 (d, 1H), 7.97-7.95 (d, 1H), 7.61-7.54 (dd, 2H), 7.34-7.33 (d, 2H), 3.88 (s, 4H), 3.81 (s, 4H), 2.04 (s, 3H)

N-(4-{3-[4-(5 Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-methanesulfonamide



¹H NMR (500 MHz, DMSO-d₆): δ9.34 (b, 1H), 9.05 (s, 1H), 8.90 (s, 1H), 8.82 (s, 1H), 8.67 (s, 1H), 8.32-8.30 (d, 2H), 8.15-8.14 (d, 1H), 7.59-7.57 (d, 2H), 7.54 (s, 1H), 7.44-7.42 (d, 2H), 7.15-7.14 (d, 2H), 3.87 (s, 4H), 3.80 (s, 4H), 2.91 (s, 3H), 2.04 (s, 3H)

4-(5-Methylsulfanyl-2,6-di-pyridin-3-yl-pyrimidin-4-yl)-morpholine




A mixture of 4-(2,6-dichloro-5-methylsulfanyl-pyrimidin-4-yl)-morpholine (200 mg, 0.717 mmol.), 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (323 mg, 1.577 mmol, 2.2 eq), (PPh₃)₄Pd (165 mg, 0.2 mmol.), 2M Na₂CO₃ (2.1 ml, 4.302 mmol, 6.0 eq.) and dioxane (20 ml) was added to a reaction vessel which was then flushed with argon. The reaction mixture was stirred at 90° C. for 16 hrs, and then dried. The solvent was removed by evaporation. The mixture was partitioned between EA and water. The organic layer was collected, washed with brine, dried over MgSO₄, filtered and evaporated in vacuo. The resulting residue was purified using flash chromatography on silica gel (Hexane/EtOAc 2:3, few Et₃N), recrystallized by (Hexane/EtOAc) to give a pale yellow solid 57.8 mg (22.1%).

¹H NMR (500 MHz, CDCl₃-d₁): δ9.52-9.51 (d, 1H), 8.94-8.94 (d, 1H), 8.70-8.66 (m, 3H), 8.19-8.18 (d, 1H), 7.57-7.53 (dd, 2H), 3.92-3.90 (d, 4H), 3.81-3.80 (d, 4H), 2.05 (s, 3H)


196

4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenylamine




¹H NMR (500 MHz, DMSO-d₆): δ1.07-1.10 (m, 3H), 3.66-3.70 (m, 2H), 3.78-3.79 (m, 8H), 5.54 (s, 2H), 6.60-6.62 (m, 2H), 7.52-7.54 (m, 1H), 8.06-8.07 (m, 2H), 8.36-8.38 (m, 1H), 8.65-8.66 (m, 1H), 9.17 (s, 1H)

4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenylamine



¹H NMR (500 MHz, DMSO-d₆): δ1.09-1.13 (m, 3H), 3.67-3.71 (m, 2H), 3.78-3.79 (m, 8H), 5.55 (s, 2H), 6.60-6.62 (m, 2H), 7.97-7.98 (m, 2H), 8.05-8.07 (m, 2H), 8.71-8.73 (m, 2H)


1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea



¹H NMR (500 MHz, DMSO-d₆): δ1.08-1.11 (m, 3H), 3.69-3.73 (m, 2H), 3.79-3.80 (m, 4H), 3.84-3.85 (m, 4H), 7.46-7.47 (m, 2H), 7.56-7.60 (m, 3H), 8.30-8.32 (m, 2H), 8.37-8.41 (m, 3H), 8.67-8.68 (m, 1H), 9.20 (s, 1H), 9.24 (s, 1H), 9.32 (s, 1H)


197

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea




¹H NMR (500 MHz, DMSO-d₆): δ 1.08-1.11 (m, 3H), 3.69-3.73 (m, 2H), 3.79-3.80 (m, 4H), 3.84-3.85 (m, 4H), 7.33 (m, 1H), 7.57-7.59 (m, 3H), 7.98 (m, 1H), 8.20-8.21 (m, 1H), 8.29-8.31 (m, 2H), 8.39 (m, 1H), 8.61-8.62 (m, 1H), 8.67-8.68 (m, 1H), 8.89 (s, 1H), 9.07 (s, 1H), 9.19-9.20 (m, 1H)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea



¹H NMR (500 MHz, DMSO-d₆): δ 1.11-1.14 (m, 3H), 3.70-3.74 (m, 2H), 3.78-3.79 (m, 4H), 3.84-3.85 (m, 4H), 7.45-7.46 (m, 2H), 7.58-7.59 (m, 2H), 7.99-8.00 (m, 2H), 8.30-8.32 (m, 2H), 8.37-8.38 (m, 2H), 8.74-8.75 (m, 2H), 9.20 (s, 1H), 9.25 (s, 1H)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea




¹H NMR (500 MHz, DMSO-d₆): δ 1.11-1.14 (m, 3H), 3.70-3.74 (m, 2H), 3.79-3.80 (m, 4H), 3.84-3.85 (m, 4H), 7.32-7.34 (m, 1H), 7.58-7.59 (m, 2H), 7.95-8.00 (m, 3H),

198


8.20-8.21 (m, 1H), 8.29-8.31 (m, 2H), 8.61-8.62 (m, 1H), 8.74-8.75 (m, 2H), 8.89 (s, 1H), 9.07 (s, 1H)

[4-(4-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester




¹H NMR (500 MHz, DMSO-d₆): δ 1.31-1.34 (m, 3H), 3.73-3.75 (m, 4H), 3.84-3.86 (m, 4H), 3.91-3.96 (m, 4H), 7.24-7.29 (m, 3H), 7.43-7.46 (m, 2H), 7.61-7.63 (m, 2H), 8.18-8.20 (m, 2H)

[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester



¹H NMR (500 MHz, DMSO-d₆): δ 1.11-1.13 (m, 3H), 3.70-3.74 (m, 2H), 3.78-3.79 (m, 4H), 3.84-3.85 (m, 4H), 7.24-7.29 (m, 3H), 7.43-7.46 (m, 2H), 7.62-7.64 (m, 2H), 7.99-8.00 (m, 2H), 8.32-8.34 (m, 2H), 8.73-8.74 (m, 2H), 10.46 (s, 1H)


[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester



199


¹H NMR (500 MHz, DMSO-d₆): δ1.08-1.11 (m, 3H), 3.69-3.73 (m, 2H), 3.78-3.79 (m, 4H), 3.84-3.85 (m, 4H), 7.24-7.29 (m, 3H), 7.43-7.46 (m, 2H), 7.53-7.57 (m, 1H), 7.62-7.64 (m, 2H), 8.32-8.34 (m, 2H), 8.39-8.41 (m, 1H), 8.66-8.67 (m, 1H), 9.20 (s, 1H), 10.45 (s, 1H)

1-[4-(4-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea




¹H NMR (500 MHz, DMSO-d₆): δ1.31-1.34 (m, 3H), 3.68-3.70 (m, 2H), 3.73-3.75 (m, 4H), 3.83-3.84 (m, 4H), 3.91-3.97 (m, 4H), 4.18 (s, 2H), 7.28-7.30 (m, 2H), 7.48-7.49 (m, 2H), 7.55-7.57 (m, 2H), 8.14-8.16 (m, 2H), 8.84 (s, 1H), 9.00 (s, 1H)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea



¹H NMR (500 MHz, DMSO-d₆): δ1.11-1.14 (m, 3H), 3.68-3.74 (m, 4H), 3.78-3.79 (m, 4H), 3.84-3.85 (m, 4H), 3.95-3.97 (m, 2H), 4.18 (s, 2H), 7.28-7.30 (m, 2H), 7.48-7.50 (m, 2H), 7.57-7.59 (m, 2H), 7.99-8.00 (m, 2H), 8.28-8.30 (m, 2H), 8.73-8.74 (m, 2H), 8.83 (s, 1H), 8.98 (s, 1H)

1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea



200

¹H NMR (500 MHz, DMSO-d₆): δ1.08-1.11 (m, 3H), 3.69-3.73 (m, 4H), 3.78-3.79 (m, 4H), 3.84-3.85 (m, 4H), 3.95-3.97 (m, 2H), 4.18 (s, 2H), 7.28-7.30 (m, 2H), 7.48-7.50 (m, 2H), 7.54-7.58 (m, 3H), 8.28-8.30 (m, 2H), 8.39-8.41 (m, 1H), 8.66-8.67 (m, 1H), 8.84 (s, 1H), 8.98 (s, 1H), 9.19-9.20 (m, 1H)

Biological Testing

The compounds of the invention, prepared as described in the examples, were subject to the following series of biological assays. Brief descriptions of different assays are as follows.

(1) Antiproliferation Assay

A549, HCT-116, and A498 cell lines were purchased from the Bioresource Collection and Research Center (BCRC, Taiwan) and cultured in Dulbecco's Modified Eagle Medium (DMEM, Sigma) containing 10% fetal bovine serum (FBS, HyClone). PC-3, U-87 MG, HCC1954 and BT474 cell lines were purchased from the American Type Culture Collection (ATCC, USA). PC-3 was grown in F12K medium (GIBCO) containing 10% FBS. U-87 MG was grown in Eagle's minimum essential medium (MEM, Sigma) supplemented with nonessential amino acids, L-glutamine, sodium pyruvate, and 10% FBS. HCC1954 was grown in RPMI medium (GIBCO) containing 10% FBS. BT-474 was grown in DMEM medium (GIBCO) containing 10% FBS. All cells were incubated in a humidified atmosphere containing 5% CO₂ at 37° C.

Cell lines were inoculated to the wells of a 96-well culture plate with 1000-5000 cells per well. The compounds were prepared and used to treat cells with less than 1% DMSO. After 72 hours of drug treatment, the viability of each cell line was examined by CellTiter 96 Aqueous Non-Radioactive Cell Proliferation Assay (Promega). The living cells were detectable by conversion of MTS into aqueous, soluble formazan. MTS/PMS solution was freshly prepared and 20 μl of MTS/PMS solution was added to each well of the 96-well cultured plate. The assay plate was incubated for 3 hours and the absorbance at 490 nm was measured by EMax ELISA Reader (Molecular Devices). The IC₅₀ values of compounds were determined after carrying out assays at eight serially diluted concentrations of each compound in triplicate, and are the mean of three separate determinations. The results were analyzed using linear regression software (GraphPad Prism 5; GraphPad Software Inc.).

(2) Immunoblotting

For immunoblotting experiments, A549 cells were inoculated to the wells of a 6-well culture plate and cultured overnight. Cells were incubated with the compound for 30 min and 24 hours and whole cell lysate was harvested by adding 1×SDS Sample Buffer (62.5 mM Tris-HCl (pH 6.8 at 25° C.), 2% w/v SDS, 10% glycerol, 50 mM DTT, and 0.01% w/v bromophenol blue or phenol red). Proteins were separated by SDS-PAGE electrophoresis and transferred to PVDF membrane. Protein expression was immunoblotted by various primary antibodies and detected by Immobilon Western Chemiluminescent HRP Substrate (Millipore). Antibody against Akt, phosphor-Akt (Ser473), phosphor-Akt (Thr308), p70 S6 Kinase, phosphor-p70 S6 Kinase (Thr389), 4E-BP1, and phosphor-4E-BP1 (Thr37/46) were purchased from Cell Signaling Technology.

(3) mTOR Kinase Assay

The activity of the compound against mTOR was examined by measuring the incorporation of ³³P from [γ-³³P]-ATP into 4EBP1. His-tagged, recombinant human 4EBP1 was expressed in *E. coli*, purified by nickel-nitrilotriacetic acid (Ni-NTA) resins, and stored at -80° C. Phosphorylation of 4EBP1 by mTOR was assayed in the presence or absence of the compound and performed in a final volume of 25 μl

201

reaction buffer containing 300 ng 4EBP1, 50 ng recombinant mTOR (Invitrogen), 50 mM HEPES (pH 7.5), 1 mM EGTA, 0.01% Polysorbate 20, 10 mM MnCl₂, 2.5 mM DTT, 10 μ M ATP, and 0.5 μ Ci [γ -³³P]-ATP (PerkinElmer) for 30 min at 30° C. The reactions were terminated by adding 3% phosphoric acid. The ³³P labeled 4EBP1 was transferred onto UniFilter-96 GF/B plate (PerkinElmer) and quantified by Top Count Microplate Scintillation Counter (PerkinElmer). For primary screening of kinase activity inhibition, each test compound was evaluated at 10 M in duplicate. The results were the average of duplicate measurements and expressed as percentage inhibition (compound treatment versus DMSO control). The IC₅₀ values of the compounds were determined after carrying out assays at eight serially diluted concentrations of each compound in duplicate. The results were analyzed using linear regression software (GraphPad Prism 5; GraphPad Software Inc.).

(4) PI3K Enzyme Assay

The ability of compounds at 500 nM against phosphatidylinositol 3-kinase (PI3K) was verified by PI3K HTRF Assay. Recombinant PI3K (p110 α /p85 α) and PI3K (p110 β /p85 α) were from Millipore. Enzyme reactions were performed according to the manufacturer's suggestions. Briefly, the compounds of interest dissolved in 100% DMSO were dispensed in all wells, following which the mixture of PIP2 and enzyme was added. PI3K (p110 α /p85 α) and PI3K (p110 β /p85 α) enzyme reactions were initiated in the presence of 10 μ M and 100 μ M ATP, respectively, and incubated for 30 min at room temperature. STOP solution prepared 2 hours prior to use was added to stop reactions, and mixed well. Detection solution prepared 2 hours prior to use was added, and incubated at room temperature for 17 hours in the dark. Fluorescence was measured by Packard Fusion Alpha Microplate Analyzer with excitation at 340 nm and emission at 620 nm and 665 nm. HTRF ratio was calculated as follows:

$$HTRF \text{ ratio} = \frac{\text{emission at } 665 \text{ nm}}{\text{emission at } 620 \text{ nm}} \times 10000$$

Biological Activity

Inhibition of mTOR kinase by various compounds listed above are summarized and shown in Table 1 and Table 2 below. IC₅₀ value is defined as the concentration of the test compound which achieves a half-maximal inhibition of the kinase activity.

Inhibition of PI3K α and PI3K β by mTOR compounds at 500 nM are summarized and shown in Table 3.

Effects on cellular proliferation by the various compounds of the present invention are summarized and shown in Table 4 below.

Inhibition effects of compounds of interest on PI3K/Akt/mTOR signal transduction pathway in PC-3 cell are summarized and shown in Table 5 below.

TABLE 1

Compound	% Inhibition @ 10 μ M
MTR-0037	98%
MTR-0043	96%
MTR-0054	81%
MTR-0069	98%
MTR-0073	86%
MTR-0077	87%
MTR-0079	83%
MTR-0081	95%

202

TABLE 1-continued

Compound	% Inhibition @ 10 μ M
MTR-0083	99%
MTR-0091	92%
MTR-0098	86%
MTR-0102	81%
MTR-0106	96%
MTR-0108	89%
MTR-0116	81%
MTR-0118	94%
MTR-0120	98%
MTR-0121	94%
MTR-0123	91%
MTR-0124	99%
MTR-0125	96%
MTR-0128	99%
MTR-0129	96%
MTR-0132	98%
MTR-0133	93%
MTR-0135	90%
MTR-0137	84%
MTR-0139	85%
MTR-0141	96%
MTR-0142	98%
MTR-0145	100%
MTR-0147	95%
MTR-0149	93%
MTR-0150	87%
MTR-0152	98%
MTR-0159	97%
MTR-162	95%
MTR-0164	91%
MTR-0165	99%
MTR-0167	89%
MTR-0168	91%
MTR-0169	96%
MTR-0172	86%
MTR-173	92%
MTR-0176	96%
MTR-0177	84%
MTR-0178	90%
MTR-0182	99%
MTR-0183	95%
MTR-0184	98%
MTR-0185	99%
MTR-0186	96%
MTR-0191	100%
MTR-0193	81%
MTR-0195	101%
MTR-0197	97%
MTR-0198	87%
MTR-0199	99%
MTR-0200	84%
MTR-0202	96%
MTR-0205	100%
MTR-0210	96%
MTR-0211	98%
MTR-0212	85%
MTR-0213	84%
MTR-0215	96%
MTR-0216	94%
MTR-0220	102%
MTR-0221	101%
MTR-0222	93%
MTR-0225	101%
MTR-0226	90%
MTR-0227	99%
MTR-0228	101%
MTR-0229	100%
MTR-0231	97%
MTR-0232	95%
MTR-0233	91%
MTR-0236	98%
MTR-0237	85%
MTR-0238	95%
MTR-0239	89%
MTR-0241	99%
MTR-0242	96%
MTR-0243	92%

203

TABLE 1-continued

Compound	% Inhibition @ 10 μ M	
MTR-0244	99%	5
MTR-0245	94%	
MTR-0246	96%	
MTR-0248	95%	
MTR-0249	97%	
MTR-0250	98%	10
MTR-0251	98%	
MTR-0252	82%	
MTR-0253	99%	
MTR-0256	90%	
MTR-0257	93%	15
MTR-0261	98%	
MTR-0262	90%	
MTR-0264	86%	
MTR-0265	96%	
MTR-0266	96%	20
MTR-0267	98%	
MTR-0268	97%	
MTR-0270	96%	
MTR-0271	98%	
MTR-0273	99%	25
MTR-0274	99%	
MTR-0275	97%	
MTR-0276	82%	
MTR-0277	99%	
MTR-0278	100%	30
MTR-0280	95%	
MTR-0282	99%	
MTR-0283	100%	
MTR-0286	93%	
MTR-0288	98%	35
MTR-0289	91%	
MTR-0291	89%	
MTR-0293	83%	
MTR-0297	83%	
MTR-0299	97%	40
MTR-0303	90%	
MTR-0304	90%	
MTR-0305	82%	
MTR-0306	90%	
MTR-0307	82%	45
MTR-0308	93%	
MTR-0310	86%	
MTR-0312	94%	
MTR-0313	98%	
MTR-0314	98%	50
MTR-0316	99%	
MTR-0317	93%	
MTR-0318	91%	
MTR-0319	84%	
MTR-0322	94%	55
MTR-0323	100%	
MTR-0324	81%	
MTR-0325	100%	
MTR-0326	101%	
MTR-0328	85%	60
MTR-0329	97%	
MTR-0334	93%	
MTR-0335	89%	
MTR-0336	89%	
MTR-0337	96%	65
MTR-0338	94%	
MTR-0341	83%	
MTR-0343	92%	
MTR-0344	92%	
MTR-0345	95%	
MTR-0347	96%	
MTR-0350	95%	
MTR-0351	99%	
MTR-0352	94%	
MTR-0353	96%	
MTR-0356	82%	
MTR-0358	100%	
MTR-0360	95%	
MTR-0361	95%	
MTR-0362	84%	
MTR-0363	93%	

204

TABLE 1-continued

Compound	% Inhibition @ 10 μ M	
MTR-0364	100%	
MTR-0365	99%	
MTR-0366	85%	
MTR-0368	97%	
MTR-0369	98%	
MTR-0370	97%	
MTR-0372	96%	
MTR-0373	81%	
MTR-0376	96%	
MTR-0377	97%	
MTR-0378	96%	
MTR-0380	91%	
MTR-0381	92%	
MTR-0382	99%	
MTR-0385	98%	
MTR-0387	96%	
MTR-0388	100%	
MTR-0391	100%	
MTR-0392	102%	
MTR-0394	101%	
MTR-0398	89%	
MTR-0400	85%	
MTR-0401	101%	
MTR-0403	88%	
MTR-0404	97%	
MTR-0405	97%	
MTR-0406	97%	
MTR-0407	96%	
MTR-0410	85%	
MTR-0411	93%	
MTR-0412	98%	
MTR-0415	81%	
MTR-0419	82%	
MTR-0420	92%	
MTR-0421	99%	
MTR-0423	95%	
MTR-0424	100%	
MTR-0426	97%	
MTR-0427	97%	
MTR-0430	86%	
MTR-0435	90%	
MTR-0436	99%	
MTR-0438	99%	
MTR-0440	89%	
MTR-0442	96%	
MTR-0443	100%	
MTR-0446	101%	
MTR-0449	100%	
MTR-0452	100%	
MTR-0453	101%	
MTR-0454	100%	
MTR-0455	100%	
MTR-0456	101%	
MTR-0457	98%	
MTR-0459	88%	
MTR-0461	99%	
MTR-0462	99%	
MTR-0463	97%	
MTR-0464	96%	
MTR-0465	97%	
MTR-0468	97%	
MTR-0469	93%	
MTR-0470	88%	
MTR-0471	94%	
MTR-0472	95%	
MTR-0474	88%	
MTR-0476	99%	
MTR-0477	99%	
MTR-0478	99%	
MTR-0479	95%	
MTR-0480	90%	
MTR-0481	98%	
MTR-0483	88%	
MTR-0484	95%	
MTR-0485	100%	
MTR-0486	88%	
MTR-0487	98%	

205

TABLE 1-continued

Compound	% Inhibition @ 10 μ M
MTR-0488	94%
MTR-0489	96%
MTR-0490	102%
MTR-0491	100%
MTR-0492	100%
MTR-0493	99%
MTR-0494	86%
MTR-0495	100%
MTR-0496	95%

TABLE 2

Compounds	IC ₅₀ of Kinase [nM]
MTR-0069	142.6
MTR-0081	292.3
MTR-0083	107.2
MTR-0106	234.9
MTR-0124	102.5
MTR-0128	253.4
MTR-0132	116.3
MTR-0142	257.3
MTR-0152	206.8
MTR-0165	233.2
MTR-0173	198.1
MTR-0176	210.1
MTR-0182	275.3
MTR-0184	102.3
MTR-0185	223.5
MTR-0195	116.4
MTR-0197	110.4
MTR-0199	161.8
MTR-0202	253.6
MTR-0205	116.4
MTR-0211	226.9
MTR-0215	52.5
MTR-0216	133.9
MTR-0220	72.2
MTR-0221	134.6
MTR-0222	11.5
MTR-0225	101.2
MTR-0226	212.6
MTR-0227	18.8
MTR-0228	82.6
MTR-0229	241
MTR-0231	105.2
MTR-0232	184.7
MTR-0233	80.1
MTR-0236	80.7
MTR-0238	225.2
MTR-0241	8.1
MTR-0242	107.2
MTR-0244	18
MTR-0246	228.1
MTR-0248	96.4
MTR-0249	116.7
MTR-0250	150.2
MTR-0251	146.1
MTR-0253	62.5
MTR-0257	74.4
MTR-0261	86.8
MTR-0262	8
MTR-0265	171.9
MTR-0266	256.7
MTR-0267	109.1
MTR-0268	237.7
MTR-0271	89.6
MTR-0273	81.1
MTR-0274	105.6
MTR-0277	78.3
MTR-0278	67
MTR-0282	47.7

206

TABLE 2-continued

Compounds	IC ₅₀ of Kinase [nM]
MTR-0283	30.5
MTR-0288	78.9
MTR-0299	177
MTR-0303	237.9
MTR-0308	48.6
MTR-0313	214.3
MTR-0314	169.7
MTR-0316	93.9
MTR-0322	93
MTR-0323	50.1
MTR-0325	24.8
MTR-0326	40.9
MTR-0337	272.2
MTR-0347	174.5
MTR-0351	65.7
MTR-0353	194.6
MTR-0358	26.8
MTR-0364	70.4
MTR-0365	179.9
MTR-0368	83.4
MTR-0369	214.1
MTR-0370	86
MTR-0372	202.4
MTR-0376	130.3
MTR-0378	133.2
MTR-0380	221.2
MTR-0382	67.7
MTR-0385	215
MTR-0387	140.8
MTR-0388	79.6
MTR-0391	5.4
MTR-0392	19.6
MTR-0394	70.4
MTR-0401	64.8
MTR-0404	148.5
MTR-0405	49.5
MTR-0406	179.7
MTR-0407	219.5
MTR-0412	136.93
MTR-0420	65.7
MTR-0421	58.8
MTR-0424	30.1
MTR-0426	173.9
MTR-0427	97.7
MTR-0436	69.4
MTR-0438	51.1
MTR-0442	39
MTR-0443	76.8
MTR-0446	56.6
MTR-0449	79.2
MTR-0452	155.1
MTR-0453	30.2
MTR-0454	110.9
MTR-0455	42.3
MTR-0456	119.3
MTR-0457	5.8
MTR-0461	138.2
MTR-0462	101.1
MTR-0463	47.8
MTR-0464	75.1
MTR-0465	124.1
MTR-0469	193.5
MTR-0471	252
MTR-0472	117.4
MTR-0476	175.9
MTR-0477	23.7
MTR-0478	25.1
MTR-0481	169.3
MTR-0484	58.1
MTR-0487	86.7
MTR-0488	255.5
MTR-0489	267.7

207

TABLE 3

Compound	% inhibition @ 500 nM	
	p110 α	p110 β
MTR-0043	85.40%	44.80%
MTR-0069	53.80%	4.10%
MTR-0083	44.80%	9.00%
MTR-0091	60.90%	16.20%
MTR-0124	49.10%	-2.60%
MTR-0128	85.80%	17.30%
MTR-0133	70.00%	4.50%
MTR-0135	70.10%	14.20%
MTR-0147	52.40%	1.20%
MTR-0152	83.00%	35.70%
MTR-0162	102.20%	33.90%
MTR-0164	84.70%	42.10%
MTR-0165	100.80%	66.20%
MTR-0168	65.00%	20.20%
MTR-0169	52.70%	3.20%
MTR-0173	82.10%	32.40%
MTR-0178	48.80%	8.20%
MTR-0197	59.80%	20.30%
MTR-0205	42.60%	6.00%
MTR-0211	42.00%	24.40%
MTR-0215	53.60%	24.10%
MTR-0222	44.30%	16.50%
MTR-0225	43.80%	14.30%
MTR-0226	45.70%	25.20%
MTR-0227	87.00%	38.20%
MTR-0228	74.40%	33.20%
MTR-0236	61.20%	17.60%
MTR-0241	94.80%	53.70%
MTR-0242	96.30%	22.80%
MTR-0244	60.00%	26.30%
MTR-0246	43.30%	10.90%
MTR-0248	45.80%	3.60%
MTR-0257	95.50%	62.40%
MTR-0261	58.40%	8.80%
MTR-0262	56.30%	18.00%
MTR-0267	83.70%	54.10%
MTR-0270	41.10%	20.00%
MTR-0271	94.50%	53.80%
MTR-0273	50.20%	28.30%
MTR-0274	59.00%	28.40%
MTR-0277	61.90%	56.50%
MTR-0278	58.60%	31.00%
MTR-0282	76.80%	53.90%
MTR-0283	72.90%	53.40%
MTR-0288	79.20%	32.60%
MTR-0299	44.30%	6.70%
MTR-0308	106.50%	37.50%
MTR-0313	86.90%	30.40%
MTR-0316	55.90%	27.90%
MTR-0322	44.20%	13.20%
MTR-0323	90.80%	49.50%
MTR-0325	91.50%	36.90%
MTR-0326	90.70%	42.30%
MTR-0347	80.60%	63.50%
MTR-0351	52.30%	32.00%
MTR-0358	97.80%	97.80%
MTR-0368	101.40%	60.90%
MTR-0370	61.80%	40.80%
MTR-0372	95.50%	47.90%
MTR-0376	79.40%	47.90%
MTR-0378	84.50%	30.00%
MTR-0380	100.60%	67.00%
MTR-0382	95.90%	53.30%
MTR-0385	64.00%	36.80%
MTR-0387	83.10%	27.70%
MTR-0391	51.70%	29.10%
MTR-0392	46.60%	31.30%
MTR-0394	70.50%	67.20%
MTR-0398	47.00%	21.80%
MTR-0401	41.70%	16.20%
MTR-0404	66.90%	22.70%
MTR-0405	52.50%	23.60%
MTR-0406	78.40%	56.90%
MTR-0407	63.40%	48.60%
MTR-0412	82.70%	53.00%

208

TABLE 3-continued

Compound	% inhibition @ 500 nM	
	p110 α	p110 β
MTR-0420	58.60%	8.30%
MTR-0421	98.70%	51.30%
MTR-0424	71.50%	27.10%
MTR-0426	58.40%	39.70%
MTR-0427	71.10%	76.20%
MTR-0436	71.50%	51.90%
MTR-0442	79.10%	48.50%
MTR-0446	66.30%	40.70%
MTR-0449	75.40%	92.50%
MTR-0454	74.40%	30.30%
MTR-0455	48.90%	19.80%
MTR-0456	52.60%	30.60%
MTR-0457	47.40%	22.30%
MTR-0461	93.40%	46.70%
MTR-0462	67.90%	38.30%
MTR-0469	81.70%	70.70%

TABLE 4

Compound	IC ₅₀ , nM	
	A549	PC3
MTR-0124	566 \pm 12	390 \pm 9
MTR-0128	914 \pm 29	499 \pm 45
MTR-0142	488 \pm 56	416 \pm 90
MTR-0145	1358 \pm 340	432 \pm 20
MTR-0191	373 \pm 14	374 \pm 18
MTR-0197	115 \pm 2	18 \pm 1
MTR-0205	673 \pm 19	130 \pm 5
MTR-0221	710 \pm 76	327 \pm 43
MTR-0222	88 \pm 17	30 \pm 7
MTR-0225	1535 \pm 113	333 \pm 24
MTR-0227	223 \pm 20	49 \pm 3
MTR-0228	1029 \pm 114	193 \pm 8
MTR-0241	129 \pm 5	61 \pm 4
MTR-0242	748 \pm 140	286 \pm 1
MTR-0244	891 \pm 64	70 \pm 9
MTR-0246	1449 \pm 29	433 \pm 10
MTR-0248	145 \pm 5	178 \pm 27
MTR-0250	834 \pm 37	305 \pm 24
MTR-0251	2062 \pm 75	456 \pm 23
MTR-0253	2478 \pm 367	268 \pm 44
MTR-0257	501 \pm 23	119 \pm 14
MTR-0261	512 \pm 25	111 \pm 2
MTR-0262	561 \pm 49	130 \pm 4
MTR-0267	3492 \pm 156	207 \pm 7
MTR-0271	155 \pm 1	97 \pm 4
MTR-0273	660 \pm 40	162 \pm 6
MTR-0274	135 \pm 11	57 \pm 2
MTR-0277	561 \pm 30	86 \pm 3
MTR-0278	67 \pm 3	39 \pm 0
MTR-0280	382 \pm 23	126 \pm 8
MTR-0282	7571 \pm 755	148 \pm 14
MTR-0283	3721 \pm 341	283 \pm 31
MTR-0288	3982 \pm 275	252 \pm 27
MTR-0299	2499 \pm 49	384 \pm 39
MTR-0303	223 \pm 8	46 \pm 1
MTR-0308	219 \pm 11	62 \pm 2
MTR-0322	302 \pm 1	109 \pm 0
MTR-0323	1041 \pm 23	86 \pm 3
MTR-0325	53 \pm 2	16 \pm 1
MTR-0326	1029 \pm 19	147 \pm 10
MTR-0337	457 \pm 38	154 \pm 6
MTR-0347	313 \pm 3	154 \pm 8
MTR-0350	614 \pm 28	188 \pm 6
MTR-0353	557 \pm 7	396 \pm 6
MTR-0358	144 \pm 1	34 \pm 1
MTR-0364	462 \pm 8	142 \pm 14
MTR-0365	74 \pm 5	23 \pm 1
MTR-0368	214 \pm 2	74 \pm 2
MTR-0369	280 \pm 6	44 \pm 4

TABLE 4-continued


Compound	IC ₅₀ , nM	
	A549	PC3
MTR-0370	287 ± 5	92 ± 4
MTR-0372	332 ± 20	204 ± 7
MTR-0376	595 ± 33	142 ± 10
MTR-0377	598 ± 18	205 ± 6
MTR-0378	186 ± 3	113 ± 11
MTR-0382	74 ± 5	19 ± 1
MTR-0385	361 ± 13	396 ± 6
MTR-0387	148 ± 10	88 ± 3
MTR-0388	250 ± 8	136 ± 3
MTR-0391	6 ± 0	8 ± 0
MTR-0392	188 ± 7	22 ± 0
MTR-0394	431 ± 10	71 ± 2
MTR-0398	460 ± 28	179 ± 14
MTR-0401	685 ± 35	169 ± 3
MTR-0404	273 ± 1	109 ± 0
MTR-0405	88 ± 6	74 ± 2
MTR-0406	>10000	285 ± 5
MTR-0407	>10000	386 ± 16
MTR-0420	234 ± 1	148 ± 2
MTR-0421	87 ± 1	37 ± 1
MTR-0424	118 ± 2	51 ± 1
MTR-0426	962 ± 58	282 ± 11
MTR-0427	1798 ± 52	125 ± 2
MTR-0436	299 ± 10	78 ± 4
MTR-0438	116 ± 4	48 ± 1
MTR-0442	392 ± 15	110 ± 10
MTR-0443	529 ± 7	131 ± 6
MTR-0446	379 ± 13	130 ± 8
MTR-0449	1050 ± 12	95 ± 7
MTR-0452	82 ± 3	32 ± 1
MTR-0453	206 ± 1	56 ± 2
MTR-0454	215 ± 1	54 ± 1
MTR-0455	309 ± 16	72 ± 2
MTR-0456	2088 ± 34	362 ± 25
MTR-0457	11 ± 0.2	6.5 ± 0.3
MTR-0461	177 ± 3	64 ± 2
MTR-0462	448 ± 23	182 ± 6
MTR-0463	222 ± 15	32 ± 1
MTR-0464	12 ± 0.8	2.4 ± 0
MTR-0465	197 ± 3	137 ± 2
MTR-0469	39 ± 2	5.7 ± 0
MTR-0471		34 ± 2
MTR-0472		226 ± 19
MTR-0476	137 ± 6	15 ± 0
MTR-0477	122 ± 1	102 ± 1
MTR-0478	107 ± 4	102 ± 1
MTR-0481	608 ± 14	424 ± 26

TABLE 5


compound	IC ₅₀ (nM)		
	p-S6K (mTORC1)	p-Akt S473 (mTORC2)	p-Akt T308 (PI3K)
PI-103	64.86	35.61	86.73
GDC-0941	451.34	11.49	158.49
BEZ-235	<10	28.91	892.1
Rapamycin	<10	<10	>1000
MTR-0227	11.64	184.73	141.29
MTR-0325	<10	74.49	>1000
MTR-0358	<10	491.45	>1000
MTR-0391	<10	19.14	290.64
MTR-0392	35.03	17.91	<10
MTR-0394	<10	11.11	<10

What is claimed is:

1. A compound of formula (I):

R¹ is selected from:

(i) a group of the following formula:



wherein

P is C₆aryl which is optionally substituted by halogen, —OH, —NH₂ or —OC₁-C₆alkyl; an unsubstituted 6-membered heteroaryl comprising two N atoms; an unsubstituted indole group; or an unsubstituted indazole group;

Q is selected from —H, —OR, —SR, —Halo, —NR₃R₄, —OS(O)_mR, —OC(O)R, —OC(O)NHR, —S(O)_mNR₃R₄, —NRC(O)R, —NRS(O)_mR, —NRC(O)NR₃R₄, and —NRC(S)NR₃R₄, wherein each R, R₃, and R₄ is independently selected from

H,

C₁-C₆ alkyl, optionally substituted by halogen, —N(C₁-C₃alkyl)₂,

a 5- or 6-membered carbocyclic group comprising 1-2 heteroatoms selected from N and O,

a 6-membered heterocyclic group comprising 1-2 heteroatoms selected from N and O, optionally substituted by C₁-C₆alkyl, C₆ aryl group optionally substituted by one or two substituents selected from halogen, —OC₁-C₃alkyl, —CF₃, —NH₂, —C(O)NH₂, —NHC(O)C₁-C₃alkyl, —N(C₁-C₃alkyl)₂, —COOH, —SO₂NH₂, —SO₂C₁-C₃alkyl, —NHSO₂C₁-C₃alkyl, —CO₂C₁-C₆alkyl, dioxymethylene group, —NHC(O)CF₃, —C(O)NH(CH₂)₂-N(C₁-C₃alkyl)₂, —O(CH₂)₂N(C₁-C₃alkyl)₂,

a 6-membered heterocyclyl comprising 1-2 heteroatoms selected from N, O and S, optionally substituted oxo, C₁-C₃alkyl, or —SO₂C₁-C₃alkyl, —C(O)-6-membered heterocyclyl, optionally substituted C₁-C₃alkyl,

a 6-membered heteroaryl comprising 1-2 N heteroatoms, optionally substituted by one or two substituents selected from a 6-membered heterocyclyl and —SC₁-C₃alkyl, or

a 5- or 6-membered heteroaryl group comprising 1-2 heteroatoms selected from N, O and S, optionally condensed with benzene ring and optionally substituted by halogen, —CO₂C₁-C₃alkyl, oxo, —NHC(O)C₁-C₃alkyl, C₁-C₃alkyl, a 6-membered heterocyclyl comprising 2 heteroatoms selected from N and O, optionally substituted C₁-C₃alkyl; m is 1 or 2; or R₃ and R₄ together with the nitrogen atom to which they are attached form a saturated 5- or 6-membered N-containing heterocyclic group which is optionally substituted by C₁-C₃alkyl, —SO₂C₁-C₃alkyl, or oxo;

—C(O)R, —C(O)N(R)₂ and —S(O)_mR wherein R and m are as defined above;

Y is selected from —O—(CH₂)_n—, —S—(CH₂)_n—, and —S(O)_m(CH₂)_n— wherein m is 1 or 2, n is 0 or an integer of 1 to 2, and R² is selected from H or a group —NR₃R₄ wherein R₃ and R₄ are as defined above;

Z is selected from halo, —(CH₂)_sCOOR, —(CH₂)_sCONR₃R₄, and —(CH₂)_sCH₂NR₃R₄, wherein s is 0 or an integer of 1 to 2 and wherein R, R₃ and R₄ are as defined above; an unsubstituted 6-membered heteroaryl comprising one heteroatom N, an optionally substituted 6-membered heterocyclyl comprising two heteroatoms selected from N and O, wherein the substituent is selected from C₁-C₃alkyl and C₁-C₃alkylsulfonyl; and

W is selected from a morpholine ring and a pyridine ring; or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1, wherein P is a C₆ aryl optionally substituted by halogen, —OH, —NH₂ or —OC₁-C₆alkyl.

3. A compound according to claim 1, wherein W is a morpholine ring.

4. A compound according to claim 1, wherein Z is selected from halo and —(CH₂)_sCOOR.

5. A compound according to claim 1 wherein Q is —NRC(O)N—R₃R₄.

6. A compound according to claim 1, wherein Y is —O—(CH₂)_n— or —S—(CH₂)_n—.

7. A compound selected from:

5-Ethoxy-2-(3-hydroxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0037)

5-Ethoxy-2-(4-hydroxy-3-methoxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0038)

5-Ethoxy-2-(3-fluoro-4-methoxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0039)

2-(3-Amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0040)

3-[4-(2-Dimethylamino-ethoxy)-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl]-phenol (MTR-0043)

3-[4-(2-Dimethylamino-ethoxy)-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl]-phenol (MTR-0046)

3-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0047)

2-(3,5-Difluoro-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0053)

5-Ethoxy-2-(1H-indol-5-yl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0054)

5-Ethoxy-2-(1H-indol-6-yl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0055)

2-Benzo[1,3]dioxol-5-yl-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0056)

5-Ethoxy-2-(1H-indazol-4-yl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0057)

5-Ethoxy-2-(2-fluoro-3-methoxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0058)

2-{3-[3-(4-Chloro-3-trifluoromethyl-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0059)

5-Ethoxy-6-morpholin-4-yl-2-[3-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0060)

2-(4-Amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0068)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0069)

2-{4-[3-(4-Chloro-3-trifluoromethyl-phenyl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0070)

4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0071)

2-(3-Hydroxy-phenyl)-5-methoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0073)

5-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-1H-indole (MTR-0074)

2-(1H-Indazol-4-yl)-5-methoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0075)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-chloro-3-trifluoromethyl-phenyl)-urea (MTR-0076)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(6-chloro-pyridin-3-yl)-urea (MTR-0077)

N-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-acetamide (MTR-0078)

2-{4-[3-(6-Chloro-pyridin-3-yl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0079)

5-Ethoxy-2-[4-(3-ethyl-ureido)-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0081)

[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0083)

4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-1H-indazole (MTR-0084)

Methanesulfonic acid 3-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester (MTR-0086)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea (MTR-0091)

3-(4-Chloro-5-methanesulfonyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenol (MTR-0094)

N-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-methanesulfonamide (MTR-0096)

4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenol (MTR-0098) Methanesulfonic acid 4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester (MTR-0100)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-thioureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0102)

2-(4-Benzenesulfonylamino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0103)

1-(2-Chloro-ethyl)-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0108)

5-Ethoxy-2-(4-hydroxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0111)

5-Ethoxy-2-(4-ethylcarbamoyloxy-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0112)

5-Ethoxy-6-morpholin-4-yl-2-(4-phenylcarbamoyloxy-phenyl)-pyrimidine-4-carboxylic acid ethyl ester (MTR-0113)

(2-Chloro-ethyl)-carbamic acid 4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester (MTR-0114)

213

Benzenesulfonic acid 4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester (MTR-0115)

3-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenol (MTR-0116)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid (MTR-0118)

N-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-benzenesulfonamide (MTR-0119)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid diethylamide (MTR-0120)

5-Ethoxy-6-morpholin-4-yl-2-(4-ureido-phenyl)-pyrimidine-4-carboxylic acid ethyl ester (MTR-0121)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-dimethylamino-phenyl)-urea (MTR-0123)

1-[4-[5-Ethoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl]-3-phenyl-urea (MTR-0124)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid amide (MTR-0125)

Phenyl-carbamic acid 4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester (MTR-0127)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea (MTR-0128)

1-Ethyl-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0129)

Ethyl-carbamic acid 4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl ester (MTR-0130)

4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-benzenesulfonamide (MTR-0131)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-ethyl-urea (MTR-0132)

5-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-pyridin-2-ylamine (MTR-0133)

1-[5-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-pyridin-2-yl]-3-phenyl-urea (MTR-0134)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-fluoro-phenyl)-urea (MTR-0135)

1-[5-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-pyridin-2-yl]-3-ethyl-urea (MTR-0136)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3-fluoro-phenyl)-urea (MTR-0137)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3,4-difluoro-phenyl)-urea (MTR-0138)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid (2-morpholin-4-ylethyl)-amide (MTR-0139)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid (2-pyrrolidin-1-ylethyl)-amide (MTR-0140)

1-[4-[5-Ethoxy-4-morpholin-4-yl-6-(pyrrolidine-1-carbonyl)-pyrimidin-2-yl]-phenyl]-3-phenyl-urea (MTR-0141)

1-[4-[5-Ethoxy-4-morpholin-4-yl-6-(piperidine-1-carbonyl)-pyrimidin-2-yl]-phenyl]-3-phenyl-urea (MTR-0142)

1-[5-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-pyridin-2-yl]-3-(3-fluoro-phenyl)-urea (MTR-0143)

5-Ethoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid (2-diethylaminoethyl)-amide (MTR-0144)

214

1-[4-[5-Ethoxy-4-(4-methanesulfonyl-piperazine-1-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl]-3-phenyl-urea (MTR-0145)

2-(6-Amino-pyridin-3-yl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0146)

5-Ethoxy-2-[4-[3-(3-fluoro-phenyl)-ureido]-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0147)

5-Ethoxy-6-morpholin-4-yl-2-[6-(3-phenyl-ureido)-pyridin-3-yl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0148)

5-Ethoxy-2-[4-[3-(4-fluoro-phenyl)-ureido]-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0149)

2-[4-[3-(3,4-Difluoro-phenyl)-ureido]-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0150)

1-(4-Fluoro-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0152)

4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0153)

4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0154)

4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenol (MTR-0155)

5-Ethoxy-2-[6-[3-(4-fluoro-phenyl)-ureido]-pyridin-3-yl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0156)

2-[6-[3-(3,4-Difluoro-phenyl)-ureido]-pyridin-3-yl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0157)

5-Ethoxy-2-[6-[3-(3-fluoro-phenyl)-ureido]-pyridin-3-yl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0158)

1-Ethyl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0159)

4-[5-Ethoxy-4-(4-methanesulfonyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-1H-indazole (MTR-0160)

4-[5-Ethoxy-4-(4-methanesulfonyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenylamine (MTR-0161)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea (MTR-0162)

4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-benzenesulfonamide (MTR-0163)

1-(3-Fluoro-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0164)

1-(3,4-Difluoro-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0165)

[2-(4-Amino-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidin-4-yl]-methanol (MTR-0166)

1-(4-Fluoro-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thiourea (MTR-0167)

1-(3-Fluoro-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thiourea (MTR-0168)

1-[4-[5-Ethoxy-4-(4-methanesulfonyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl]-3-phenyl-urea (MTR-0169)

4-[5-Ethoxy-4-(4-methyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-1H-indazole (MTR-0170)

4-[5-Ethoxy-4-(4-methyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenylamine (MTR-0171)

1-[4-[5-Ethoxy-4-(4-methyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl]-3-phenyl-urea (MTR-0172)

215

- 1-{4-[4-(4-Methanesulfonyl-piperazin-1-yl)-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-phenyl-urea (MTR-0173)
- 4-(5-Ethoxy-4-morpholin-4-yl-6-morpholin-4-ylmethylpyrimidin-2-yl)-1H-indazole (MTR-0174)
- 4-(5-Ethoxy-4-morpholin-4-yl-6-morpholin-4-ylmethylpyrimidin-2-yl)-phenylamine (MTR-0175)
- 1-[4-(5-Ethoxy-4-morpholin-4-yl-6-morpholin-4-ylmethylpyrimidin-2-yl)-phenyl]-3-phenyl-urea (MTR-0176)
- 1-(3,4-Difluoro-phenyl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0177)
- 1-(4-Fluoro-phenyl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0178)
- 3-[5-Ethoxy-4-(4-methanesulfonyl-piperazin-1-ylmethyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenol (MTR-0180)
- 2-(4-Amino-phenyl)-5-methoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0181)
- 5-Methoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0182)
- 5-Methoxy-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidine-4-carboxylic acid (MTR-0183)
- 1-[4-[5-Methoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl]-3-phenyl-urea (MTR-0184)
- 1-{4-[4-(4-Methanesulfonyl-piperazine-1-carbonyl)-5-methoxy-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl}-3-phenyl-urea (MTR-0185)
- 1-[4-[5-Ethoxy-4-(4-methyl-piperazine-1-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl]-3-phenyl-urea (MTR-0186)
- [2-(4-Amino-phenyl)-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-4-yl]-acetic acid methyl ester (MTR-0187)
- {5-Methylsulfanyl-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidin-4-yl}-acetic acid methyl ester (MTR-0188)
- {5-Methylsulfanyl-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidin-4-yl}-acetic acid (MTR-0189)
- 2-[5-Methylsulfanyl-6-morpholin-4-yl-2-[4-(3-phenyl-ureido)-phenyl]-pyrimidin-4-yl]-acetamide (MTR-0192)
- 1-[4-(5-Ethoxy-4-morpholin-4-yl-6-piperidin-1-ylmethylpyrimidin-2-yl)-phenyl]-3-phenyl-urea (MTR-0193)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-methyl-urea (MTR-0195)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-morpholin-4-yl-urea (MTR-0196)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0197)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-morpholin-4-yl-ethyl)-urea (MTR-0198)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(2-dimethylamino-ethoxy)-phenyl]-urea (MTR-0199)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-thiazol-2-yl-urea (MTR-0200)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-methyl-piperazin-1-yl)-urea (MTR-0201)

216

- 5-Ethoxy-2-[4-(3-methyl-ureido)-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0202)
- 5-Ethoxy-2-[4-[(morpholine-4-carbonyl)-amino]-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0203)
- 2-[4-[3-(3-Dimethylamino-propyl)-ureido]-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0204)
- 5-Ethoxy-6-morpholin-4-yl-2-[4-(3-pyridin-3-yl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0205)
- 5-Ethoxy-2-[4-[3-(4-methyl-piperazin-1-yl)-ureido]-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0206)
- 1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-morpholin-4-yl-ethyl)-urea (MTR-0207)
- 1-(3-Dimethylamino-propyl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0208)
- 1-(4-Methyl-piperazin-1-yl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0209)
- 1-Methyl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0210)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3-fluoro-4-morpholin-4-yl-phenyl)-urea (MTR-0211)
- 4-[3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido]-benzoic acid ethyl ester (MTR-0212)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0213)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3-trifluoromethyl-phenyl)-urea (MTR-0214)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-methyl-piperazin-1-yl)-phenyl]-urea (MTR-0215)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-thiomorpholin-4-yl-phenyl)-urea (MTR-0216)
- 5-Ethoxy-6-morpholin-4-yl-2-[4-[3-(2-morpholin-4-yl-ethyl)-ureido]-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0217)
- 5-Ethoxy-6-morpholin-4-yl-2-(2-phenylamino-1H-benzimidazol-5-yl)-pyrimidine-4-carboxylic acid ethyl ester (MTR-0218)
- 5-Ethoxy-6-morpholin-4-yl-2-[4-(3-morpholin-4-yl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0219)
- 5-Ethoxy-6-morpholin-4-yl-2-[4-(3-thiazol-2-yl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0220)
- 5-Ethoxy-2-[4-[3-(3-fluoro-4-morpholin-4-yl-phenyl)-ureido]-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0221)
- 5-Ethoxy-6-morpholin-4-yl-2-(4-[3-(3-oxo-morpholin-4-yl)-phenyl]-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0222)
- 5-Ethoxy-6-morpholin-4-yl-2-[4-[3-(3-trifluoromethyl-phenyl)-ureido]-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0223)
- 1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-morpholin-4-yl-urea (MTR-0224)

217

- 1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0225)
- 1-(3-Fluoro-4-morpholin-4-yl-phenyl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0226)
- 1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0227)
- 1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0228)
- 1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-methyl-urea (MTR-0229)
- 5-Ethoxy-6-morpholin-4-yl-2-[4-(3-pyridin-2-yl-ureido)-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0230)
- 5-Ethoxy-2-(4-{3-[4-(4-methyl-piperazin-1-yl)-phenyl]-ureido}-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0231)
- 5-Ethoxy-6-morpholin-4-yl-2-[4-[3-(4-thiomorpholin-4-yl-phenyl)-ureido]-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0232)
- 5-Ethoxy-6-morpholin-4-yl-2-[4-[3-(4-morpholin-4-yl-phenyl)-ureido]-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0233)
- 2-[4-[3-(2-Amino-phenyl)-thioureido]-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0234)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-morpholin-4-yl-phenyl)-urea (MTR-0235)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-urea (MTR-0236)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(6-morpholin-4-yl-pyridin-3-yl)-urea (MTR-0237)
- 1-[4-[4-(4-Methyl-piperazin-1-yl)-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl]-3-pyridin-3-yl-urea (MTR-0238)
- 1-(3,4-Dimethoxy-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0239)
- 1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3-trifluoromethyl-phenyl)-urea (MTR-0240)
- 1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0241)
- 1-(3-Fluoro-4-morpholin-4-yl-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0242)
- 2-[4-(1H-Benzimidazol-2-ylamino)-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0243)
- 2-[4-[3-(4-Carbamoyl-phenyl)-ureido]-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0244)
- 5-Ethoxy-6-morpholin-4-yl-2-[4-(3-pyridin-3-yl-ureido)-phenyl]-pyrimidine-4-carboxylic acid (MTR-0245)
- 1-[4-(4-Methyl-piperazin-1-yl)-phenyl]-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0246)
- 1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-thiomorpholin-4-yl-phenyl)-urea (MTR-0247)
- 1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-morpholin-4-yl-phenyl)-urea (MTR-0248)

218

- 1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-thiazol-2-yl-urea (MTR-0249)
- 2-[4-[3-(3-Acetylamino-phenyl)-ureido]-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0250)
- 2-[4-[3-(3-Carbamoyl-phenyl)-ureido]-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0251)
- 5-Ethoxy-6-morpholin-4-yl-2-(4-{3-[3-(2,2,2-trifluoroacetylamino)-phenyl]-ureido}-phenyl)-pyrimidine-4-carboxylic acid ethyl ester (MTR-0252)
- 1-[4-[5-Ethoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl]-3-pyridin-3-yl-urea (MTR-0253)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-2-yl-urea (MTR-0254)
- 2-[3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido]-thiophene-3-carboxylic acid methyl ester (MTR-0255)
- 1-Benzo[1,3]dioxol-5-yl-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0256)
- 4-[3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido]-benzamide (MTR-0257)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3,4-dimethoxy-phenyl)-urea (MTR-0258)
- 1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-morpholin-4-yl-ethyl)-urea (MTR-0259)
- 1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-penta-2,4-dienyl]-3-(4-methyl-piperazin-1-yl)-urea (MTR-0260)
- 1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-morpholin-4-yl-phenyl)-urea (MTR-0261)
- 1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-thiomorpholin-4-yl-phenyl)-urea (MTR-0262)
- 1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(3-trifluoromethyl-phenyl)-urea (MTR-0263)
- 4-[3-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido]-benzoic acid ethyl ester (MTR-0264)
- 1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(6-morpholin-4-yl-pyridin-3-yl)-urea (MTR-0265)
- 1-[6-(4-Methyl-piperazin-1-yl)-pyridin-3-yl]-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0266)
- 4-[3-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido]-benzamide (MTR-0267)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[5-(4-methyl-piperazin-1-yl)-pyridin-2-yl]-urea (MTR-0268)
- N-(3-[3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido]-phenyl)-2,2,2-trifluoro-acetamide (MTR-0269)
- N-(3-[3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido]-phenyl)-acetamide (MTR-0270)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-methyl-piperazine-1-carbonyl)-phenyl]-urea (MTR-0271)
- 4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester (MTR-0272)

219

- 5-Ethoxy-6-morpholin-4-yl-2-[4-[3-(4-sulfamoyl-phenyl)-ureido]-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0273)
- 5-Ethoxy-2-(4-[3-[4-(4-methyl-piperazine-1-carbonyl)-phenyl]-ureido]-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0274)
- 5-Ethoxy-2-(4-[3-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-ureido]-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0275)
- 5-Ethoxy-6-morpholin-4-yl-2-[4-[3-(6-morpholin-4-yl-pyridin-3-yl)-ureido]-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0276)
- 2-(4-[3-[4-(2-Dimethylamino-ethylcarbamoyl)-phenyl]-ureido]-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0277)
- 5-Ethoxy-2-(4-[3-[4-(morpholine-4-carbonyl)-phenyl]-ureido]-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0278)
- 4-[5-Ethoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl]-carbamic acid phenyl ester (MTR-0279)
- 1-[4-[5-Ethoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0280)
- 4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester (MTR-0281)
- 4-[3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido]-benzamide (MTR-0282)
- 4-[3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido]-benzenesulfonamide (MTR-0283)
- 4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-2-nitrophenylamine (MTR-0284)
- 1-(2-Amino-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thiourea (MTR-0285)
- 4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0286)
- 1-(2-Amino-phenyl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thiourea (MTR-0287)
- 4-[3-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido]-benzenesulfonamide (MTR-0288)
- (1H-Benzimidazol-2-yl)-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-amine (MTR-0289)
- 4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester (MTR-0290)
- 1-(6-Bromo-pyridin-3-yl)-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0291)
- 1-(2-Amino-phenyl)-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thiourea (MTR-0292)
- 4-[3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-thiourea]-benzoic acid (MTR-0293)
- 5-Ethoxy-2-(4-[3-[5-(4-methyl-piperazin-1-yl)-pyridin-2-yl]-ureido]-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0294)
- 5-Ethoxy-2-[4-[3-(2-methoxycarbonyl-thiophen-3-yl)-ureido]-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0296)
- 5-Ethoxy-2-[4-[3-(1H-indazol-4-yl)-ureido]-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0297)
- 5-Ethoxy-2-[4-[3-(4-methyl-1H-benzotriazol-5-yl)-ureido]-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0298)

220

- 5-Ethoxy-6-morpholin-4-yl-2-[4-[3-(2-oxo-2,3-dihydrobenzooxazol-5-yl)-ureido]-phenyl]-pyrimidine-4-carboxylic acid ethyl ester (MTR-0299)
- 2-(4-Amino-3-fluoro-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0300)
- 5-Ethoxy-2-(3-fluoro-4-phenoxy-carbonylamino-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0301)
- 5-Ethoxy-2-(3-fluoro-4-[3-[4-(3-oxo-morpholin-4-yl)-phenyl]-ureido]-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0302)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-thiourea (MTR-0303)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-thiourea (MTR-0304)
- (1H-Benzimidazol-2-yl)-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-amine (MTR-0305)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-isopropyl-urea (MTR-0306)
- 4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid (MTR-0307)
- 1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-carbonyl)-phenyl]-urea (MTR-0308)
- 4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-benzene-1,2-diamine (MTR-0309)
- (1H-Benzimidazol-2-yl)-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-amine (MTR-0310)
- 5-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-1H-benzimidazol-2-ylamine (MTR-0311)
- 1-[4-(2-Dimethylamino-ethoxy)-phenyl]-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0312)
- 3-[3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido]-benzamide (MTR-0313)
- 1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-thiazol-2-yl-urea (MTR-0314)
- 1-(3,4-Dimethoxy-phenyl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0315)
- 4-[3-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido]-benzoic acid (MTR-0316)
- 1-Benzo[1,3]dioxol-5-yl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0317)
- 1-[5-(4-Methyl-piperazin-1-yl)-pyridin-2-yl]-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0318)
- 1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-thiourea (MTR-0319)
- 2-(4-Amino-2-fluoro-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0320)
- 5-Ethoxy-2-(2-fluoro-4-phenoxy-carbonylamino-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0321)
- 5-Ethoxy-2-(2-fluoro-4-[3-[4-(3-oxo-morpholin-4-yl)-phenyl]-ureido]-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0322)
- 4-[3-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido]-benzamide (MTR-0323)

221

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(4-thiomorpholin-4-yl-phenyl)-urea (MTR-0324)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0325)

4-{3-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzenesulfonamide (MTR-0326)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-2-fluoro-phenylamino-urea (MTR-0327)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(5-morpholin-4-yl-pyridin-2-yl)-urea (MTR-0328)

1-(4-Amino-phenyl)-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0329)

1-tert-Butyl-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0330)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-amino-urea (MTR-0331)

4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenylamine (MTR-0332)

1-tert-Butylamino-3-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0333)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[5-(4-methyl-piperazin-1-yl)-pyridin-2-yl]-urea (MTR-0334)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-thiourea (MTR-0335)

4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0336)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-methyl-piperazin-1-yl)-phenyl]-urea (MTR-0337)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[6-(4-methyl-piperazin-1-yl)-pyridin-3-yl]-urea (MTR-0338)

1-Isopropyl-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0339)

1-(6-Bromo-pyridin-3-yl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0340)

1-Isopropyl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0341)

1-tert-Butyl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0342)

1-[4-(2-Dimethylamino-ethoxy)-phenyl]-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0343)

2-[4-[3-(4-Carbamoyl-phenyl)-ureido]-3-fluoro-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0344)

2-[4-[3-(4-Carbamoyl-phenyl)-ureido]-2-fluoro-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0345)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(1H-indazol-4-yl)-urea (MTR-0346)

N-(4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-methanesulfonamide (MTR-0347)

1,3-Bis-[4-(4-chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0349)

1-(4-Amino-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0350)

222

4-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzoic acid (MTR-0351)

1-(6-Bromo-pyridin-3-yl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0352)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(6-morpholin-4-yl-pyridin-3-yl)-urea (MTR-0353)

4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenyl]-carbamic acid phenyl Ester (MTR-0354)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0355)

4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenyl]-urea (MTR-0356)

4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-2-fluoro-phenylamine (MTR-0357)

4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-benzenesulfonamide (MTR-0358)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-3-fluoro-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0359)

4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0360)

2-[4-(3-Benzo[1,3]dioxol-5-yl-ureido)-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0361)

5-Ethoxy-2-[4-[3-(4-ethoxycarbonyl-phenyl)-ureido]-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0362)

2-(4-{3-[4-(2-Dimethylamino-ethoxy)-phenyl]-ureido}-phenyl)-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0363)

4-(3-{4-[5-Ethoxy-4-(morpholine-4-carbonyl)-6-morpholin-4-yl-pyrimidin-2-yl]-phenyl]-ureido)-benzamide (MTR-0364)

5-Ethoxy-2-[4-[3-(4-methanesulfonyl-phenyl)-ureido]-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0365)

5-Ethoxy-2-[4-(3-isopropyl-ureido)-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0366)

2-[4-(3-tert-Butyl-ureido)-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0367)

N-(4-{3-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-acetamide (MTR-0368)

5-Ethoxy-2-[4-[3-(4-methanesulfonylamino-phenyl)-ureido]-phenyl]-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0369)

2-[4-[3-(4-Acetylamino-phenyl)-ureido]-phenyl]-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0370)

N-(4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-acetamide (MTR-0371)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzooxazol-5-yl)-urea (MTR-0372)

6-Chloro-5-methylsulfanyl-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-ylamine (MTR-0373)

1,3-Bis-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0374)

3-Fluoro-4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0375)

223

N-(4-{3-[4-(5-Methylsulfanyl-4-morpholin-4-yl)-pyrimidin-2-yl]-phenyl}-ureido)-phenyl-methanesulfonamide (MTR-0376)

N-(3-{3-[4-(5-Methylsulfanyl-4-morpholin-4-yl)-pyrimidin-2-yl]-phenyl}-ureido)-phenyl)-acetamide (MTR-0377)

1-[3-Fluoro-4-(5-methylsulfanyl-4-morpholin-4-yl)-pyrimidin-2-yl]-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0378)

N-(4-{3-[3-Fluoro-4-(5-methylsulfanyl-4-morpholin-4-yl)-pyrimidin-2-yl]-phenyl}-ureido)-phenyl)-acetamide (MTR-0379)

N-(4-{3-[4-(5-Methylsulfanyl-4-morpholin-4-yl)-pyrimidin-2-yl]-phenyl}-ureido)-phenyl)-acetamide (MTR-0380)

1-[4-(5-Methanesulfonyl-4-morpholin-4-yl)-pyrimidin-2-yl]-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0381)

1-[4-(5-Methoxy-4-morpholin-4-yl)-pyrimidin-2-yl]-phenyl]-3-[4-(morpholine-4-carbonyl)-phenyl]-urea (MTR-0382)

2-Fluoro-4-(5-methoxy-4-morpholin-4-yl)-pyrimidin-2-yl)-phenylamine (MTR-0383)

3-Fluoro-4-(5-methoxy-4-morpholin-4-yl)-pyrimidin-2-yl)-phenylamine (MTR-0384)

4-{3-[2-Fluoro-4-(5-methoxy-4-morpholin-4-yl)-pyrimidin-2-yl]-phenyl}-ureido)-benzamide (MTR-0385)

1-[4-(5-Methoxy-4-morpholin-4-yl)-pyrimidin-2-yl]-phenyl]-3-pyridin-2-yl-urea (MTR-0386)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl)-pyrimidin-2-yl]-phenyl]-3-(4-methanesulfonyl-phenyl)-urea (MTR-0387)

[4-(5-Methylsulfanyl-4-morpholin-4-yl)-6-pyridin-4-yl-pyrimidin-2-yl]-phenyl]-urea (MTR-0388)

1-(6-Chloro-5-methylsulfanyl-4-morpholin-4-yl)-[2,5']bipyrimidinyl-2'-yl)-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0389)

[4-(5-Methylsulfanyl-4-morpholin-4-yl)-6-pyridin-4-yl-pyrimidin-2-yl]-phenyl]-carbamic acid phenyl ester (MTR-0390)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl)-6-pyridin-4-yl-pyrimidin-2-yl]-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0391)

4-{3-[4-(5-Methylsulfanyl-4-morpholin-4-yl)-6-pyridin-4-yl-pyrimidin-2-yl]-phenyl}-ureido)-benzamide (MTR-0392)

N-(4-{3-[4-(5-Ethoxy-4-morpholin-4-yl)-pyrimidin-2-yl]-phenyl}-ureido)-phenyl)-methanesulfonamide (MTR-0394)

1-[4-(5-Ethoxy-4-morpholin-4-yl)-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzooxazol-5-yl)-urea (MTR-0395)

4-(5-Ethoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0396)

[4-(5-Ethoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-carbamic acid phenyl ester (MTR-0397)

1-[4-(5-Ethoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0398)

1-[4-(5-Ethoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea (MTR-0399)

4-{3-[4-(5-Ethoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-ureido)-benzamide (MTR-0400)

1-[4-(5-Methanesulfonyl-4-morpholin-4-yl)-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0401)

224

2'-Amino-5-ethoxy-6-morpholin-4-yl-[2,5']bipyrimidinyl-4-carboxylic acid ethyl ester (MTR-0402)

5-Ethoxy-6-morpholin-4-yl-2'-[3-[4-(3-oxo-morpholin-4-yl)-phenyl]-ureido)-[2,5']bipyrimidinyl-4-carboxylic acid ethyl ester (MTR-0403)

1-(4-Methanesulfonyl-phenyl)-3-[4-(5-methylsulfanyl-4-morpholin-4-yl)-pyrimidin-2-yl)-phenyl]-urea (MTR-0404)

1-[2-Fluoro-4-(5-methoxy-4-morpholin-4-yl)-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0405)

1-[3-Fluoro-4-(5-methoxy-4-morpholin-4-yl)-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0406)

4-{3-[3-Fluoro-4-(5-methoxy-4-morpholin-4-yl)-pyrimidin-2-yl)-phenyl]-ureido)-benzamide (MTR-0407)

1-[4-(5-Methoxy-4-morpholin-4-yl)-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-thiourea (MTR-0408)

4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl)-pyrimidin-2-yl)-2-fluoro-phenyl]-ureido)-benzamide (MTR-0409)

4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl)-pyrimidin-2-yl)-2-fluoro-phenyl]-ureido)-benzene-sulfonamide (MTR-0410)

4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl)-pyrimidin-2-yl)-3-fluoro-phenyl]-ureido)-benzamide (MTR-0411)

4-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl)-pyrimidin-2-yl)-3-fluoro-phenyl]-ureido)-benzene-sulfonamide (MTR-0412)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl)-pyrimidin-2-yl)-phenyl]-3-[4-(4-methanesulfonyl-piperazin-1-yl)-phenyl]-urea (MTR-0413)

N-(5-{3-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl)-pyrimidin-2-yl)-phenyl]-ureido)-pyridin-2-yl)-acetamide (MTR-0415)

2'-[3-(4-Carbamoyl-phenyl)-ureido]-5-ethoxy-6-morpholin-4-yl-[2,5']bipyrimidinyl-4-carboxylic acid ethyl ester (MTR-0417)

2'-[3-(4-Acetylamino-phenyl)-ureido]-5-ethoxy-6-morpholin-4-yl-[2,5']bipyrimidinyl-4-carboxylic acid ethyl ester (MTR-0418)

5-Ethoxy-2-(4-{3-[4-(4-methanesulfonyl-piperazin-1-yl)-phenyl]-ureido)-phenyl)-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0419)

1-[4-(5-Ethoxy-4-morpholin-4-yl)-pyrimidin-2-yl)-phenyl]-3-[4-(4-methanesulfonyl-piperazin-1-yl)-phenyl]-urea (MTR-0420)

1-[4-(5-Ethoxy-4-morpholin-4-yl)-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-carbonyl)-phenyl]-urea (MTR-0421)

1-[4-(5-Ethoxy-4-morpholin-4-yl)-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzooxazol-6-yl)-urea (MTR-0422)

5-Ethoxy-6-morpholin-4-yl-2-{4-[3-(2-oxo-2,3-dihydro-benzooxazol-6-yl)-ureido]-phenyl}-pyrimidine-4-carboxylic acid ethyl ester (MTR-0423)

1-[4-(5-Ethoxy-4-morpholin-4-yl)-pyrimidin-2-yl)-phenyl]-3-(4-methanesulfonyl-phenyl)-urea (MTR-0424)

5-(5-Ethoxy-4-morpholin-4-yl)-pyrimidin-2-yl)-1,3-dihydro-benzimidazol-2-one (MTR-0425)

N-(3-{3-[4-(5-Methoxy-4-morpholin-4-yl)-pyrimidin-2-yl)-phenyl]-ureido)-phenyl)-acetamide (MTR-0426)

N-(3-Dimethylamino-propyl)-4-{3-[4-(5-methoxy-4-morpholin-4-yl)-pyrimidin-2-yl)-phenyl]-ureido)-benzamide (MTR-0427)

225

5-Methoxy-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-ylamine (MTR-0428)

4-[3-(5-Methoxy-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-yl)-ureido]-benzamide (MTR-0429)

1-[4-(4-Methanesulfonyl-piperazin-1-yl)-phenyl]-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0430)

5-Methylsulfanyl-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-ylamine (MTR-0431)

4-[3-(6-Chloro-5-methylsulfanyl-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-yl)-ureido]-benzenesulfonamide (MTR-0432)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzooxazol-6-yl)-urea (MTR-0433)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-sulfonyl)-phenyl]-urea (MTR-0435)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(4-dimethylamino)piperidine-1-carbonyl]-phenyl]-urea (MTR-0436)

4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenylamine (MTR-0437)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea (MTR-0438)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-thiourea (MTR-0440)

N-(5-{3-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-pyridin-2-yl)-acetamide (MTR-0442)

2-{4-[3-(6-Acetylamino-pyridin-3-yl)-ureido]-phenyl}-5-ethoxy-6-morpholin-4-yl-pyrimidine-4-carboxylic acid ethyl ester (MTR-0443)

N-[3-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-methanesulfonamide (MTR-0444)

N-[3-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-2,4-difluoro-benzenesulfonamide (MTR-0445)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0446)

1-[4-(5-Ethoxy-4-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0447)

4-(5-Methoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0448)

N-(4-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-methanesulfonamide (MTR-0449)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzooxazol-6-yl)-urea (MTR-0450)

1-(5-Methoxy-4-morpholin-4-yl-[2,5']bipyrimidinyl-2'-yl)-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0451)

1-[4-(5-Ethoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-sulfonyl)-phenyl]-urea (MTR-0452)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0453)

1-[4-(Morpholine-4-carbonyl)-phenyl]-3-{4-[4-morpholin-4-yl-5-(2-morpholin-4-yl-ethoxy)-pyrimidin-2-yl]-phenyl}-urea (MTR-0454)

1-{4-[4-Morpholin-4-yl-5-(2-morpholin-4-yl-ethoxy)-pyrimidin-2-yl]-phenyl}-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0455)

226

4-(3-{4-[4-Morpholin-4-yl-5-(2-morpholin-4-yl-ethoxy)-pyrimidin-2-yl]-phenyl}-ureido)-benzamide (MTR-0456)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0457)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-(2-oxo-2,3-dihydro-benzooxazol-6-yl)-urea (MTR-0459)

4-(4-Chloro-5-methoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0460)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-carbonyl)-phenyl]-urea (MTR-0461)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0462)

N-(4-{3-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-ureido}-phenyl)-methanesulfonamide (MTR-0463)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0464)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-urea (MTR-0465)

4-(5-Methylsulfanyl-2,6-di-pyridin-3-yl-pyrimidin-4-yl)-morpholine (MTR-0466)

N-[3-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-methanesulfonamide (MTR-0467)

1-[4-(5-Methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-phenyl-thiourea (MTR-0468)

1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0469)

1-(4-Methanesulfonyl-phenyl)-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0470)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-sulfonyl)-phenyl]-urea (MTR-0471)

1-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0472)

1-[4-(5-Methoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(morpholine-4-carbonyl)-phenyl]-urea (MTR-0473)

1-[4-(5-Methoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0474)

1-[4-(4-Chloro-5-methoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0476)

1-Ethyl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0477)

1-Methyl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0478)

1-Isopropyl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0479)

1-[4-(5-Methoxy-4-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0480)

1-{4-[4-Morpholin-4-yl-5-(2-morpholin-4-yl-ethoxy)-pyrimidin-2-yl]-phenyl}-3-pyridin-4-yl-urea (MTR-0481)

4-(5-Methoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenylamine (MTR-0482)

4-(5-Methoxy-4-morpholin-4-yl-6-(4-aminophenyl)-pyrimidin-2-yl)-phenylamine (MTR-0483)


1-[4-(4-Chloro-5-methylsulfanyl-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-isoxazol-3-yl-urea (MTR-0484)

227


- 1-Isloxazol-3-yl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0485)
- N-(5-{3-[4-(5-Methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-ureido}-pyridin-2-yl)-acetamide (MTR-0486)
- 1-[4-(4-Methanesulfonyl-piperazin-1-yl)-phenyl]-3-[4-(5-methoxy-4-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0487)
- 1-[4-(4-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0488)
- 1-[4-(4-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0489)
- 1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0490)
- 1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0491)
- 1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0492)
- 1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0493)
- 1-[4-(4-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0494)
- 1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0495)
- 1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0496)
- 1-[4-(5-Methoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0497)
- 1-[4-(5-Methoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-4-yl-urea (MTR-0498)
- 1-[4-(5-Methoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-pyridin-3-yl-urea (MTR-0499)
- 1-[4-(4-Chloro-5-methoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0500)
- 1-[4-(5-Methoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0501)
- 1-[4-(5-Methoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-[4-(3-oxo-morpholin-4-yl)-phenyl]-urea (MTR-0502)
- 1-[4-(4-Chloro-5-methoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-isoxazol-3-yl-urea (MTR-0503)
- 1-Isloxazol-3-yl-3-[4-(5-methoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0504)
- 1-[4-(4-Chloro-5-ethoxy-6-morpholin-4-yl-pyrimidin-2-yl)-phenyl]-3-isoxazol-3-yl-urea (MTR-0505)

228

- 1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-4-yl-pyrimidin-2-yl)-phenyl]-3-isoxazol-3-yl-urea (MTR-0506)
- 1-Isloxazol-3-yl-3-[4-(5-methoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0507)
- 1-[4-(5-Ethoxy-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-3-isoxazol-3-yl-urea (MTR-0508) and
- 1-Isloxazol-3-yl-3-[4-(5-methylsulfanyl-4-morpholin-4-yl-6-pyridin-3-yl-pyrimidin-2-yl)-phenyl]-urea (MTR-0509)
- or a pharmaceutically acceptable salt thereof.
8. A pharmaceutical composition which comprises an effective amount of a compound of claim 1 and a pharmaceutically acceptable carrier or diluent.
9. A pharmaceutical composition which comprises an effective amount of a compound of claim 7 and a pharmaceutically acceptable carrier or diluent.
10. A process for preparing the compound of formula (I) according to claim 1, comprising:
- (i) reacting a compound of formula (7)



wherein Z' is halo, and Z, R², and Y are as defined in claim 1, with an amine of formula HW, wherein W is defined in claim 1, and in the presence of an inert solvent and a base to form a compound of formula (8),



- and
- ii) subjecting the compound of formula (8) to Suzuki coupling in the presence of a boronic acid or a boronic ester to form a compound of formula (I).

* * * * *